

**Priest Rapids Hatchery Monitoring and Evaluation
Annual Report for 2020 - 2021**

Steven P. Richards
Washington Department of Fish & Wildlife

And

Todd N. Pearsons
Public Utility District Number 2 of Grant County, Washington

November 30, 2021

Disclaimer

This report is provided as an annual data update of the Public Utility District No. 2 of Grant County, Washington's (Grant PUD's) monitoring and evaluation plan for Priest Rapids Hatchery. All data are provisional and subject to change as new data and analyses become available. Readers are cautioned to use data at their own risk and should consult the most current report to obtain the most current and accurate information. Data sets will become final when they are published in peer reviewed scientific journals.

This report should be cited as:

Richards, S.P. and T.N. Pearsons. 2021. Priest Rapids Hatchery Monitoring and Evaluation Annual Report for 2020-2021. Public Utility District No. 2 of Grant County, Ephrata, Washington.

Executive Summary

This report is the eleventh annual report dedicated to monitoring and evaluating the Priest Rapids Hatchery (PRH) production of fall Chinook Salmon. The PRH is located below Priest Rapids Dam adjacent to the Columbia River and has been in operation since 1963. The monitoring and evaluation program associated with PRH is intended to evaluate the performance of the program in meeting hatchery and natural production goals. This report is intended to be cumulative, but also focus attention on the most recent year of data collection and production (2020-2021).

The PRH was originally built to mitigate for the construction and operation of Priest Rapids and Wanapum dams. The hatchery is operated as an integrated program for the purpose of increasing harvest while limiting undesirable risks to the naturally spawning population. The hatchery produces 5.6 million subyearling fall Chinook Salmon for Public Utility District No. 2 of Grant County, Washington's (GPUD) mitigation requirement and 1.7 million subyearling fall Chinook Salmon under contract with the United States Army Corps of Engineers for mitigation for the construction and operation of John Day Dam. These fish contribute significantly to a variety of fisheries, such as fisheries off the coasts of Alaska and Canada and fisheries in the Columbia River.

The estimated total escapement of fall Chinook Salmon to the Hanford Reach in 2020 was 74,834 fish. This is higher than the mean abundances of the past few decades. The mean and median escapement for 1991 through 2020 was 73,879 and 58,779 fish, respectively.

The 2020 returns to PRH volunteer trap totaled 38,333 fall Chinook Salmon. A total of 9,219 fish that returned to the volunteer trap at PRH were ponded at the hatchery for broodstock of which many were later surplus or spawned on-site by staff from the Yakama Nation for their off-site production. An additional 1,166 fish from the Angler Broodstock Collection (ABC) fishery were included in the broodstock in an effort to increase the number of natural origin broodstock. In total, 5,606 fish were spawned to meet egg-take goals for multiple hatchery programs. Greater than 68% of the fish that were surplus to broodstock needs in recent years were provided to foodbanks and tribes for consumption.

There were several similarities and differences of hatchery and natural origin fall Chinook Salmon. The hatchery origin fish appeared to return at a younger age than natural origin fish. It appears that age-2 hatchery origin fish tend to be larger than natural origin fish of the same age. Likewise, age-4 and 5 natural origin fish tend to be larger than their hatchery origin counterparts. The number of eggs, egg size, and egg mass produced by hatchery and natural origin females increased with fish length and were generally similar when standardized for length. Except for one year (2013), egg retention in female carcasses in the Hanford Reach has been low.

Hatchery origin fish released from PRH spawned throughout the Hanford Reach. In addition, the hatchery origin proportions of spawners relative to total spawners in the different sections of the Hanford Reach were similar. Recent evidence suggests that adult carcasses drift downstream of their spawning location and bias estimates of downstream spawning distribution. Stray rates into other populations appeared to be low based upon coded-wire tag recoveries and PIT tag detections of PRH adults in the Snake River were also low. However, in some years there have been notable numbers of PIT tag detections of PRH adults above Priest Rapids Dam.

The PRH continued to contribute substantially to ocean and Columbia River fisheries and tends to have higher adult recruitment rates than the natural spawning fall Chinook Salmon to the

Hanford Reach of the Columbia River. Adult recruitment rate of the latest complete brood year (2014) for PRH was 6.79 versus 0.38 for fish spawning in the Hanford Reach.

PRH origin fish were estimated to make up 22.0% of the natural spawning population in the Hanford Reach during 2020. All hatchery fish combined (including fish released from Ringold Hatchery and strays from outside the Hanford Reach) comprised 26.8% of the fall Chinook Salmon on the spawning grounds. Otolith recoveries at the PRH volunteer trap indicated that a very high percentage of fish returning to the PRH were of PRH origin. The proportionate natural influence (PNI) for Hanford Reach fall Chinook Salmon including all hatcheries was 0.546 in 2020. This value was calculated using a gene flow model based on the Ford model and was lower than the PNI target of 0.67 for the first time in seven consecutive years. Adult management of fish at the PRH volunteer trap and alternative broodstock collection techniques to increase natural origin fish in the broodstock have contributed to improvements in PNI for the PRH program, but high numbers of hatchery origin spawners can make management of PNI challenging when natural return abundance is relatively low.

Table of Contents

1.0 Introduction..... 1

2.0 Objectives 3

3.0 Project Coordination 3

4.0 Life History – Hanford Reach Fall Chinook Salmon 4

5.0 Sample Size Considerations..... 4

6.0 Current Operation at Priest Rapids Hatchery..... 5

7.0 Origin of Adult Returns to Priest Rapids Hatchery 7

 7.1 Origin Based on Otolith Marks..... 8

 7.2 Origin Based on Coded-Wire Tag Recoveries..... 10

8.0 Broodstock Collection and Sampling 11

 8.1 Broodstock Age Composition..... 11

 8.2 Length by Age Class of Broodstock 13

 8.3 Gender Ratios..... 14

 8.4 Fecundity..... 15

9.0 Hatchery Rearing 20

 9.1 Number of Eggs Taken 20

 9.2 Number of Acclimation Days 20

 9.3 Annual Releases, Tagging, and Marking..... 21

 9.4 Fish Size and Condition of Release 23

 9.5 Survival Estimates 24

 9.6 Juvenile PIT Tag Detections at the Priest Rapids Hatchery Array 25

10.0 Adult Fish Pathogen Monitoring 28

 10.1 Juvenile Fish Health Inspections 30

11.0 Redd Survey..... 31

 11.1 Hanford Reach Aerial Redd Counts 31

 11.2 Redd Distribution..... 32

 11.3 Spawn Timing..... 33

 11.4 Escapement 33

 11.5 Hatchery Discharge Channel Redd Counts..... 35

12.0 Carcass Surveys 35

 12.1 Hanford Reach Carcass Survey: Section 1 – 5 36

 12.2 Proportion of Escapement Sampled: Section 1-5 37

12.3	Carcass Distribution and Origin.....	38
12.4	Priest Rapids Dam Pool Carcass Survey: Section 6	41
12.5	Number Sampled: Section 6	41
	12.5.1 Proportion of Escapement Sampled: Section 6.....	41
	12.5.2 Carcass Origin: Section 6.....	42
12.6	Hatchery Discharge Channel: Section 7 and 8 Carcass Survey	43
12.7	Number sampled: Section 7 and 8	43
	12.7.1 Proportion of Escapement Sampled: Section 7 and 8.....	44
	12.7.2 Carcass Distribution and Origin: Section 7 and 8.....	44
13.0	Life History Monitoring.....	45
	13.1 Adult Migration Timing.....	46
	13.2 Age at Maturity	48
	13.3 Size at Maturity.....	51
	13.4 Gender Composition for Adult Escapement	55
	13.5 Egg Retention.....	56
14.0	Contribution to Fisheries.....	58
15.0	Straying.....	60
16.0	Genetics.....	64
17.0	Proportionate of Natural Influence	64
	17.1 Estimate of pNOB.....	65
	17.2 Estimates of pHOS.....	66
	17.3 Estimates of PNI	67
18.0	Natural and Hatchery Replacement Rates	69
19.0	Smolt-to-Adult Survivals	70
20.0	ESA/HCP Compliance.....	72
	20.1 Broodstock Collection	72
	20.2 Hatchery Rearing and Release	72
	20.3 Hatchery Effluent Monitoring.....	73
	20.4 Ecological Risk Assessment	73
21.0	Distribution of Adult fall Chinook Salmon carcass from Priest Rapids Hatchery	73
22.0	Acknowledgments.....	74
23.0	Literature Cited	75

List of Figures

Figure 1. Location of Priest Rapids and Ringold Spring hatcheries and the city of Richland Washington (indicated by stars)..... 2

Figure 2. Priest Rapids Hatchery facility and Priest Rapids Dam Off-Ladder Adult Fish Trap. 2

Figure 3 Proportion of annual returns by week (Sunday through Saturday) adult Chinook Salmon collected at the Priest Rapids Hatchery Volunteer Trap. First week generally begins between September 7 and 15. 6

Figure 4 Linear relationship between fecundity and fork length for combined samples of natural and hatchery origin fall Chinook Salmon spawned at Priest Rapids Hatchery, Return Years 2010-2020. 17

Figure 5 Fecundity versus fork length for natural- and hatchery- origin fall Chinook Salmon sub-sampled at Priest Rapids Hatchery, Return Years 2013-2020..... 18

Figure 6 Mean skein weight versus fork length for natural- and hatchery- origin fall Chinook Salmon sub-sampled at Priest Rapids Hatchery, Return Years 2013-2020..... 19

Figure 7 Total egg mass weight versus fork length for natural- and hatchery- origin fall Chinook Salmon sub-sampled at Priest Rapids Hatchery, Return Years 2013-2020..... 19

Figure 8 Distribution of fall Chinook Salmon redd counts by location for the 2020 aerial surveys in the Hanford Reach, Columbia River. (Data provided by Mission Support Alliance). 33

Figure 9 Location of aerial redd index areas (green area numbers) and river boat carcass survey sections in the Hanford Reach..... 36

List of Tables

Table 1 Source and disposition of Chinook Salmon collected for Grant PUD and USACE broodstock at Priest Rapids Hatchery, Return Year 2020. ABC = Angler broodstock collection. 6

Table 2 Total fish handled, numbers sampled, and estimates of hatchery and natural origin Chinook Salmon collected at Priest Rapids Hatchery, Priest Rapids Dam Off-Ladder Adult Fish Trap, and Angler Broodstock Collection fishery. Origin determined by otolith thermal marks, presence of coded-wire tags, and/or adipose clips, Return Years 2013-2020. 9

Table 3 Estimated proportion of hatchery and natural origin adult Chinook Salmon returning to the Priest Rapids Hatchery Volunteer Trap based on coded-wire tag expansion. The entire collection was sampled for coded-wire tags, Return Years 2005-2020. 10

Table 4 Age composition for hatchery and natural origin fall Chinook Salmon spawned at Priest Rapids Hatchery (includes all sources of broodstock), Return Years 2007-

	2020. Proportions calculated from expanded age compositions by origin for each source of broodstock to account for differing sample rates.....	11
Table 5	Age composition for hatchery and natural origin fall Chinook broodstock collected from the Priest Rapids Hatchery volunteer trap, Return Years 2012-2020.....	12
Table 6	Age composition for hatchery and natural origin fall Chinook Salmon broodstock collected from Angler Broodstock Collection, Return Years 2012-2020.....	13
Table 7	Mean fork length (cm) at age (total age) of fall Chinook Salmon sampled from each source of broodstock spawned at Priest Rapids Hatchery, Return Year 2020. N = sample size and SD = 1 standard deviation.	13
Table 8	Mean fork length (cm) at age (total age) of hatchery and natural origin fall Chinook Salmon collected from broodstock originating from the Priest Rapids Hatchery volunteer trap. N = sample size and SD = 1 standard deviation, Return Years 2012-2020.....	14
Table 9	Number of male and female hatchery fall Chinook Salmon broodstock at Priest Rapids Hatchery, Return Years 2001-2020. Ratios of males to females are also provided.	14
Table 10	Mean fecundity of fall Chinook Salmon collected for broodstock at Priest Rapids Hatchery, Return Years 2001-2020. SD = 1 standard deviation.....	15
Table 11	Mean fecundity at age for fall Chinook Salmon sampled at the Priest Rapids Hatchery, Return Years 2010-2020. N = sample size and SD = 1 standard deviation.....	17
Table 12	Number of eggs taken from fall Chinook Salmon broodstock collected at Priest Rapids Hatchery, Return Years 1984-2020.	20
Table 13	Number of days fall Chinook Salmon fry were reared at Priest Rapids Hatchery prior to release, Brood Year 2020.....	21
Table 14	Number of marked, unmarked, and tagged fall Chinook Salmon smolts released from Priest Rapids Hatchery, Brood Years 1977-2020.	22
Table 15	Mean length (FL, mm), weight (g and fish/pound), and coefficient of variations (CV) of fall Chinook smolts released from Priest Rapids Hatchery, Brood Years 1991-2020.	23
Table 16	Hatchery life stage survival (Proportion) for fall Chinook Salmon at Priest Rapids Hatchery, Brood Years 1989-2020.	24
Table 17	Number of sub-yearlings PIT tagged, mark, and release dates, and the number of unique tags detected at the array in the Priest Rapids discharge channel, Brood Years 2011-2018 and 2020.	26
Table 18	Viral inspections of fall Chinook Salmon broodstock at Priest Rapids Hatchery, Return Years 1991-2020.	28

Table 19	ELISA test results to determine risk of bacterial kidney disease of adult female fall Chinook Salmon broodstock at Priest Rapids Hatchery, Return Years 2008-2020.....	29
Table 20	Juvenile fish health inspections for Priest Rapids Hatchery fall Chinook Salmon, Brood Years 2006-2020.....	30
Table 21	Summary of fall Chinook Salmon peak redd counts for the 1948-2020 aerial surveys in the Hanford Reach, Columbia River.	32
Table 22	Number of fall Chinook Salmon redds counted in different reaches on the Hanford Reach area of the Columbia River during aerial counts, October 2020 through November 2020. (Data provided by Mission Support Alliance).....	32
Table 23	Calculation of escapement estimates for fall Chinook Salmon in the Hanford Reach, Columbia River 2020.....	34
Table 24	Escapement for fall Chinook Salmon in the Hanford Reach, Return Years 1991-2020.....	34
Table 25	Numbers and percentage of total escapement of fall Chinook Salmon carcasses surveyed (1:1 rate) for coded-wire tags within each survey section on the Hanford Reach, Return Years, 2010-2020.....	36
Table 26	Number of carcass surveys conducted by section in the Hanford Reach, Return Years 2010-2020.....	37
Table 27	Number of redds and carcasses, total spawning escapement, and proportion of escapement sampled for fall Chinook Salmon in Section 1 through 5 of the Hanford Reach, Return Year 2020.....	38
Table 28	Numbers of natural and hatchery origin fall Chinook Salmon carcasses sampled within Section 1 through 5 of Hanford Reach based on expansions of coded-wire tag recoveries, Return Years 2010-2020.....	39
Table 29	Origin of Chinook Salmon carcasses recovered in the Hanford Reach by section based on recoveries of marked and unmarked carcasses within the biological sample, Return Years 2012-2020.....	40
Table 30	Carcasses sampled, total spawning escapement and proportion of escapement for fall Chinook Salmon in Section 6 (Priest Rapids Dam Pool), Return Years 2010-2020.....	41
Table 31	Origin of fall Chinook Salmon spawning in Section 6 (Priest Rapids Dam Pool), Return Years 2012-2020.....	42
Table 32	The number of fall Chinook Salmon carcass surveys within Section 7 (Priest Rapids Hatchery Discharge Channel) and Section 8 (Columbia River at the confluence of the hatchery discharge channel), Return Years 2010-2020.	43
Table 33	Number of carcasses sampled, total spawning escapement and proportion of escapement sampled for fall Chinook Salmon within Section 7 (Priest Rapids Hatchery Discharge Channel) and Section 8 (Columbia River at confluence of the hatchery discharge channel), Return Year 2020.....	44

Table 34	The origin of Chinook Salmon carcasses recovered within Section 7 (Priest Rapids Hatchery Discharge Channel) and Section 8 (Columbia River at the confluence of the hatchery discharge channel), Return Years 2012-2020.	44
Table 35	The week that 10%, 50% (median), and 90% of the natural and hatchery origin fall Chinook Salmon passed Bonneville Dam, 2010-2020. Migration timing is based on PIT tag passage of Hanford natural origin and Priest Rapids Hatchery in the adult fish ladder at Bonneville Dam.	47
Table 36	Age compositions for fall Chinook Salmon sampled in the Hanford Reach escapement compared to fall Chinook Salmon sampled at Priest Rapids Hatchery (genders combined), Brood Years 1998-2015.	49
Table 37	Age compositions for male natural and hatchery origin fall Chinook Salmon sampled in the Hanford Reach escapement, Brood Years 2007-2015.	50
Table 38	Age compositions for female natural and hatchery origin fall Chinook Salmon sampled in the Hanford Reach escapement, Brood Years 2007-2015.	50
Table 39	Age compositions for natural and hatchery origin fall Chinook Salmon sampled in the Hanford Reach escapement, Brood Years 2007-2015.	51
Table 40	Mean fork length (cm) at age (total age) of fall Chinook Salmon sampled in the Hanford Reach escapement compared to fall Chinook Salmon sampled at Priest Rapids Hatchery, Brood Years 1999-2015. N = sample size and SD = 1 standard deviation.	52
Table 41	Mean fork length (cm) at age (total age) of male natural and hatchery origin fall Chinook Salmon that spawned naturally in the Hanford Reach, Brood Years 2007-2015. N = sample size and SD = 1 standard deviation.	53
Table 42	Mean fork length (cm) at age (total age) of female natural and hatchery origin fall Chinook Salmon that spawned naturally in the Hanford Reach, Brood Years 2007-2015. N = sample size and SD = 1 standard deviation.	53
Table 43	Mean fork length (cm) at age (total age) of natural and hatchery origin fall Chinook Salmon that spawned naturally in the Hanford Reach, Brood Years 2007-2015. N = sample size and SD = 1 standard deviation.	54
Table 44	Comparisons of male to female ratio of fall Chinook Salmon sampled at Priest Rapids Hatchery and in the Hanford Reach stream surveys, Brood Years 1996-2015.	55
Table 45	Comparisons of male to female ratio of fall Chinook Salmon sampled in the Hanford Reach stream surveys, Brood Years 2007-2015.	56
Table 46	Comparison of egg retention of natural and hatchery origin fall Chinook sampled in the Hanford Reach stream survey, Return Years 2010-2020.	57
Table 47	Hatchery fall Chinook Salmon contributions to harvest in the Hanford Reach fall Chinook Salmon fishery. Coded-wire tag recoveries provided from RMIS database were expanded by sample rate and juvenile tag rate, Return Years 2003-2020.	59

Table 48	Priest Rapids Hatchery coded-wire tag recoveries provided from RMIS by brood year and harvest type expanded by sample rate and juvenile tag rate, Brood Years 1997-2014. Data only includes coded-wire tag recoveries from adipose clipped fish expanded by the juvenile tag rate.....	60
Table 49	Percent of fall/summer Chinook spawning populations by return year (2000-2019) comprised of Priest Rapids Hatchery fall Chinook from 1998-2016 brood releases based on coded-wire tag recoveries.	61
Table 50	Estimated number and percentage of Priest Rapids Hatchery fall Chinook Salmon spawning escapement to Priest Rapids Hatchery and stream within and outside of the presumptive target stream by brood year (1992-2014). Coded-wire tag recoveries are expanded by juvenile mark rate and survey sample rate for each brood year.	62
Table 51	Last observations of unique PIT tagged adult fall Chinook from Priest Rapids Hatchery at detection sties upstream of McNary Dam, Brood Years 1999-2016.	63
Table 52	Origin of broodstock and pNOB apportioned to program for fall Chinook Salmon spawned at Priest Rapids Hatchery, Brood Year 2020.	65
Table 53	Number of broodstock spawned and pNOB apportioned to program for fall Chinook Salmon spawned at Priest Rapids Hatchery, Return Years 2012-2020.	66
Table 54	Proportion of hatchery Chinook Salmon on the spawning grounds (pHOS) in the Hanford Reach, Brood Years 2012-2020.....	66
Table 55	Origin of pHOS apportioned by program source for fall Chinook Salmon spawning naturally in the Hanford Reach, Return Years 2012-2020.	67
Table 56	PNI of the Hanford Reach fall Chinook Salmon supplementation program based on expanded coded-wire tag recoveries of all fish surveyed, Return Years 2001-2020.....	68
Table 57	PNI estimates for the Hanford Reach fall Chinook Salmon supplementation programs based on otoliths, Return Years 2012-2020. Calculated from multiple population gene flow model based on the Ford model which has been extended to three or more populations.	69
Table 58	Broodstock spawned at Priest Rapids Hatchery, estimated escapement to the Hanford Reach, natural and hatchery origin recruits (NOR and HOR), and natural and hatchery replacement rates (NRR and HRR, with and without harvest) for natural origin fall Chinook Salmon in the Hanford Reach, Brood Years 1996-2013.....	70
Table 59	Smolt-to-adult Survival Ratios (SAR) for Priest Rapids Hatchery fall Chinook Salmon, Brood Years 1992-2014. Data includes all coded-wire tag recoveries from adipose clipped fish.....	71
Table 60	Smolt-to-adult Survival Ratios (SAR) for Hanford Reach natural origin fall Chinook Salmon, Brood Years 1992-2014. Data includes all coded-wire tag recoveries from adipose clipped fish. Source Regional Mark Processing Center.	71

Table 61	Recoveries and disposition of steelhead at the Priest Rapids Hatchery volunteer trap, Return Year 2020.....	72
Table 62	Disposition of Chinook Salmon removed from Priest Rapids Hatchery volunteer trap, Return Year 2001-2020. Surplus numbers do not include shipped to other hatchery programs.....	73

List of Appendices

Appendix A	Evaluation of Coded-Wire Tag Bias	A-1
Appendix B	Recovery of coded-wire tags collected from adult returns to the Priest Rapids Hatchery Volunteer Trap during Return Year 2020	B-1
Appendix C	Juvenile fish health inspections for Priest Rapids Hatchery fall Chinook Salmon, Brood Years 1998-2020. The description in the Condition column indicates the presence of a certain condition within at least one of the fish examined.	C-1
Appendix D	Number and percent of fall Chinook Salmon redds counted in different reaches of the Columbia River, 2001-2020. Data for years 2001-2010 was collected by staff with Pacific Northwest National Laboratory. Data for years 2011-2020 was collected by staff with Environmental Assessment Services, LLC.	D-1
Appendix E	Historical numbers of Chinook Salmon carcasses recovered during the annual Hanford Reach fall Chinook Salmon carcass survey, Return Years 1991-2020.	E-1
Appendix F	Estimated escapements for fall Chinook spawning in Hanford Reach and Priest Rapids Dam pool, Return Year 2020.	F-1
Appendix G	Demographic comparisons for double index tag groups released from Priest Rapids Hatchery, Brood Years 2009-2017.	G-1
Appendix H	Explanation of methods for calculating adult-to-adult expansions based on coded-wire tag recoveries at Priest Rapids Hatchery	H-1

1.0 Introduction

The Public Utility District No. 2 of Grant County, Washington (Grant PUD) produces and releases 5.6 million subyearling fall Chinook Salmon smolts from Priest Rapids Hatchery (PRH) as part of its mitigation for the construction and operation of Priest Rapids and Wanapum dams. The mitigation is the result of three components: 1) inundation of historic spawning habitat (5,000,000), annual losses of fish that migrate through the project (325,543), and flow fluctuation impacts in the Hanford Reach (273,961). The PRH is located at the top end of the Hanford Reach on the east bank of the Columbia River immediately downstream of Priest Rapids Dam (Figure 1 and Figure 2). The Washington Department of Fish & Wildlife (WDFW) operates PRH which is owned by the Grant PUD. Funding for operations and maintenance is provided by both the Grant PUD and the U.S. Army Corps of Engineers (USACE). This report describes the monitoring and evaluation of the PRH M&E program.

PRH also produces fish for other programs. PRH produces and releases 1.7 million subyearling smolts on-site for the USACE John Day Mitigation. An additional 4.1 million eyed eggs are targeted to provide fish for the USACE John Day Mitigation released at Ringold Springs Hatchery (RSH) located at the lower portion of the Hanford Reach. The eggs for the RSH program are first transferred to Bonneville Hatchery for marking and ultimately ~3.7 million subyearlings are transported to, acclimated, and released as subyearling smolts from RSH. A separate companion report of the USACE program is provided by the USACE. In recent years, PRH has accommodated egg-takes for fall Chinook Salmon programs managed by either Yakama Nation (YN) or Umatilla Tribe as well as the WDFW's Salmon in the Classroom program and to support various research projects.

A Monitoring and Evaluation Plan for all Grant, Douglas, and Chelan County Public Utility Districts Hatchery Programs has been updated and approved by the committees that oversee the PUD hatchery programs (Hillman et al. 2017). This document provides guiding principles and approaches for the monitoring and evaluation (M&E) of all PUD hatchery programs including PRH. Objectives, hypotheses, measured and derived variables, and field methods that were used to collect data are described in this document.

This report of the PRH M&E program is the eleventh annual report (Hoffarth and Pearsons 2012a, 2012b, Richards et al. 2013, Richards and Pearsons 2014, 2015, 2016, 2017, 2018, 2019 and 2020) and encompasses data collected during the Washington State fiscal year (FY) 2020 - 2021 as well as earlier years where data were available. The data presented in this report are preliminary and subject to change as new data and analyses become available. Readers are encouraged to consult the most recent annual report to obtain the most current and accurate information.




Figure 1. Location of Priest Rapids and Ringold Spring hatcheries and the city of Richland Washington (indicated by stars).



Figure 2. Priest Rapids Hatchery facility and Priest Rapids Dam Off-Ladder Adult Fish Trap.

2.0 Objectives

The objective of the PRH M&E plan is to evaluate the performance of the PRH program relative to the goals and objectives of the PRH program. The overarching goal of the PRH program is to meet Grant PUD's hatchery mitigation by producing fish for harvest while keeping genetic and ecological impacts within acceptable limits. The M&E objectives of the PRH program are described below.

- Objective 1: Determine if conservation programs have increased the number of naturally spawning and naturally produced adults of the target population and if the program has reduced the natural replacement rate (NRR) of the supplemented population.
- Objective 2: Determine if the proportion of hatchery fish on the spawning ground affects the freshwater productivity of supplemented stocks.
- Objective 3: Determine if the hatchery adult-to-adult survival (i.e., hatchery replacement rate, HRR) is greater than the natural adult-to-adult survival (i.e., natural replacement rate, NRR) and the target hatchery survival rate.
- Objective 4: Determine if the proportion of hatchery origin spawners (pHOS or PNI) is meeting management targets.
- Objective 5: Determine if the run timing, spawn timing, and spawning distribution of the hatchery component is similar to the natural component of the target population or is meeting programs-specific objectives.
- Objective 6: Determine if stray rate of hatchery fish is below the acceptable levels to maintain genetic variation among stocks.
- Objective 7: Determine if genetic diversity, population structure, and effective population size have changed in natural spawning populations as a result of the hatchery program.
- Objective 8: Determine if hatchery programs have caused changes in phenotypic characteristics of natural populations.
- Objective 9: Determine if hatchery fish were released at programmed size and number.
- Objective 10: Determine if appropriate harvest rates have been applied to the conservation, safety-net, and segregated harvest programs to meet the HCP/SSSA goal of providing harvest opportunities while also contributing to population management and minimalizing risk to natural populations.

We also present information in this report about two regional objectives that relate to disease and ecological interactions.

3.0 Project Coordination

WDFW M&E staff partially assigned to PRH also conducts similar work at RSH. The M&E staff also works in conjunction with multiple WDFW groups that include PRH fish culture staff, the Columbia River Coded-Wire Tag Recovery Program (CRCWTRP), Region 3 Fish Management staff, the Supplementation Research Team in Wenatchee, and the Grant PUD biological science staff to complete many of the tasks included in the M&E Plan. In addition, samples collected at the hatchery and in the field were transported to and analyzed by WDFW laboratories including the WDFW Scale Reading Lab and the WDFW Otolith Lab. Coded-wire tags (CWT) were

processed by WDFW staff. Data and analyses collected in association with the PRH M&E and Hanford Reach population monitoring are incorporated into the WDFW Traps, Weirs, and Surveys (TWS) database which is administered by the WDFW staff stationed in the Region 5 Headquarters in Vancouver. Agency managers use these data for forecasting and managing fall Chinook Salmon populations in the Columbia and Snake rivers and tributaries. WDFW and Grant PUD secured and held all environmental permits necessary for the work described in this report.

4.0 Life History – Hanford Reach Fall Chinook Salmon

The Hanford Reach is one of the last non-impounded reaches of the Columbia River and the location of the largest and most productive natural spawning fall Chinook Salmon population in the United States (Harnish et al. 2012, Langshaw et al. 2015, Harnish 2017, Langshaw et al. 2017). The Hanford Reach extends ~81 km from the city of Richland to the base of Priest Rapids Dam. Natural origin fall Chinook Salmon emerge from the substrate in the spring and rear in the Hanford Reach until outmigration in the summer. Egg-to-fry survival has been estimated to be about 71% in the Hanford Reach (Oldenburg et al. 2012) and egg-to-pre-smolt survival has been estimated to be about 40.2% (Harnish et al. 2012) or more recently at 63.4% (Harnish, 2017). Both estimates are high when compared to other Chinook Salmon populations and flow management within the Hanford Reach has resulted in improvements in survival (Harnish et al. 2012, Harnish 2017, Langshaw et al. 2017).

The age at maturity for naturally produced fish in the Hanford Reach varies between age-1 mini-jack and age-6 adults: albeit recoveries of age-1 and 6 fish are generally rare. The age of fish reported in this document begins with the first birthday occurring the year after the parents spawned. The abundance of mini-jacks which mature as age-1 males is currently not known. Age-2 male fall Chinook Salmon (a.k.a jacks) return to the Hanford Reach after spending roughly one year in the ocean. The majority of the natural origin adults return after spending three to four years in the ocean (age-4 and 5). A small portion, typically less than 2%, will spend up to five years in the ocean and return as age-6. The ocean distribution of natural and hatchery origin Hanford Reach upriver brights are similar and range from the northern California coast to the Gulf of Alaska (Norris et al. 2000, Weitkamp 2010). The majority of the adults migrate north of the Columbia River with the harvest primarily occurring in non-selective ocean and freshwater fisheries (Norris et al, 2000). Adults return to the mouth of the Columbia River between August and October and spawn in large cobble substrate between October and December (Langshaw et al. 2017).

5.0 Sample Size Considerations

We attempted to strike an appropriate balance between objectives, statistical precision, logistics, and financial investment when setting sample size targets. A variety of approaches were used for setting sample sizes and this depended upon the objective. For example, a phased subsampling approach was used in some cases to determine age and origin and 100% sampling occurred in others (e.g., CWT, otoliths in fecundity samples). A phased approach was used to collect some biological samples with sufficient accuracy and precision. In the phased approach, we attempted to collect an excess number of raw samples such as carcasses and trap recoveries and then use post season analysis to determine sub-sampling strategies for otolith decoding where appropriate. The sample size target of systematic field sampling for later otolith reading was 2,500 of the carcasses in the Hanford Reach, 1,000 at the hatchery trap, and 1,000 of the hatchery volunteer

broodstock, and 200 broodstock collected from each other source such as off ladder adult fish trap at Priest Rapids Dam (OLAFT) and angler broodstock collection fishery (ABC).

All adult fall Chinook Salmon recovered at PRH, in the Hanford Reach sport fishery, and in the stream surveys were screened for the presence of CWT to increase the number of CWT recoveries and maximize the precision of estimates generated from these data. Representative otolith samples by survey type were randomly selected as a sub-sample for decoding to estimate origin by age class if numbers allowed. In some cases, all otolith samples for a survey type were processed if the sampling rate provided relatively low numbers of otoliths collected or if there was a need for higher precision or accuracy. During return year 2020, randomly selected sub-samples of otoliths collected from the PRH volunteer trap and volunteer broodstock were submitted for decoding. The methodologies for selecting otolith sub-samples have differed between return years as field methods changed and as new analyses facilitated improvements in approaches. In general, we randomly select otoliths from various survey types to obtain roughly 120 otoliths for each age and gender. In some cases, all otoliths were submitted for stratified groups (age/gender) when specific age classes contain less than 100 samples. For example, typically all samples of age-5 and 6 fish were submitted because of the low number of fish represented in the field collected sample. The stratified sub-sample size refinement process is described in Richards and Pearsons (2014). The sub-sample groups often included fish possessing a CWT within the biological sample which increased the number of fish sampled for origin with no additional cost.

6.0 Current Operation at Priest Rapids Hatchery

The 2020 broodstock for PRH were collected at the hatchery volunteer trap, and from the ABC fishery. The majority of the broodstock were collected from the PRH volunteer trap which was operated from September 14 through December 14, 2020. A total of 39,999 mature fall Chinook Salmon were handled during broodstock collection activities (Table 1). To increase pNOB for the Grant PUD program, to the extent possible, the broodstock ponded excluded adipose intact fish with a fork length less than 74 cm and known hatchery origin fish (i.e., possessing an adipose clip and or CWT). Data collected at PRH in prior years suggests that age 3 adults recovered from the volunteer trap included lower proportions of natural origin fish than age 4 and 5 adults. Prioritizing these fish for the Grant PUD program results in some known hatchery origin fish included as broodstock for the USACE program due to a lack of sufficient numbers of adipose fin intact/non CWT fish to meet the broodstock needs for both programs. A portion of these known hatchery origin fish ponded were surplus as they were replaced by adipose fin intact/non CWT fish during subsequent trapping and ponding operations. In total, 5,606 adults were spawned for the Grant PUD and USACE programs. Of these, 1,709 (509 males and 1,200 females) were adipose clipped.

A portion of the fish intended for surplus from PRH were utilized for broodstock to support other fall Chinook Salmon production by the Yakama Nation. These fish included 263 males and 241 females spawned on-site by staff from the Yakama Nation. The PRH monitoring and evaluation (M&E) staff categorized and sampled these fish as surplus from PRH. The carcasses used for broodstock were utilized for pet food since they were treated with formalin during the period in which they were held for broodstock.

Table 1 Source and disposition of Chinook Salmon collected for Grant PUD and USACE broodstock at Priest Rapids Hatchery, Return Year 2020. ABC = Angler broodstock collection.

Collection Location	Gender	Collected	Trap Surplused	Trap Mortalities	Ponded	Spawned ¹	Pond Surplused	Pond Mortalities
Volunteer Trap (Sept 10 – Dec 6)	Males ¹	24,326	21,648	138	2,540	1,311	745	484
	Females	10,368	4,821	49	5,498	3,287	1,344	867
	Jacks	4,139	3,946	178	15	1	9	5
	Total	38,833	30,415	365	8,053	4,599	2,098	1,356
ABC (Oct 25, 26 & 27)	Males	626	0	0	626	566	17	43
	Females	538	0	0	538	441	28	72
	Jacks	2	0	0	2	0	2	0
	Total	1,166	0	0	1,166	1,007	44	115
Facility	Total	39,999	30,415	365	9,219	5,606	2,145	1,471

¹ There were 59 males taken directly from the trap and spawned. These fish were not included in the total fish ponded.

The pattern of arrival timing by week (Sunday through Saturday) for adult fall Chinook Salmon to the PRH Volunteer Trap was determined to help schedule future sampling and broodstock activities. Trap operations during 2020 should have provided unimpeded access to the trap during most of each week. Rarely was the trap closed for an entire day because of exceeding its holding capacity during large spikes in returns. The collection numbers suggest that peak arrival to the PRH Volunteer Trap during 2020 occurred during the last week of October and first week of November which is similar to the average timing for years 2014-2019 (Figure 3).




Figure 3 Proportion of annual returns by week (Sunday through Saturday) adult Chinook Salmon collected at the Priest Rapids Hatchery Volunteer Trap. First week generally begins between September 7 and 15.

PRH has four adult Salmon holding ponds. Ponds 2 and 3 were used to hold broodstock collected at the PRH Volunteer Trap. Pond 1 was used to hold broodstock for the Yakama Nation to supplement production at other facilities. Pond 4 was used to hold broodstock collected from the ABC. The PRH staff generally transported fish from the volunteer trap on non-spawn days to collect broodstock and or to surplus the excess fish. Male fall Chinook Salmon typically comprised the majority of fish surplus from the trap. Spawn days generally occurred on Tuesdays and Wednesdays each week from October 27 through December 1 (N = 8). The electro-anesthesia (EA) system was regularly used during surplus and spawning activities during 2020.

The egg-take goal from the 2020 PRH brood was 13,530,000 eggs. The actual egg-take for the Grant PUD and USACE programs was 13,971,184 (~103% of the goal). In general, the spawning protocol includes stripping eggs from two females into a five-gallon bucket and then adding milt from a single male. Two buckets of fertilized eggs were then combined to help ensure fertilization. These buckets were then transferred to the incubation room, weighed to estimate numbers of eggs, and placed in vertical incubation trays at roughly 7,000 eggs per tray.

The alternative mating strategy of crossing 1 male with 4 females that was implemented during prior years (Richards and Pearsons 2018) did not occur during 2020 because there were sufficient broodstock collected from the ABC to allow mating of males and females at 1:2 ratio for the Grant PUD program. A total of 566 males from the ABC were crossed with 378 females from the ABC and 752 females from the PRH volunteer trap. Of the total 441 ABC females spawned, 63 were spawned with males collected from the PRH volunteer trap.

There were 4,338,660 eyed eggs shipped to Bonneville Hatchery for hatching and early rearing. For the combined programs at PRH, 7,759,659 fry were moved from the vertical trays in the incubation building to outdoor raceways between January 26 and February 24, 2021. The fry were reared in the raceways until they were of sufficient size that a portion of them could be marked in some manner (i.e., adipose clipped and or tagged). Fish receiving marks and or tags were collected directly from the raceways banks and then released into the corresponding concrete rearing ponds (e.g., fish moved from raceway bank E to channel pond E). Fish not selected for marking were transferred from the raceway banks into the corresponding rearing ponds. The growth of smolts from ponds E and D was accelerated for early releases that occurred on May 24 and 27, respectively. The remaining smolts were released between June 09 and June 17. All releases occurred at night. These fish migrate down the old 1.6 km long spawning channel and then down the hatchery discharge channel to the Columbia River.

7.0 Origin of Adult Returns to Priest Rapids Hatchery

The origin of fish collected from the different locations was determined by examination of hatchery marks (i.e., otolith thermal marks, adipose clips, and CWTs) for the fish within the demographic sample groups. PRH origin fish were identified by their otolith mark or a CWT. The fish that did not possess an otolith mark or other hatchery marks and tags were classified as natural origin. Historically, the very low recovery (<1%) of non-adipose clipped CWT strays at PRH suggests that a high percentage of the fish not possessing any type of hatchery mark may be of natural origin. In some sections of this report, we make a simplifying assumption that fish without hatchery marks are of natural origin.

Similar to that observed in previous years, there is a discrepancy between estimates of origin based on CWT and those based on otoliths marks. It's believed that estimates of origin based on

otolith sampling may provide the most accurate data under the current marking regime at PRH due to discrepancies in the data associated with CWT results (Appendix A).

An examination of thermal mark accuracy was conducted for 2020 where 288 known origin otoliths were blindly examined amongst the Hanford Reach spawning survey. An overall error rate of 2.4% was detected from the known origin samples. The decoding errors included 6 (2.6%) that were found to be false negatives (no mark was assigned when a mark should have been present N= 231). There was 1 (1.8%) false positive (thermo mark reported when it should not have been present, N=57). The implication of this error is that natural-origin fish are likely overestimated, and hatchery-origin fish are underestimated. Preliminary results suggest a directional bias which were like results found in Volk et al. (1999) for false negative error (1-5%).

We present estimates of abundance based on CWTs (1:1 sample rate) and estimates based on sub-samples of hatchery marked fish collected from specific groups (varying sample rates) to illustrate differences in the estimates for the proportions of natural and hatchery origin fish recovered at PRH as well as the potential for creating a method to correct the historical database that was generated using CWT recoveries.

7.1 Origin Based on Otolith Marks

For return year 2020, the proportion of broodstock obtained from the PRH volunteer trap that was natural origin was estimated as 0.045. Overall, the proportion of natural origin fish surplused or removed as mortalities that originated from the PRH volunteer trap was estimated at 0.033. The proportion of natural origin fish used as broodstock from the ABC was estimated to be 0.742. The estimated numbers of natural and hatchery origin broodstock spawned annually since return year 2013 are given in Table 2.

For return years 2014 through 2020, a minimum fork-length threshold of ~73 cm was generally used to reduce the number of age-2 and 3 male broodstock collected at OLAFT and the PRH volunteer trap along with the exclusion of hatchery marks and tags. Historical data suggests that age-2 and 3 fall Chinook Salmon returning to the Hanford Reach comprise a greater proportion of hatchery origin fish compared to age-4 and 5 fall Chinook Salmon returning to the Hanford Reach.

Table 2 Total fish handled, numbers sampled, and estimates of hatchery and natural origin Chinook Salmon collected at Priest Rapids Hatchery, Priest Rapids Dam Off-Ladder Adult Fish Trap, and Angler Broodstock Collection fishery. Origin determined by otolith thermal marks, presence of coded-wire tags, and/or adipose clips, Return Years 2013-2020.

Priest Rapids Hatchery Broodstock ¹			Estimate (95% CI)	
Return Year	Total	(N)	Hatchery Origin	Natural Origin ²
2013	4,476	503	4,395 [4,319, 4,436]	81 [40, 157]
2014	4,427	574	4,228 [4,130, 4,294]	199 [133, 297]
2015	4,875	682	4,482 [4,368, 4,573]	393 [302, 507]
2016	4,324	827	4,067 [4,034, 4,095]	257 [227, 290]
2017	4,511	533	4,093 [3,967, 4,197]	417 [414, 543]
2018	4,039	717	3,478 [3,389, 3,594]	556 [440, 645]
2019	3,691	641	3,259 [3,167, 3,351]	432 [340, 524]
2020	4,599	817	4,393 [4,325, 4,456]	206 [143, 274]
Priest Rapids Hatchery Surplused from Trap			Estimate (95% CI)	
Return Year	Total	(N)	Hatchery Origin	Natural Origin ²
2013 ^a	37,355	608	36,085 [35,375, 36,533]	1,270 [822, 1,980]
2014 ^b	73,352	639	69,024 [67,484, 70,271]	4,328 [3,081, 5,868]
2015 ^b	57,625	619	54,646 [53,418, 55,551]	2,979 [2,075, 4,207]
2016 ^a	24,461	1,033	23,790 [23,737, 23,837]	668 [619, 719]
2017 ^a	13,301	1,426	11,954 [10,680, 10,803]	1,348 [1218, 1492]
2018 ^a	12,137	1,001	9,811 [9,443, 10,051]	2,326 [2,086, 2,694]
2019 ^a	15,735	1,068	14,637 [14,240, 14,744]	1,098 [991, 1,495]
2020 ^a	34,234	1,460	33,132 [32,776, 33,423]	1,139 [811, 1,458]
Off Ladder Fish Trap Broodstock ¹			Estimate (95% CI)	
Return Year	Total	(N)	Hatchery Origin	Natural Origin ²
2013	763	169	343 [242, 370]	420 [392, 416]
2014	825	225	143 [122, 166]	682 [659, 703]
2015	348	164	45 [29, 66]	303 [282, 319]
2016	366	211	99 [83, 117]	267 [249, 283]
2017	809	226	108 [78, 148]	701 [661, 731]
2018	710	195	84 [57, 121]	626 [589, 653]
2019	568	94	73 [34, 111]	495 [457, 534]
2020	0	0		
Angler Broodstock Collection Broodstock ¹			Estimate (95% CI)	
Return Year	Total	(N)	Hatchery Origin	Natural Origin ²
2013	308	293	59 [46, 75]	249 [233, 262]
2014	221	111	17 [9, 34]	204 [187, 212]
2015	301	141	11 [4, 26]	290 [275, 297]
2016	247	94	11 [6, 20]	236 [227, 241]
2017	348	171	33 [20, 52]	315 [296, 328]
2018	1,087	499	78 [56, 108]	1,007 [977, 1029]
2019	1,223	216	159 [136, 182]	1,064 [1,041, 1087]
2020	1,004	229	259 [203, 321]	745 [683, 801]

¹ Includes only fish that were spawned.

² Origin based on the absence of otolith marks, coded-wire tags, or adipose clips.

^a These data were collected from samples intermittently high-graded for broodstock and may not be representative of the entire return to the Priest Rapids Hatchery volunteer trap.

^b These data are representative of the entire volunteer return to the Priest Rapids Hatchery volunteer trap.

7.2 Origin Based on Coded-Wire Tag Recoveries

The expansions of CWT recoveries at PRH until recent years have notably underestimated the returns of PRH origin fish by return year and brood year. This bias and steps taken to identify the source are provided in Appendix A. The majority of CWTs recovered at PRH originate from PRH, however CWTs originate from hatcheries throughout the basin as well as from tagging of natural origin fall Chinook Salmon in the Hanford Reach.

All Chinook Salmon returning to PRH and broodstock collected from the OLAFT and ABC were sampled for the presence of CWT. A total of 5,679 CWT fish were recovered from Chinook Salmon sampled at PRH in 2020, of which 486 were recovered from the broodstock obtained from the PRH volunteer trap (Appendix B). The broodstock collected from the PRH volunteer trap were generally culled to exclude CWT fish for the purpose of increasing natural origin broodstock. Therefore, this CWT group is not representative of the volunteer broodstock. The ABC fish were not screened for a CWT during collection but were later scanned for CWT at the hatchery. There were 44 CWT recovered from the ABC collection of which 40 were spawned and the rest were surplus. The juvenile mark rate expansions of CWT recovered adults at PRH in 2020 suggest that 90.6% of the returns to the PRH volunteer trap were hatchery origin fish. If we were to assume that these CWT expansions accurately reflected the proportion of hatchery origin fish, then the remaining 9.4% of the unaccounted fish could potentially be natural origin (Table 3).

There were 19 natural origin CWT Hanford Reach fall Chinook Salmon recovered at the hatchery in 2020 of which 18 were excluded from the broodstock while sorting out adipose clipped fish to increase the proportion of natural origin broodstock. There is not an expansion factor for the natural origin CWT fish so there was no attempt to estimate the proportion of natural origin fish based on these CWT recoveries.

Table 3 Estimated proportion of hatchery and natural origin adult Chinook Salmon returning to the Priest Rapids Hatchery Volunteer Trap based on coded-wire tag expansion. The entire collection was sampled for coded-wire tags, Return Years 2005-2020.

Return Year	Returns to Priest Rapids Hatchery Volunteer Trap	Origin based on Coded-Wire Tag expansions		Natural Origin ¹
		Priest Rapids Hatchery	Other Hatchery	
2005	10,616	0.622	0.006	0.372
2006	8,223	0.490	0.006	0.504
2007	6,000	0.671	0.004	0.325
2008	19,586	0.491	0.008	0.501
2009	12,778	0.428	0.003	0.569
2010	19,169	0.602	0.003	0.395
2011	20,823	0.613	0.006	0.381
2012	28,039	0.692	0.004	0.304
2013	41,831	0.713	0.034	0.253
2014	77,259	0.809	0.020	0.171
2015	63,978	0.914	0.015	0.071
2016	28,786	0.912	0.024	0.064
2017	17,812	0.868	0.046	0.086
2018	16,171	0.737	0.023	0.240
2019	19,426	0.802	0.034	0.164

Return Year	Returns to Priest Rapids Hatchery Volunteer Trap	Origin based on Coded-Wire Tag expansions		Natural Origin ¹
		Priest Rapids Hatchery	Other Hatchery	
2020	39,948	0.844	0.062	0.094
Mean	26,903	0.701	0.019	0.281
Median	19,506	0.703	0.012	0.279

¹ The proportion not accounted for by coded-wire tag expansion is assumed to be of natural origin.

8.0 Broodstock Collection and Sampling

The broodstock collected at the PRH volunteer trap were systematically sampled at a rate of 1:5 for otoliths (origin), scales (age), gender, and length. The broodstock collected from the ABC were sampled at a 1:4 rate for otoliths (origin), scales (age), gender, and length.

8.1 Broodstock Age Composition

A combined total of 5,630 fish were spawned from the PRH Volunteer Trap and ABC to provide green eggs for the combined Grant PUD and USACE production at PRH and the USACE production at RSH. The broodstock age compositions reported for years prior to 2012 are not directly comparable to the 2012 through 2020 broodstock age compositions due to inconsistent methodology for assigning origin and selecting broodstock based on fork length (Table 4, Table 5, and Table 6). Prior to 2012, the origin of broodstock was estimated by adult CWT recoveries which in turn were expanded by the specific juvenile tag rates. In addition, the broodstock age compositions for 2014 through 2020 are influenced by the inconsistent selection of broodstock based on a 73 cm minimum fork length at OLAFT and the volunteer trap. In addition, jacks collected in the ABC fishery were seldom used for broodstock.

Table 4 Age composition for hatchery and natural origin fall Chinook Salmon spawned at Priest Rapids Hatchery (includes all sources of broodstock), Return Years 2007-2020. Proportions calculated from expanded age compositions by origin for each source of broodstock to account for differing sample rates.

Return Year	Origin	Age Composition				
		Age-2	Age-3	Age-4	Age-5	Age-6
2007	Natural ¹	0.000	1.000	0.000	0.000	0.000
	Hatchery ¹	0.081	0.274	0.486	0.138	0.020
2008	Natural ¹	--	--	--	--	--
	Hatchery ¹	0.011	0.848	0.100	0.039	0.002
2009	Natural ¹	--	--	--	--	--
	Hatchery ¹	0.012	0.086	0.883	0.019	0.000
2010	Natural ¹	--	--	--	--	--
	Hatchery	0.016	0.755	0.111	0.118	0.000
2011	Natural ¹	--	--	--	--	--
	Hatchery ¹	0.010	0.229	0.753	0.008	0.000
2012	Natural ²	0.032	0.435	0.400	0.131	0.002
	Hatchery ²	0.006	0.487	0.376	0.130	0.000
2013	Natural ²	0.000	0.446	0.517	0.037	0.000
	Hatchery ²	0.001	0.658	0.339	0.002	0.000
2014	Natural ²	0.000	0.045	0.886	0.070	0.000
	Hatchery ²	0.000	0.064	0.897	0.039	0.000
2015	Natural ²	0.000	0.183	0.506	0.305	0.006

Return Year	Origin	Age Composition				
		Age-2	Age-3	Age-4	Age-5	Age-6
2016	Hatchery ²	0.000	0.210	0.680	0.110	0.000
	Natural ²	0.000	0.101	0.761	0.138	0.000
	Hatchery ²	0.000	0.099	0.700	0.196	0.007
2017	Natural ²	0.000	0.130	0.618	0.252	0.000
	Hatchery ²	0.000	0.074	0.663	0.258	0.005
2018	Natural ²	0.000	0.448	0.419	0.130	0.003
	Hatchery ²	0.000	0.361	0.526	0.105	0.008
2019	Natural ²	0.000	0.263	0.691	0.046	0.000
	Hatchery ²	0.000	0.231	0.745	0.023	0.000
2020	Natural ²	0.000	0.260	0.635	0.105	0.000
	Hatchery ²	0.000	0.382	0.592	0.025	0.000
Mean (2012-19)	Natural ²	0.004	0.257	0.604	0.135	0.001
	Hatchery ²	0.001	0.285	0.613	0.099	0.002

¹ Origin determined from coded-wire tag expansions of juvenile mark rate.

² Origin determined from presence of hatchery marks (i.e., coded-wire tags, adipose clips, and otoliths)

Table 5 Age composition for hatchery and natural origin fall Chinook broodstock collected from the Priest Rapids Hatchery volunteer trap, Return Years 2012-2020.

Return Year	Origin ¹	N	Age Composition				
			Age-2	Age-3	Age-4	Age-5	Age-6
2012	Natural	249	0.000	0.295	0.585	0.121	0.000
	Hatchery	4,158	0.000	0.477	0.389	0.134	0.000
2013	Natural	81	0.000	0.390	0.610	0.000	0.000
	Hatchery	4,476	0.000	0.656	0.342	0.002	0.000
2014	Natural	574	0.000	0.115	0.885	0.000	0.000
	Hatchery	4,427	0.000	0.065	0.899	0.036	0.000
2015	Natural	682	0.000	0.218	0.491	0.273	0.018
	Hatchery	4,875	0.000	0.215	0.668	0.116	0.000
2016	Natural	827	0.000	0.102	0.776	0.122	0.000
	Hatchery	4,324	0.000	0.100	0.763	0.136	0.000
2017	Natural	533	0.000	0.290	0.544	0.167	0.000
	Hatchery	4,511	0.000	0.075	0.662	0.258	0.005
2018	Natural	556	0.000	0.391	0.449	0.160	0.000
	Hatchery	3,478	0.000	0.351	0.535	0.106	0.008
2019	Natural	432	0.000	0.110	0.877	0.014	0.000
	Hatchery	3,259	0.000	0.197	0.781	0.023	0.000
2020	Natural	206	0.000	0.164	0.744	0.092	0.000
	Hatchery	4,393	0.000	0.368	0.608	0.024	0.000
Mean	Natural	460	0.000	0.231	0.662	0.105	0.002
	Hatchery	4,211	0.000	0.278	0.627	0.093	0.001

¹ Origin determined from “in-sample” otoliths, adipose clips and/or coded-wire tags.

Table 6 Age composition for hatchery and natural origin fall Chinook Salmon broodstock collected from Angler Broodstock Collection, Return Years 2012-2020.

Return Year	Origin ¹	Age Composition					
		N	Age-2	Age-3	Age-4	Age-5	Age-6
2012	Natural	59	0.000	0.542	0.339	0.119	0.000
	Hatchery	6	0.000	0.667	0.333	0.000	0.000
2013	Natural	249	0.000	0.511	0.468	0.021	0.000
	Hatchery	59	0.000	0.839	0.161	0.000	0.000
2014	Natural	204	0.000	0.126	0.830	0.044	0.000
	Hatchery	17	0.059	0.369	0.572	0.000	0.000
2015	Natural	290	0.000	0.196	0.499	0.305	0.000
	Hatchery	11	0.000	0.397	0.603	0.000	0.000
2016	Natural	236	0.000	0.156	0.656	0.189	0.000
	Hatchery	11	0.000	0.250	0.750	0.000	0.000
2017	Natural	315	0.000	0.127	0.561	0.312	0.000
	Hatchery	33	0.000	0.055	0.649	0.296	0.000
2018	Natural	1,007	0.000	0.433	0.417	0.143	0.006
	Hatchery	78	0.000	0.493	0.425	0.082	0.000
2019	Natural	1,064	0.000	0.175	0.788	0.037	0.000
	Hatchery	159	0.000	0.713	0.287	0.000	0.000
2020	Natural	768	0.000	0.104	0.734	0.161	0.000
	Hatchery	263	0.000	0.294	0.613	0.092	0.000
Mean	Natural	466	0.000	0.263	0.588	0.148	0.001
	Hatchery	71	0.007	0.453	0.488	0.052	0.000

¹ Origin determined from “in-sample” otoliths, adipose clips and/or coded-wire tags.

8.2 Length by Age Class of Broodstock

The mean fork length (cm) by age for each source of broodstock is provided in Table 7 and Table 8. For the most part, natural origin fish were larger for all ages for each source of broodstock. The exceptions were hatchery origin age-3 fish collected at the PRH volunteer trap appear to be larger than age-3 fish at the OLAFT and age-4 natural origin fish were longer than hatchery origin fish collected at the hatchery trap.

Table 7 Mean fork length (cm) at age (total age) of fall Chinook Salmon sampled from each source of broodstock spawned at Priest Rapids Hatchery, Return Year 2020. N = sample size and SD = 1 standard deviation.

Source of Broodstock	Origin ¹	Fall Chinook Fork Length (cm)														
		Age-2			Age-3			Age-4			Age-5			Age-6		
		N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
Volunteer Returns	Natural	0	--	--	6	74	3	26	79	5	3	89	3	--	--	--
	Hatchery	0	--	--	274	71	4	451	80	5	16	85	7	--	--	--
ABC	Natural	0	--	--	45	69	6	94	81	7	16	89	8	--	--	--
	Hatchery	0	--	--	32	67	4	16	78	3	2	75	1	--	--	--

¹ It is assumed for this analysis that all fish not possessing an otolith mark, ad-clipped or hatchery origin coded-wire tag were natural origin.

Table 8 Mean fork length (cm) at age (total age) of hatchery and natural origin fall Chinook Salmon collected from broodstock originating from the Priest Rapids Hatchery volunteer trap. N = sample size and SD = 1 standard deviation, Return Years 2012-2020.

Return Year	Origin ¹	Fall Chinook Fork Length (cm)														
		Age-2			Age-3			Age-4			Age-5			Age-6		
		N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
2012	Natural	0	--	--	12	71	4	25	82	4	5	86	4	0	--	--
	Hatchery	0	--	--	298	70	4	253	81	5	91	88	7	0	--	--
2013	Natural	0	--	--	4	76	4	7	78	4	0	--	--	0	--	--
	Hatchery	0	--	--	288	71	4	200	80	5	2	85	4	0	--	--
2014	Natural	0	--	--	3	74	2	23	80	5	0			0	--	--
	Hatchery	0	--	--	36	70	3	491	78	5	21	87	6	0	--	--
2015	Natural	0	--	--	12	74	7	30	79	6	15	86	4	1	87	0
	Hatchery	0	--	--	133	71	4	437	80	4	79	84	5	0	--	--
2016	Natural	0	--	--	78	73	3	594	79	4	106	85	6	0	--	--
	Hatchery	0	--	--	133	71	4	437	80	4	79	84	5	0	--	--
2017	Natural	0	--	--	15	73	4	26	79	4	8	81	8	0	--	--
	Hatchery	0	--	--	39	72	4	315	77	4	127	82	6	3	84	3
2018	Natural	0	--	--	50	71	4	53	79	5	17	84	5	0	--	--
	Hatchery	0	--	--	230	70	4	342	78	5	69	81	5	7	82	3
2019	Natural	0	--	--	46	70	5	288	81	5	11	84	8	--	--	--
	Hatchery	0	--	--	133	71	4	459	79	5	14	83	6	--	--	--
2020	Natural	0	--	--	6	74	3	26	79	5	3	89	3	--	--	--
	Hatchery	0	--	--	274	70	5	451	80	5	16	85	7	--	--	--
Mean	Natural	0	--	--	25	73	4	119	80	5	18	85	5	0	87	0
	Hatchery	0	--	--	174	71	4	376	79	5	55	84	6	1	83	3

¹It is assumed for this analysis that all fish not possessing an otolith mark, ad-clipped or hatchery origin coded-wire tag were natural origin.

8.3 Gender Ratios

PRH staff sort and select broodstock from the volunteer trap to meet their egg-take goals and male-to-female spawner ratio of 1:2. The broodstock collected from the ABC were spawned at a 1:2 male-to-female ratio as well. The 2020 broodstock population was comprised of 66.4% females, resulting in an overall male to female ratio of 0.51:1.00, which is lower than the historic mean ratio of 0.53:1.00 (Table 9).

Table 9 Number of male and female hatchery fall Chinook Salmon broodstock at Priest Rapids Hatchery, Return Years 2001-2020. Ratios of males to females are also provided.

Return Year	Males (M)	Females (F)	M/F Ratio
2001	1,697	3,289	0.52:1.00
2002	1,936	3,628	0.53:1.00
2003	1,667	3,176	0.52:1.00
2004	1,688	3,099	0.54:1.00
2005	1,962	3,326	0.59:1.00
2006	1,777	3,322	0.53:1.00
2007	850	1,301	0.65:1.00

Return Year	Males (M)	Females (F)	M/F Ratio
2008	1,823	3,195	0.57:1.00
2009	1,531	3,000	0.51:1.00
2010	1,809	3,447	0.52:1.00
2011	1,858	3,000	0.62:1.00
2012	1,749	3,225	0.54:1.00
2013	1,865	3,578	0.52:1.00
2014 ^a	1,805	3,688	0.49:1.00
2015 ^a	1,697	3,827	0.44:1.00
2016 ^a	1,537	3,401	0.45:1.00
2017 ^a	1,835	3,835	0.48:1.00
2018 ^a	1,863	3,955	0.47:1.00
2019	1,835	3,640	0.50:1.00
2020	1,891	3,740	0.51:1.00
Mean	1,725	3,312	0.53:1.00

^a Includes broodstock used in the 1-male x 4-females alternative mating strategy.

8.4 Fecundity

The annual mean fecundity for PRH was calculated as the proportion of the total number of females spawned to the total estimated take of green eggs. The total number of green eggs is calculated after the first pick of both live and dead eggs from the incubation trays. Fish culture staff weigh large lots of either dead or live eggs and then sub-sample the lots to calculate a mean individual egg weight. The number of eggs per lot is estimated by dividing the weight of each egg lot by the calculated mean individual egg weight. The egg count for each lot is summed to estimate the facility egg-take. Each egg lot likely contained small and varying amounts of interstitial water which might overestimate the egg count.

Fecundity for the 2020 broodstock averaged 3,748 eggs per female which is similar to that observed in recent years but less than the historical mean of 3,925 (Table 10). Pre-spawn egg loss was often observed during the electro-anesthetic and pneumatic fish euthanizing process (a physical strike to the head) and may contribute to the reduced fecundity of fish in recent years. In addition, the size and associated fecundity of Chinook Salmon populations has been declining coast-wide and the reduction in fecundity at PRH may be the result of larger regional factors (Ohlberger et. al. 2018).

Table 10 Mean fecundity of fall Chinook Salmon collected for broodstock at Priest Rapids Hatchery, Return Years 2001-2020. SD = 1 standard deviation.

Return Year	Egg-Take	Viable Females	Fecundity/Female
2001	10,750,000	3,161	3,401
2002	12,180,000	3,489	3,491
2003	12,814,000	3,078	4,163
2004	12,753,500	3,019	4,224
2005	14,085,000	3,211	4,386
2006	13,511,200	3,217	4,200
2007 ^a	5,067,319	1,249	4,057
2008	12,643,600	3,074	4,113
2009	13,074,798	2,858	4,575
2010	11,903,407	3,342	3,562
2011	12,693,000	3,038	4,178

Return Year	Egg-Take	Viable Females	Fecundity/Female
2012	12,398,389	3,053	4,061
2013	13,316,000	3,473	3,834
2014	14,321,183	3,563	4,019
2015	13,530,988	3,706	3,651
2016	12,411,530	3,401	3,649
2017	13,738,916	3,763	3,651
2018	14,821,007	3,975	3,729
2019	13,898,011	3,642	3,815
2020	13,971,184	3,728	3,748
Mean	12,607,522	3,252	3,925
SD	2,028,767	561	317

^a Broodstock shortage due to low adult return to Priest Rapids Hatchery volunteer trap.

Fecundities of individual females were taken from sub-samples at PRH during the spawn of 2010 through 2020 broodstock to estimate fecundity by length and age. For the 2013 through 2020 brood year data, we show comparisons between hatchery and natural origin fall Chinook Salmon sampled at PRH that include fork length/fecundity, fork length/egg size (weight) and fork length, and gamete mass. For these years, we attempted to stratify the females sampled by fork length categories to obtain fecundity samples for all sizes of fish to better estimate the relationship between length and fecundity. However, the broodstock selection protocols in recent year have reduced the availability of females under 64 cm. Some fecundity data were obtained from females not used for broodstock (i.e., surplus) to bolster sample sizes. Therefore, comparisons between age classes are not representative of the females spawned from 2013 through 2020 broodstock populations.

M&E staff performed the fecundity estimates on green eggs. The entire gamete mass was drained of most all ovarian fluid and weighed within 0.1 gram. Sub-sample sizes ranged between years from 60 or 100 green eggs which were counted out and weighed within 0.01 gram to estimate individual egg weight (g) for each female. Post brood year 2013, sample sizes were 100 eggs, which was determined to be sufficient based upon previous work that examined different samples sizes (Richards and Pearsons, 2014). The total fecundity of each female was estimated by dividing the weight of the total egg mass by the calculated mean individual egg weight. Each sample of the total egg mass likely contained slight varying amounts of ovarian fluid which might overestimate fecundity.

The fecundity data was pooled for return year 2010 through 2020 to provide a simple linear regression to predict fecundity based on fork-length (natural and hatchery origin females combined). This data shows a strong positive correlation between size and fecundity (Figure 4). The regression formula may be useful for coarse predictions of egg production for different size fish.




Figure 4 Linear relationship between fecundity and fork length for combined samples of natural and hatchery origin fall Chinook Salmon spawned at Priest Rapids Hatchery, Return Years 2010-2020.

Fecundity samples collected at PRH for years 2010 through 2012 were not identified as to the origin of the females. For years 2013 through 2020, fecundity samples were taken from the broodstock at PRH to collect data associated with fecundity by size, age and origin (hatchery or natural).

Females were selected from both the PRH volunteer broodstock as well as from ponds which possessed broodstock primarily from the OLAFT and ABC. For the most part, the origin of fish during sampling was unknown. Therefore, we made a concerted effort to select females that were not adipose clipped to increase the chances of obtaining natural origin fish which were less common than hatchery origin fish. The origins of females sampled for fecundity were determined by hatchery marks (i.e., otoliths, adipose clips and CWTs). We assume that fish not possessing any type of hatchery marks were of natural origin.

The mean fecundity by age is given in Table 11. This information is useful for forecasting potential egg-takes based on the numbers and age composition of the forecasted return.

Table 11 Mean fecundity at age for fall Chinook Salmon sampled at the Priest Rapids Hatchery, Return Years 2010-2020. N = sample size and SD = 1 standard deviation.

Return Year	Age-3			Age-4			Age-5		
	N	Mean	SD	N	Mean	SD	N	Mean	SD
2010	273	3,658	834	17	3,664	585	1	4,217	--
2011	30	3,538	842	206	4,276	884	1	4,380	--
2012	2	3,639	882	3	4,282	1089	0	--	--
2013	105	3,488	768	68	4,152	788	4	5,339	805
2014	1	3,358	--	73	4,126	755	5	4,416	407

2015	5	3,169	382	53	3,662	606	25	4,746	691
2016	14	3,192	559	101	3,676	639	36	4,173	693
2017	0	--	--	65	3,754	689	31	4,163	712
2018	17	3,997	771	80	3,876	757	26	3,850	689
2019	4	3,458	434	89	3,949	721	8	4,323	689
2020	3	3,802	340	14	3,757	563	7	4,341	1,041
Mean	41	3,530	646	70	3,925	734	13	4,395	716

The data collected from return years 2013 through 2020 were pooled to increase the number of samples for a given fork length. The linear relationships between fork length and variables including fecundity, total egg mass weight, and mean egg weight for natural and hatchery origin females subsampled are plotted Figure 5, Figure 6 and Figure 7. All relationships show a positive correlation with fork length. In addition, the relationships between fish size and egg data were similar for hatchery and natural origin fish.




Figure 5 Fecundity versus fork length for natural- and hatchery- origin fall Chinook Salmon sub-sampled at Priest Rapids Hatchery, Return Years 2013-2020.




Figure 6 Mean skein weight versus fork length for natural- and hatchery- origin fall Chinook Salmon sub-sampled at Priest Rapids Hatchery, Return Years 2013-2020.




Figure 7 Total egg mass weight versus fork length for natural- and hatchery- origin fall Chinook Salmon sub-sampled at Priest Rapids Hatchery, Return Years 2013-2020.

9.0 Hatchery Rearing

9.1 Number of Eggs Taken

In 2020, an estimated 13,971,184 eggs were collected at PRH (Table 12) to meet the production targets at PRH and RSH. The egg-take target for return year 2020 was 13,530,000 based on current program needs. An additional 915,800 green eggs were collected by staff from the Yakama Nation for their fall Chinook Salmon production at Prosser Hatchery adjacent to the Yakima River.

PRH incubates approximately 8.4 million eyed eggs to produce the 7.3 million smolt release at the hatchery. An additional 4.1 million eyed eggs are needed to meet the program goal of eyed eggs delivered to Bonneville Hatchery for the 3.5 million subyearling releases from RSH. There were 798,767 green eggs taken which after picking dead eggs produced a surplus of 748,692 eyed eggs of which 517,474 were culled and the remaining 231,218 provided for transfer to off-site production for other programs.

Table 12 Number of eggs taken from fall Chinook Salmon broodstock collected at Priest Rapids Hatchery, Return Years 1984-2020.

Return Year	Number of Eggs Taken	Return Year	Number of Eggs Taken
1984	10,342,000	2004	12,753,500
1985	10,632,000	2005	14,085,000
1986	22,126,100	2006	13,511,200
1987	24,123,000	2007	5,067,319
1988	16,682,000	2008 ^a	12,643,600
1989	13,856,500	2009	13,074,798
1990	9,605,000	2010	11,903,407
1991	6,338,000	2011	12,693,000
1992	11,156,400	2012	12,398,389
1993	14,785,000	2013	13,276,000
1994	16,074,600	2014	14,321,818
1995	17,345,900	2015	13,530,988
1996	14,533,500	2016	12,411,530
1997	17,007,000	2017	13,738,916
1998	13,981,300	2017	13,738,916
1999	16,089,600	2018	14,821,007
2000	15,359,500	2019	13,898,011
2001	10,750,000	2020	13,971,184
2002	12,180,000	10-year (11-20) Mean	13,339,271
2003	12,814,000	10-year (11-20) SD	775,085

^a Began annual egg-takes starting in return year 2008 for the 3.5 million Ringold Springs Hatchery Program

9.2 Number of Acclimation Days

The 2020 brood were incubated on a combination of well and river water before being transferred to intermediate concrete raceways and then transferred to the concrete holding ponds for final acclimation before release into the Columbia River in late May and June 2021. The egg-takes for the 2020 brood were distributed into nine batches associated with the dates in which fish were spawned. The rearing-acclimation days ranged from 106 to 118 (Table 13).

Table 13 Number of days fall Chinook Salmon fry were reared at Priest Rapids Hatchery prior to release, Brood Year 2020.

Batch	Egg Tray to Raceway Transfer Date	Release Date	Number of Days
1	January 26 into Bank E	May 24	118
2	Shipped off Station	--	--
3	February 8 into Bank E	May 24	118
3	February 8 into Bank D	May 27	108
3	February 8 into Bank C	June 9	106
4	February 8 into Bank C	June 9	106
5	February 23 into Bank C	June 9	106
5	February 23 into Bank B	June 14	111
6	February 24 into Bank A	June 17	113
7	February 24 into Bank A	June 17	113
8	February 24 into Bank A	June 17	113

9.3 Annual Releases, Tagging, and Marking

The annual release of fall Chinook Salmon smolts from PRH has ranged considerably since the initial release of roughly 2.38 million smolts from the 1979 brood year to over roughly 10.30 million from the 1982 brood year (Table 14). The 2021 release goal for PRH was 7,299,504 smolts. This goal includes a recent increase in the Grant PUD mitigation from 5,000,000 to 5,599,504 combined with the ongoing USACE’s John Day mitigation of 1,700,000 smolts.

In 2021, staff at PRH released an estimated 7,542,518 subyearling fall Chinook Salmon from the 2020 broodstock. Fish were released between May 24 and June 17.

Various mark types and rates have occurred at PRH over the years for both the Grant PUD and USACE mitigation fish. In 1976, PRH began adipose fin clipping and coded-wire tagging a portion of the juvenile fall Chinook released to determine PRH contributions to ocean and river fisheries. The smolt production at PRH associated with the USACE mitigation increased the number of adipose clipped smolts released by ~1.7 million starting with brood year 2006. The number of coded-wire tagged fish released from PRH increased to >1.2 million fish starting with brood year 2009 of which ~600,000 were adipose clipped. An additional 1 million adipose clipped smolts were included in the release since brood year 2011. Fish released in 2020 (BY 2019 production) were not PIT tagged because of human health risks associated with the COVID-19 pandemic.

All PRH releases for both mitigation programs were 100% otolith marked beginning with the 2008 release. All intra-annual releases from PRH have the same annual otolith pattern, but the pattern differs between years. The eyed eggs produced for the RSH program have received an otolith mark for brood years 2010 through 2016. Otolith marking of the RSH production was discontinued beginning with the 2017 brood due to the problematic timing of completing the thermal mark prior to shipping eyed eggs to Bonneville Hatchery for hatching and early rearing, as well as all the fish to be released from RSH are adipose clipped so they can be identified as hatchery origin without examining otoliths.

Since 1987, the U.S. Section of the Pacific Salmon Commission (PSC) has supported a coordinated project which seeks to capture and tag 200,000 naturally produced juvenile fall Chinook Salmon in the Hanford Reach (Fryer 2017). Fish were collected with seines over a ten-day period between late May and early June. Fish were approximately 40-80 mm long at the time

of capture. Recoveries from these tagged fish were used to estimate harvest exploitation rates and interception rates for Hanford Reach natural origin fall Chinook Salmon. These data have also more recently been used to estimate the number of natural origin juveniles produced in the Hanford Reach (Harnish et al. 2012, Harnish 2017).

Table 14 Number of marked, unmarked, and tagged fall Chinook Salmon smolts released from Priest Rapids Hatchery, Brood Years 1977-2020.

Brood Year	Total Released	Non Ad-Clip Released	AD/CWT	CWT Only	AD Only	PIT
1977	150,625	0	147,338	0	3,287	
1978	153,840	0	152,532	0	1,308	
1979	3,005,654	2,858,509	147,145	0		
1980	4,832,591	4,581,054	251,537	0		
1981	5,509,241	5,198,365	310,876	0		
1982	10,296,700	9,888,989	407,711	0		
1983	9,742,700	9,517,263	222,055	0	3,382	
1984	6,363,000	6,253,240	106,960	0	2,800	
1985	6,048,000	5,843,176	203,534	0	1,290	
1986	7,709,000	7,506,142	201,843	0	1,015	
1987	7,709,000	7,501,578	196,221	0	11,201	
1988	5,404,550	5,200,080	201,608	0	2,862	
1989	6,431,100	6,224,770	194,530	0	11,800	
1990	5,333,500	5,134,031	199,469	0		
1991	7,000,100	6,798,453	201,647	0		
1992	7,134,159	6,939,537	194,622	0		
1993	6,705,836	6,520,153	185,683	0		
1994	6,702,000	6,526,120	175,880	0		1,500 ^c
1995	6,700,000	6,503,811	196,189	0		3,000 ^c
1996	6,644,100	6,450,885	193,215	0		3,000 ^c
1997	6,737,600	6,541,351	196,249	0		3,000 ^c
1998	6,504,800	6,311,140	193,660	0		3,000 ^c
1999	6,856,000	6,651,664	204,336	0		3,000 ^c
2000	6,862,550	6,661,771	200,779	0		3,000 ^c
2001	6,779,035	6,559,109	219,926	0		3,000 ^c
2002	6,777,605	6,422,232	355,373	0		3,000 ^c
2003	6,814,560	6,415,444	399,116	0		3,000 ^c
2004	6,599,838	6,399,766	200,072	0		3,000 ^c
2005	6,876,290	6,676,845	199,445	0		3,000 ^c
2006	6,743,101	4,912,487	202,000	0	1,628,614	3,000 ^c
2007 ^a	4,548,307	4,344,926	202,568	0	813 ^b	3,000 ^c
2008 ^a	6,788,314	4,850,844	218,082	0	1,719,388	2,994 ^c
2009 ^a	6,776,651	3,413,334	619,568	1,026,561	1,717,188	1,995 ^c
2010 ^a	6,798,390	3,383,859	602,580	1,108,990	1,702,961	3,000 ^c
2011 ^a	7,056,948	3,094,666	595,608	598,031	2,768,643	42,844 ^c
2012 ^a	6,822,861	2,905,694	603,930	601,009	2,712,228	42,908 ^c
2013 ^a	7,267,248	3,347,417	603,417	603,439	2,712,975	42,908 ^c
2014 ^a	7,039,544	3,125,734	600,688	600,730	2,712,392	42,621 ^c
2015 ^a	7,242,054	3,317,992	602,116	601,770	2,720,176	42,999 ^d
2016 ^a	7,006,252	3,045,689	603,539	603,864	2,710,302	42,858 ^d
2017 ^a	7,987,222	4,067,088	602,725	607,287	2,710,121	42,978 ^c

Brood Year	Total Released	Non Ad-Clip Released	AD/CWT	CWT Only	AD Only	PIT
2018 ^a	7,213,916	3,311,964	603,788	601,893	2,696,272	42,990 ^c
2019 ^a	7,611,873	3,724,820	584,402	596,500	2,706,151	0
2020 ^a	7,542,518	3,629,049	601,032	603,332	2,709,105	42,873 ^c

^a Entire release was otolith marked

^b Low returns to PRH precluded the production of the USACE adipose clipped release.

^c PIT tagged fish were included within the other mark group totals

^d PIT tagged fish were not adipose clipped and reported as a unique group.

9.4 Fish Size and Condition of Release

The data associated with fish size and condition at release from PRH prior to brood year 2013 was obtained from the hatchery staff. The mean fish weight was obtained by weighing groups of roughly 300 fish sampled from each pond to the nearest gram and then dividing the group weight by the total number of fish weighed. The fork length of each fish from the group weighed was measured to the nearest millimeter to calculate mean length and coefficient of variation. Samples from each of the rearing ponds were taken the day of release. The results were pooled to provide mean estimates for the facility. The size and condition data for the 2013 through 2020 broods were collected by M&E staff the day prior to or day of release for each pond. We attempted to collect representative samples by capturing multiple groups of fish with a cast net from three to five sections evenly distributed within each rearing pond. Each fish sampled was individually weighed to the nearest 0.1 gram and measured for fork length to the nearest millimeter. Beginning with the 2019, the smolts sampled at release were also non-lethally visually inspected for precociousness. The results for each dataset were pooled to provide mean estimates for the entire facility production.

The goal for PRH is to release fall Chinook Salmon smolts at 50 fish per pound. At release, the smolts from the 2020 brood averaged 46 fish per pound with a mean fork length of 95 mm, and a mean CV of 7.2 (Table 15). For brood years 1991 through 2020, smolts released from PRH have averaged 48 fish per pound with a mean fork length of 95 and a mean CV of 7.3. No precociously mature smolts observed.

Table 15 Mean length (FL, mm), weight (g and fish/pound), and coefficient of variations (CV) of fall Chinook smolts released from Priest Rapids Hatchery, Brood Years 1991-2020.

Brood year	Release Year	Fork Length (mm)		Mean Weight		Percent Precocious smolts	N
		Mean	CV	Grams (g)	Fish/pound		
1991	1992	93	8.7	8.3	55	--	1,500
1992	1993	92	8.6	8.3	54	--	1,500
1993	1994	95	6.9	9.3	49	--	1,500
1994	1995	96	6.7	9.7	47	--	1,500
1995	1996	97	6.6	10	45	--	1,500
1996	1997	95	11	8.7	52	--	1,500
1997	1998	103	8.9	10.1	45	--	1,500
1998	1999	95	6.5	9.6	48	--	1,500
1999	2000	93	6.6	8.9	51	--	1,500
2000	2001	97	6.3	10.2	45	--	1,500
2001	2002	96	6.9	10.1	45	--	1,500
2002	2003	95	6.9	9.5	48	--	1,500
2003	2004	96	6.8	9.6	48	--	1,500

Brood year	Release Year	Fork Length (mm)		Mean Weight		Percent Precocious smolts	N
		Mean	CV	Grams (g)	Fish/pound		
2004	2005	95	5.9	9.4	48	--	1,500
2005	2006	98	6.3	10.1	45	--	1,500
2006	2007	98	7.0	9.9	46	--	1,500
2007	2008	101	8.3	10.2	45	--	1,200
2008	2009	94	6.7	9.3	49	--	1,500
2009	2010	94	7.3	9.2	49	--	1,500
2010	2011	92	9.1	9.7	47	--	1,500
2011	2012	94	7.1	9.2	49	--	1,500
2012	2013	95	7.6	9.7	47	--	1,500
2013	2014	92	8.4	9.0	50	--	648
2014	2015	91	6.6	8.7	52	--	1,728
2015	2016	92	6.1	9.3	49	--	1,595
2016	2017	89	6.1	9.3	49	--	1,788
2017	2018	91	6.6	9.2	50	--	1,633
2018	2019	90	7.4	9.0	51	--	2,382
2019	2020	94	7.2	9.6	47	0.00	1,615
2020	2021	95	7.2	9.9	46	0.00	1,795
Mean		95	7.3	9.4	48	0.00	1,529

9.5 Survival Estimates

The survival proportion (P^{\wedge}) of fertilized egg to juvenile release for brood year 2020 was 0.859 and lower than the historic mean of 0.870 (Table 16). The green egg to eyed egg stage was the most critical life stage at PRH during incubation/juvenile rearing because generally the greatest level of reported loss annually occurs at this stage. The green egg to eyed egg survival for brood year 2020 was 0.913 and higher than the historical mean of 0.899.

In 2020, survival of fish ponded for broodstock was 0.815 and lower than the historic mean of 0.826. The mean pond survival of male and females was similar.

Table 16 Hatchery life stage survival (Proportion) for fall Chinook Salmon at Priest Rapids Hatchery, Brood Years 1989-2020.

Brood year	PRH Volunteers Ponded to Spawned				Unfertilized to Eyed Egg	Eyed egg to Ponding	Ponding to Release	Fertilized Egg to Release
	Female	Male	Jack	Total				
1989	--	--	--	0.919	0.866	0.976	0.950	0.821
1990	--	--	--	0.947	0.869	0.996	0.984	0.852
1991	--	--	--	0.973	0.948	0.993	0.998	0.922
1992	--	--	--	0.952	0.945	0.991	0.965	0.901
1993	--	--	--	0.917	0.941	0.984	0.974	0.902
1994	--	--	--	0.710	0.935	0.985	0.953	0.878
1995	--	--	--	0.897	0.914	0.980	0.962	0.862
1996	--	--	--	0.908	0.924	0.997	0.983	0.897
1997	--	--	--	0.900	0.915	0.996	0.970	0.790
1998	--	--	--	0.834	0.914	0.998	0.970	0.884
1999	--	--	--	0.759	0.897	0.997	0.995	0.888
2000	--	--	--	0.868	0.898	0.995	0.985	0.884

Brood year	PRH Volunteers Poned to Spawned				Unfertilized to Eyed Egg	Eyed egg to Poding	Poding to Release	Fertilized Egg to Release
	Female	Male	Jack	Total				
2001	0.776	0.732	0.665	0.757	0.886	0.994	0.975	0.859
2002	0.835	0.829	0.705	0.828	0.880	0.995	0.979	0.858
2003	0.893	0.817	0.698	0.858	0.882	0.989	0.989	0.868
2004	0.958	0.915	0.646	0.845	0.881	0.975	0.985	0.846
2005	0.890	0.890	0.782	0.886	0.914	0.976	0.991	0.884
2006	0.918	0.924	0.695	0.913	0.897	0.975	0.981	0.859
2007	0.967	0.748	0.642	0.861	0.858	0.996	0.981	0.898
2008	0.943	0.896	0.877	0.924	0.902	0.973	0.877	0.877
2009	0.848	0.901	0.916	0.864	0.912	0.977	0.891	0.891
2010	0.803	0.831	0.803	0.809	0.913	0.985	0.977	0.841
2011	0.611	0.847	0.737	0.679	0.903	0.985	0.985	0.875
2012	0.643	0.786	0.630	0.688	0.873	0.970	0.962	0.787
2013	0.698	0.660	0.333	0.684	0.884	0.983	0.951	0.806
2014	0.830	0.880	N/A	0.847	0.865	0.933	0.978	0.913
2015	0.841	0.810	N/A	0.830	0.917	0.934	0.985	0.919
2016	0.873	0.782	N/A	0.843	0.899	0.825	0.989	0.816
2017	0.820	0.824	N/A	0.821	0.917	0.942	0.985	0.928
2018	0.831	0.895	N/A	0.869	0.943	0.978	0.924	0.903
2019	0.923	0.867	N/A	0.902	0.934	0.939	0.971	0.912
2020	0.811	0.817	N/A	0.815	0.913	0.966	0.979	0.859
Mean	0.836	0.833	0.702	0.826	0.899	0.965	0.967	0.870

9.6 Juvenile PIT Tag Detections at the Priest Rapids Hatchery Array

Roughly 3,000 sub-yearlings at PRH were annually PIT tagged and released from PRH for brood years 1995 through 2010 to assess timing, migration speed, and juvenile survival from PRH to McNary Dam. The analysis for these measures is reported annually by the Fish Passage Center and can be found at www.fpc.org/documents/FPC_memos.html.

Beginning with the 2011 brood, approximately 40,000 additional juveniles were annually PIT tagged and released to bolster the data collected for estimation of juvenile abundance at release and adult straying. These tags can also be used to estimate adult migration timing, conversion rates from Bonneville Dam to McNary Dam to PRH, as well as fallback and re-ascension estimates at McNary, Ice Harbor, and Priest Rapids dams. The annual detections at the PRH array of unique tags and rates by tag group are given in Table 17. Prior to the 2012 release (brood year 2011), a PIT tag array consisting of six antennas was installed in the hatchery discharge channel to detect both juvenile out-migrants and adult returns. The detection rates reported below account for the relatively few shed PIT tags found in the rearing raceways. Prior to the release of the 2016 brood, the mortalities routinely recovered from the rearing ponds were not scanned for PIT tags. This prohibits us from knowing the actual total number of PIT tagged fish released. Hence, the overall proportion of released PIT tagged fish detected would likely be higher than reported if we knew the actual number of live PIT tagged fish that left the ponds. In addition, PIT tagged fish that were consumed by birds prior to release were not accounted for.

The overall detection rate for the releases of the 2011 brood year was 70.4%. The releases occurred over an eight-day period, with only two days of consecutive releases. Detection rates for the 2011 brood year release may have been reduced as a result of the array being inundated by high river elevations during portions of releases. The overall detection rate for the 2012 brood year was 3.4%. The low detection rates were likely due to force releasing all the smolts in four consecutive days which appears to have overwhelmed the PIT tag detection equipment. The restricted release period was necessitated by the construction schedule of the new hatchery.

A concerted effort was made during both the 2013 and 2014 brood year releases to improve the PIT tag detection efficiency at the PRH array. First, the automatic upload function of the array was discontinued to reduce the usage demand on the system’s processor. Secondly, the five releases from the hatchery were conducted over a fourteen-day period beginning on June 12 to spread out over time the number of PIT tags passing the array. This was managed by pulling the individual weir boards for each pond over a two-day period. The percentages of PIT tagged subyearlings detected for the 2013 and 2014 brood years were 92.9% and 94.5%, respectively.

The releases of the 2015 brood occurred every two days between June 16 and June 24, 2016 to accommodate a day versus night release evaluation. During the evaluation, all weir boards for a given pond were incrementally pulled over an eight-hour period on the date of release. Overall, 84.3% of the PIT tagged subyearlings were detected. The detected rate between release groups varied from 33.6% to 97.0%. These values are lower than the previous two years. It’s possible that forced releases over an eight-hour period may have resulted in high rates of tag collision at the array resulting in poor detection efficiency.

The releases of the 2016, 2017, 2018, and 2020 broods were initiated at 9 PM for each pond. All weir boards were pulled by 3 AM. Releases occurred on different dates between May 22 and June 20 to evaluate the influence of release time on survival. The overall detection rates of these broods ranged from 86.8% to 99.1%. There were no PIT-tag groups for the 2019 brood due to the COVID 19 pandemic, however fish were released in the same manner as those for the other broods between 2016 and 2020.

Table 17 Number of sub-yearlings PIT tagged, mark, and release dates, and the number of unique tags detected at the array in the Priest Rapids discharge channel, Brood Years 2011-2018 and 2020.

Brood Year	Tag File	Tagging Date	Release Date	# Tagged	# of Tags Recovered from Facility Mortalities	# of Unique Detections	% Detected
2011	CSM12114.A01	4/23/2012	6/20/2012	9937	No Data	6,277	63.2
2011	CSM12114.A04	4/23/2012	6/14/2012	9948	No Data	6,674	67.1
2011	CSM12114.A03	4/24/2012	6/15/2012	9997	No Data	6,963	69.7
2011	CSM12115.A02	4/24/2012	6/16/2012	9967	No Data	8,115	81.4
2011	CSM12115.A02	5/30/2012	6/20/2012	1000	No Data	499	49.9
2011	SMP12151.PR2	5/30/2012	6/16/2012	998	No Data	806	80.8
2011	SMP12152.PR3	5/31/2012	6/12/2012	996	No Data	810	81.3
Totals				42,844	N/A	30,144	70.4
2012	CSM13143.A06	5/23/2013	6/14/2013	9,982	No Data	317	3.2
2012	CSM13143.A07	5/23/2013	6/13/2013	9,983	No Data	267	2.7
2012	CSM13144.A08	5/24/2013	6/12/2013	9,974	No Data	335	3.4
2012	CSM13144.A09	5/24/2013	6/15/2013	9,977	No Data	325	3.3
2012	SMP13149.PR1	5/29/2013	6/15/2013	997	No Data	131	13.1

Brood Year	Tag File	Tagging Date	Release Date	# Tagged	# of Tags Recovered from Facility Mortalities	# of Unique Detections	% Detected
2012	SMP13149.PR2	5/29/2013	6/14/2013	996	No Data	33	3.3
2012	SMP13150.PR3	5/30/2013	6/12/2013	999	No Data	48	4.9
Totals				42,908	N/A/	1,456	3.4
2013	CSM14148.PRA	5/28/2014	6/25/2014	7,994	21	7,215	90.5
2013	CSM14148.PRB	5/28/2014	6/23/2014	7,998	14	7,215	92.5
2013	CSM14149.PRC	5/29/2014	6/18/2014	7,996	11	7,443	93.2
2013	CSM14149.PRD	5/29/2014	6/16/2014	7,993	6	7,662	95.9
2013	CSM14149.PRE	5/29/2014	6/12/2014	7,998	7	7,407	92.7
2013	SMP14148.PR1	5/29/2014	6/25/2014	996	0	914	91.8
2013	SMP14148.PR2	5/29/2014	6/18/2014	994	0	927	93.3
2013	SMP14149.PR3	5/30/2014	6/12/2014	998	0	951	95.3
Total				42,967	59	39,908	92.9
2014	CSM15147.PRE	5/27/2015	6/12/2015	7,999	169	7,438	95.0
2014	CSM15147.PRD	5/27/2015	6/15/2015	7,996	39	7,685	96.6
2014	CSM15147.PRC	5/27/2015	6/18/2015	7,996	63	7,524	94.8
2014	CSM15147.PRB	5/28/2015	6/22/2015	7,998	50	7,696	96.8
2014	CSM15147.PRA	5/28/2015	6/25/2015	7,994	31	7,447	93.5
2014	SMP15140.PR1	5/20/2015	6/25/2015	993	0	940	94.7
2014	SMP15140.PR2	5/20/2015	6/18/2015	998	0	946	94.8
2014	SMP15141.PR3	5/21/2015	6/12/2015	999	0	935	93.6
Total				42,973	352	40,611	95.3
2015	CSM16153.PRE	6/01/2016	6/16/2016	7,996	13	6,032	75.6
2015	CSM16153.PRD	6/01/2016	6/18/2016	7,998	224	7,537	97.0
2015	CSM16153.PRC	6/01/2016	6/20/2016	7,985	137	6,777	86.4
2015	CSM16154.PRB	6/02/2016	6/22/2016	7,993	13	7,136	89.4
2015	CSM16154.PRA	6/02/2016	6/24/2016	7,990	26	6,590	82.7
2015	SMP16153.PR1	6/01/2016	6/24/2016	995	88	513	56.6
2015	SMP16153.PR2	6/01/2016	6/20/2016	998	5	795	80.1
2015	SMP16154.PR3	6/02/2016	6/16/2016	1001	109	300	33.6
Totals				42,956	615	35,680	84.3
2016	BMI17129.PRE	5/09/2017	5/23/2017	7,996	18	7,279	91.2
2016	BMI17129.PRD	5/09/2017	5/25/2017	7,998	7	7,790	97.5
2016	BMI17143.PRC	5/23/2017	6/09/2017	7,981	32	7,714	97.0
2016	BMI17143.PRB	5/23/2017	6/12/2017	7,995	24	7,633	95.8
2016	BMI17144.PRA	5/24/2017	6/19/2017	7,995	46	7,633	96.0
2016	SMP17128.PR1	5/08/2017	5/23/2017	600	0	538	89.7
2016	SMP17129.PR2	5/09/2017	5/25/2017	600	0	579	96.5
2016	SMP17144.PR3	5/24/2017	6/09/2017	598	0	568	95.0
2016	SMP17144.PR4	5/24/2017	6/12/2017	601	0	581	96.7
2016	SMP17144.PR5	5/24/2017	6/19/2017	600	2	570	95.3
Totals				42,964	129	40,885	95.4
2017	BMI2018128PRE	5/08/2018	5/23/2018	7,999	24	6,681	83.5
2017	BMI2018128PRD	5/08/2018	5/25/2018	7,997	11	6,957	87.0
2017	BMI2018149PRC	5/29/2018	6/11/2018	7,997	6	7,435	93.0
2017	BMI2018150PRB	5/30/2018	6/14/2018	7,997	15	6,916	86.5
2017	BMI2018151PRA	5/31/2018	6/20/2018	7,994	16	6,725	84.1
2017	SMP2018129002	5/09/2018	5/23/2018	599	4	508	84.8
2017	SMP2018129001	5/09/2018	5/25/2018	597	1	524	87.8
2017	SMP2018149PR3	5/29/2018	6/11/2018	599	1	556	92.8

Brood Year	Tag File	Tagging Date	Release Date	# Tagged	# of Tags Recovered from Facility Mortalities	# of Unique Detections	% Detected
2017	SMP2018149PR4	5/29/2018	6/14/2018	597	0	510	85.4
2017	SMP2018150PR5	5/30/2018	6/20/2018	597	0	505	84.6
Totals				42,973	78	37,317	86.8
2018	BMI2019128PRE.XML	5/8/2019	5/22/2019	7,998	26	7240	90.5
2018	BMI2019128PRD.XML	5/8/2019	5/24/2019	8,001	61	7387	92.3
2018	BMI2019148PRC.XML	5/28/2019	6/10/2019	7,996	19	6743	84.3
2018	BMI2019149PRB.XML	5/29/2019	6/13/2109	7,998	19	7314	91.4
2018	BMI2019150PRA.XML	5/30/2019	6/17/2019	7,999	15	7665	95.8
2018	SMP2019127001.XML	5/7/2019	5/22/2019	600	2	580	96.7
2018	SMP2019128002.XML	5/8/2019	5/24/2019	599	4	577	96.3
2018	SMP2019148003.XML	5/28/2019	6/10/2019	599	0	568	94.8
2018	SMP2019149004.XML	5/29/2019	6/13/2109	599	2	580	96.8
2018	SMP2019149005.XML	5/29/2019	6/17/2019	601	2	598	99.5
Totals				42,990	150	39,252	91.3
2020	BMI-2021-125-RCE.XML	5/5/2021	5/24/2021	8,005	11	7,565	94.5
2020	BMI-2021-124-RCD.XML	5/4/2021	5/27/2021	7,969	24	7,630	95.7
2020	BMI-2021-138-RCC.XML	5/18/2021	6/9/2021	7,988	27	7,635	95.6
2020	BMI-2021-138-RCB.XML	5/19/2021	6/14/2021	7,960	25	7,620	95.7
2020	BMI-2021-139-RCA.XML	5/19/2021	6/17/2021	7,983	55	7,732	96.9
2020	SMP-2021-124-PR1.XML	5/4/2021	5/27/2021	574	1	569	99.1
2020	SMP-2021-125-PR2.XML	5/5/2021	5/24/2021	597	1	583	97.7
2020	SMP-2021-145-PR3.XML	5/25/2021	6/9/2021	600	2	585	97.5
2020	SMP-2021-145-PR4.XML	5/25/2021	6/14/2021	600	2	589	98.2
2020	SMP-2021-145-PR5.XML	5/25/2021	6/17/2021	597	4	556	93.1
Totals				42,873	152	41,064	Totals
There was no release of any PIT-tagged fish from Priest Rapids for Brood Year 2019							

10.0 Adult Fish Pathogen Monitoring

At spawning, a portion of the adult fall Chinook broodstock are sampled for infectious hemotopietic necrosis virus (IHNV), infectious pancreatic necrosis virus (IPNV), viral hemorrhagic septicemia virus (VHSV), paramyxovirus, aquaraviruses, as well as *Renibacterium salmoninarum*, the causative agent for bacterial kidney disease (BKD). Viral and bacterial screening included sampling the ovarian fluid and kidney/spleen for pathogens. All results of viral testing since 1991 were classified as negative (Table 18).

Table 18 Viral inspections of fall Chinook Salmon broodstock at Priest Rapids Hatchery, Return Years 1991-2020.

Year	Date(s)	Stock	Life stage	Ovarian Fluid	Kidney/Spleen	Results
1991	28-Oct, 4, 13-Nov	Priest Rapids	Adult	150	60	Negative
1992	2,9-Nov	Priest Rapids	Adult	150	60	Negative
1993	25-Oct, 1-Nov	Priest Rapids	Adult	150	60	Negative
1994	7-Nov	Priest Rapids	Adult	60	60	Negative
1995	9,13,19,21-Nov	Priest Rapids	Adult	160	160	Negative
1996	17-Nov	Priest Rapids	Adult	60	60	Negative
1997	17-Nov	Priest Rapids	Adult	60	60	Negative
1998	16-Nov	Priest Rapids	Adult	60	60	Negative
1999	8-Nov	Priest Rapids	Adult	60	60	Negative

Year	Date(s)	Stock	Life stage	Ovarian Fluid	Kidney/Spleen	Results
2000	13-Nov	Priest Rapids	Adult	60	60	Negative
2001	13-Nov	Priest Rapids	Adult	60	60	Negative
2002	13-Nov	Priest Rapids	Adult	60	60	Negative
2003	17-Nov	Priest Rapids	Adult	60	60	Negative
2004	8-Nov	Priest Rapids	Adult	60	60	Negative
2005	14-Nov	Priest Rapids	Adult	60	60	Negative
2006	6-Nov	Priest Rapids	Adult	60	60	Negative
2007	5-Nov	Priest Rapids	Adult	60	60	Negative
2008	3-Nov	Priest Rapids	Adult	60	60	Negative
2009	2-Nov	Priest Rapids	Adult	60	60	Negative
2010	15-Nov	Priest Rapids	Adult	60	60	Negative
2011	7,14, 21-Nov	Priest Rapids	Adult	180	180	Negative
2012	5-Nov	Priest Rapids	Adult	60	60	Negative
2013	18-Nov	Priest Rapids	Adult	60	60	Negative
2014	18-Nov	Priest Rapids	Adult	60	60	Negative
2015	11-Nov	Priest Rapids	Adult	60	60	Negative
2016	8-Nov	Priest Rapids	Adult	60	60	Negative
2017	1,3,8-Nov	Priest Rapids	Adult	268	268	Negative
2018	5-Nov	Priest Rapids	Adult	60	60	Negative
2019	4-Nov	Priest Rapids	Adult	60	60	Negative
2020	17-Nov	Priest Rapids	Adult	60	60	Negative

Annual testing for BKD was initiated with the 2008 broodstock to address concerns associated with shipping eyed eggs to Bonneville Hatchery for the USACE RSH production. The risk of BKD was assayed using the enzyme linked immunosorbent assay (ELISA) for *R. salmoninarum* antigen (Elliot 2012). Adult broodstock BKD monitoring in 2020 indicated that 98.3% females (N=60) tested had ELISA values less than an optical density of 0.10 (Below Low) (Table 19). Since 2008, tests have shown very low percentages of fish with values greater than 0.10.

Table 19 ELISA test results to determine risk of bacterial kidney disease of adult female fall Chinook Salmon broodstock at Priest Rapids Hatchery, Return Years 2008-2020.

Year	Stock	N	% < Low	% Low	% Mod	% High
2008	Priest Rapids	60	100.0%	0.0%	0.0%	0.0%
2009	Priest Rapids	60	100.0%	0.0%	0.0%	0.0%
2010	Priest Rapids	60	100.0%	0.0%	0.0%	0.0%
2011	Priest Rapids	135	100.0%	0.0%	0.0%	0.0%
2012	Priest Rapids	60	98.3%	0.0%	1.7%	0.0%
2013	Priest Rapids	60	100.0%	0.0%	0.0%	0.0%
2014	Priest Rapids	60	100.0%	0.0%	0.0%	0.0%
2015	Priest Rapids	60	98.3%	1.7%	0.0%	0.0%
2016	Priest Rapids	60	98.3%	1.7%	0.0%	0.0%
2017 ^a	Priest Rapids	268	99.6%	0.4%	0.0%	0.0%
2018	Priest Rapids	60	98.3%	1.7%	0.0%	0.0%
2019	Priest Rapids	60	100.00	0.0%	0.0%	0.0%
2020	Priest Rapids	60	98.3%	1.7%	0.0%	0.0%

^a Oregon Department of Fish and Wildlife tested 268 adults originating from PRH and incorporated into the Umatilla-John Day Mitigation Program for BKD. These fish were trapped at PRH and then transported and spawned at RSH in early November

10.1 Juvenile Fish Health Inspections

Juvenile fish are inspected for the presence of pathogens and other conditions following ponding (AFS-FHS 2014). The results of the examinations of juveniles from brood years 2010 through 2020 are summarized in Table 20. During 2020, juveniles in all raceway ponds appeared healthy upon release. Historical inspection results for brood years 1995 through 2009 are provided in Appendix C.

Table 20 Juvenile fish health inspections for Priest Rapids Hatchery fall Chinook Salmon, Brood Years 2006-2020.

Date	Stock	Brood Year	Condition
18-Feb-10	Priest Rapids	2009	Coagulated Yolk Syndrome observed in some fish sampled
1-Apr-10	Priest Rapids	2009	Healthy
19-May-10	Priest Rapids	2009	Healthy
25-Mar-11	Priest Rapids	2010	Healthy
18-Apr-11	Priest Rapids	2010	Healthy
06-Jun-11	Priest Rapids	2010	Healthy
01-Mar-12	Priest Rapids	2011	Healthy
26-Apr-12	Priest Rapids	2011	Healthy
24-May-12	Priest Rapids	2011	Healthy
11-Feb-13	Priest Rapids	2012	Healthy
3-Mar-13	Priest Rapids	2012	Healthy
29-Apr-13	Priest Rapids	2012	Healthy
28-May-13	Priest Rapids	2012	Healthy
27-Mar-14	Priest Rapids	2013	Dropout Syndrome present
23-Apr-14	Priest Rapids	2013	Dropout Syndrome present
29-May-14	Priest Rapids	2013	Healthy
26-Feb-15	Priest Rapids	2014	Coagulated Yolk Syndrome observed in some fish sampled
26-Mar-15	Priest Rapids	2014	Healthy
21-Apr-15	Priest Rapids	2014	Healthy
28-May-15	Priest Rapids	2014	Healthy
22-June-15	Priest Rapids	2014	Columnaris present in some fish sampled from Channel Pond B.
24-Feb-16	Priest Rapids	2015	Healthy
15-Mar-16	Priest Rapids	2015	Coagulated Yolk Syndrome observed in some fish sampled
15-June-16	Priest Rapids	2015	Mild Ich infection but healthy and ready for release
24-Feb-17	Priest Rapids	2016	Presence of bacterial gill disease in Raceway Bank D and E
21-Mar-17	Priest Rapids	2016	Presence of bacterial gill disease in Raceway Pond B2
6-June-17	Priest Rapids	2016	Mild Ich infection in Channel Ponds A, B, C
21-Mar-18	Priest Rapids	2017	Healthy
19-Apr-18	Priest Rapids	2017	Bacterial gill disease present in Raceway Pond C4
7-May-18	Priest Rapids	2017	Bacterial gill disease present in Raceway Ponds C2 and C3
17-May-18	Priest Rapids	2017	Re-examination of Raceway Ponds C2 and C3 found fish healthy
17-May-18	Priest Rapids	2017	Pre-release examination of Raceway Ponds D and E found fish healthy C2 and C3 found fish healthy
6-June-18	Priest Rapids	2017	Pre-release examination of Raceway Ponds A and B found fish healthy

Date	Stock	Brood Year	Condition
2-Feb-19	Priest Rapids	2018	Examinations of Raceway Banks C, D, E resulted from reports of elevated mortalities. Some fish were found to appear thin and pin-headed. Results of internal necropsies were within normal limits.
5-May-19	Priest Rapids	2018	Pre-release examinations of Raceway Pond E found fish healthy
5-May-19	Priest Rapids	2018	Pre-release examinations of Raceway Pond D resulted in no significant findings of diseases however elevated mortalities were observed. Mortalities examined showed lower levels of coelomic fat and ingesta in GI tracts compared to live fish examined.
6-June-19	Priest Rapids	2018	Pre-release examinations of Raceway Ponds A, B, and C found very low levels of bacterial gill disease
19-Mar-20	Priest Rapids	2019	Examinations of Raceway Bank A resulted from reports of flashing with minimal increase in observed mortality. Some fish examined revealed moderate infestation of <i>Trichodina sp.</i>
28-Apr-20	Priest Rapids	2019	Examinations of Raceway Banks A, B, C, D, E resulted from reports of elevated mortalities. Some fish were found to appear thin and pin-headed. Results of internal necropsies were within normal limits.
20-May-20	Priest Rapids	2019	Pre-release examination of Raceway Ponds D and E found fish generally healthy.
4-Jun-20	Priest Rapids	2019	Pre-release examination of Raceway Ponds A, B, and C found fish generally healthy.
21-May-21	Priest Rapids	2020	Pre-release examination of Raceway Ponds D and E found fish generally healthy.
7-June-21	Priest Rapids	2020	Pre-release examination of Raceway Ponds A, B, and C found fish generally healthy.

11.0 Redd Survey

Fall Chinook Salmon redd surveys were performed in the Hanford Reach during 2020 by staff with Mission Support Alliance under contract with the United States Department of Energy. WDFW M&E staff performed fall Chinook Salmon redd surveys in the PRH discharge channel during 2020.

11.1 Hanford Reach Aerial Redd Counts

Aerial redd counts in the Hanford Reach were performed by Mission Support Alliance on October 19, November 17, and 24 during 2020 (USDOE In Press). The peak fall Chinook Salmon redd count for the Hanford Reach in 2020 was 10,150 (Table 21). Redd counts should be considered an index of the total number of redds in the Hanford Reach. Redds may not be visible during flights due to wind, turbidity, ambient light, and depth. It is reported that viewing conditions for all surveys were good to excellent. The first survey occurred on a Monday and the remaining two occurred on a Tuesday. Total discharge from Priest Rapids Dam ranged from 52 to 163 kcfs during the eight-hour period prior to the surveys. None of the surveys took advantage of the low Sunday outflows at Priest Rapids Dam when flows were lowered to nearly 47 kcfs in conjunction with the Vernita Bar Settlement Agreement redd surveys performed by Grant PUD and WDFW.

Table 21 Summary of fall Chinook Salmon peak redd counts for the 1948-2020 aerial surveys in the Hanford Reach, Columbia River.

Year	Redds	Year	Redds	Year	Redds	Year	Redds	Year	Redds
1948	787	1963	1,254	1978	3,028	1993	2,863	2008	5,588
1949	313	1964	1,477	1979	2,983	1994	5,619	2009	4,996
1950	265	1965	1,789	1980	1,487	1995	3,136	2010	8,817
1951	297	1966	3,101	1981	4,866	1996	7,618	2011	8,915
1952	528	1967	3,267	1982	4,988	1997	7,600	2012	8,368
1953	139	1968	3,560	1983	5,290	1998	5,368	2013	17,398
1954	160	1969	4,508	1984	7,310	1999	6,068	2014	15,951
1955	60	1970	3,813	1985	7,645	2000	5,507	2015	20,678
1956	75	1971	3,600	1986	8,291	2001	6,248	2016	13,268
1957	525	1972	876	1987	8,616	2002	8,083	2017	8,646
1958	798	1973	2,965	1988	8,475	2003	9,465	2018	5,429
1959	281	1974	728	1989	8,834	2004	8,468	2019	7,899
1960	258	1975	2,683	1990	6,506	2005	7,891	2020	10,150
1961	828	1976	1,951	1991	4,939	2006	6,508		
1962	1,051	1977	3,240	1992	4,926	2007	4,023		
Mean (2011 – 2020)									11,195

11.2 Redd Distribution

The main spawning areas observed during the 2020 counts were located near Vernita Bar and among Islands 4-6 (Table 22 and Figure 8). Historical redd counts by location from 2001 through 2020 are included in Appendix D of this report.

Table 22 Number of fall Chinook Salmon redds counted in different reaches on the Hanford Reach area of the Columbia River during aerial counts, October 2020 through November 2020. (Data provided by Mission Support Alliance).

General Location	Start KM	End KM	Total Length	19-Oct	17-Nov	24-Nov	Max Count	Avg. Redd Per River KM
Islands 17-21	545	558	13	0	0	0	0	0
Islands 11-16	558	573	15	2	484	507	507	34
Islands 8-10	587	593	6	0	341	524	524	87
Near Island 7	593	594	1	0	550	650	650	650
Island 6 (lower half)	594	599	5	2	1,310	1,310	1,310	262
Island 4, 5 and upper 6	599	602	3	2	1,562	1,562	1,562	521
Near Island 3	602	604	2	0	650	800	800	400
Near Island 2	604	606	2	0	1,000	1,100	1,100	550
Near Island 1	606	608	2	12	100	100	100	50
Near Coyote Rapids	608	619	11	0	40	70	70	6
Midway (China Bar)	620	630	10	0	20	20	20	2
Near Vernita Bar	630	635	5	13	3,500	3,500	3,500	700
Near Priest Rapids Dam	635	638	3	0	7	7	7	2
Total	--	--	--	31	9,564	10,150	10,150	--




Figure 8 Distribution of fall Chinook Salmon redd counts by location for the 2020 aerial surveys in the Hanford Reach, Columbia River. (Data provided by Mission Support Alliance).

11.3 Spawn Timing

Based on aerial redd counts and Vernita Bar spawning ground surveys, fall Chinook Salmon spawning in the Hanford Reach during 2020 began in late October and ended in late November. River temperatures below Priest Rapids Dam varied from 15.7°C (October 20) to 10.9°C (November 25) during the spawning period which is typical to that of previous years.

11.4 Escapement

The estimated total escapement of fall Chinook Salmon to the Hanford Reach for the 2020 return year was 74,834 fish (Table 23). The historical mean and median escapement for 1991 through 2020 was 73,779 and 58,779 fish, respectively (Table 24). The estimated adult Chinook Salmon per redd was calculated by dividing the adult escapement to the Hanford Reach by peak number of redds reported in the redd survey. The estimated annual escapements to the Hanford Reach were not adjusted for pre-spawn mortality. For 2020 the estimated seven fish per redd versus the historical mean of 9 fish.

Table 23 Calculation of escapement estimates for fall Chinook Salmon in the Hanford Reach, Columbia River 2020.

Count Source	Return Year 2020		
	Adult	Jack	Total
McNary Ladder Counts	186,097	27,171	213,268
Adjusted Priest Rapids Adult Passage ¹	30,924	2,914	33,838
Ice Harbor Adult Passage	29,172	9,251	38,423
Prosser Adult Passage	2,061	118	2,179
Priest Rapids Discharge Channel	54	3	57
Priest Rapids Hatchery	34,694	4,139	38,833
Wanapum Tribal Fishery	12	0	12
Ringold Springs Hatchery	6,737	354	7,091
Yakima River Escapement (Below Prosser)	473	31	504
Yakima River Sport Harvest	230	55	285
Hanford Sport Harvest	14,651	1,395	16,046
Angler Broodstock Collection	1,164	2	1,166
Total Non-Hanford Reach Escapement	120,172	18,262	138,434
Hanford Reach Escapement	65,925	8,909	74,834

¹ Gross passage count reduced 34.15% to correct for estimated over counts resulting from fallbacks and re-ascension. The adjustments to adult fish passage were estimated by analysis of the PIT tag detections at PIT tag arrays located in the adult fish ways of the Priest Rapids Dam adult fishway and the discharge channel for Priest Rapids Hatchery.

Table 24 Escapement for fall Chinook Salmon in the Hanford Reach, Return Years 1991-2020.

Return Year	# Fish per Redd	Redds	Total Escapement ¹
1991	11	4,939	52,196
1992	9	4,926	41,952
1993	13	2,863	37,347
1994	11	5,619	63,103
1995	18	3,136	55,208
1996	6	7,618	43,249
1997	6	7,600	43,493
1998	7	5,368	35,393
1999	5	6,068	29,812
2000	9	5,507	48,020
2001	10	6,248	59,848
2002	10	8,083	84,509
2003	11	9,465	100,508
2004	10	8,468	87,696
2005	9	7,891	71,967
2006	8	6,508	51,701
2007	6	4,018	22,272
2008	5	5,618	29,058
2009	7	4,996	36,720
2010	10	8,817	87,016
2011	8	8,915	75,256
2012	7	8,368	57,710
2013	10	17,398	174,651

Return Year	# Fish per Redd	Redds	Total Escapement ¹
2014	12	15,951	183,749
2015	13	20,678	266,327
2016	9	13,268	116,388
2017	9	8,646	73,759
2018	9	5,429	46,624
2019	8	7,899	65,991
2020	7	10,150	74,834
Mean	9	8,015	73,879
Median	9	7,609	58,779

¹ Escapement includes adults and jacks

11.5 Hatchery Discharge Channel Redd Counts

The M&E team observed 28 redds during the survey in the PRH discharge channel on December 7, 2020. Similar to historical observations, the majority of spawning activity was located in a 200-meter section of the discharge channel just below the volunteer trap. We observed superimposition occurring during multiple brief site visits during November; thus, making it difficult to determine the total number of redds in the formal survey. Redds were observed throughout the suitable spawning habitat in the discharge channel (to top 200-meter section). Viewing conditions during the surveys were good.

12.0 Carcass Surveys

Prior to 2010, the carcass surveys in the Hanford Reach were generally performed by two boat crews of two staff operating seven days a week. Beginning in 2010, with support of the PRH M&E Program, the effort was increased to three boats with a three-person crew operating seven days per week. The extra staffing was necessary to maintain the overall sampling efficiency given the additional effort required to pull otoliths from fish sampled and achieve hatchery M&E objectives. The sampling goal for obtaining minimum desirable numbers of CWTs was 10% of the escapement.

Carcass surveys were performed from November 2 through December 12, 2020. All recovered carcasses were screened for the presence of a CWT which was collected if present. Roughly 33% of the fish recovered were sampled (i.e., random systematic 1:3 rate) for scales (age), otoliths, gender, length, and egg retention. All carcasses recovered were chopped in half after sampling to prevent the chance of double sampling during subsequent surveys.

Similar to methods used since 2010, the carcass survey crews recorded the sections in which carcasses were recovered in the Hanford Reach and adjacent areas. The Hanford Reach survey was divided into Sections 1 through 5 (Figure 9). The Priest Rapids Pool was designated as Section 6. The PRH discharge channel and the area of the Columbia River immediately below the discharge channel were designated as Sections 7 and 8, respectively. The fall Chinook Salmon carcasses recovered in Section 8 were likely wash outs from the hatchery discharge channel.

- Section 1. Priest Rapids Dam to Vernita Bridge (14 km)
- Section 2. Vernita Bridge to Island 2 (19 km)
- Section 3. Island 2 to Powerline Towers at Hanford town site (21 km)
- Section 4. Power line Towers to Wooded Island (21 km)

- Section 5. Wooded Island to Interstate 182 Bridge (19 km)
- Section 6. Priest Rapids Pool (34 km)
- Section 7. Priest Rapids Hatchery discharge channel (0.5 km)
- Section 8. Columbia River at the mouth of the Hatchery discharge channel (0.5 km)




Figure 9 Location of aerial redd index areas (green area numbers) and river boat carcass survey sections in the Hanford Reach.

12.1 Hanford Reach Carcass Survey: Section 1 – 5

Staff recovered 4,669 fall Chinook Salmon carcasses in the Hanford Reach in 2020; equating to 6.2% of the estimated fall Chinook Salmon escapement (Table 25). The annual number of fall Chinook Salmon carcass recovered in the Hanford Reach for the period of 1991 through 2020 is provided in Appendix E.

Table 25 Numbers and percentage of total escapement of fall Chinook Salmon carcasses surveyed (1:1 rate) for coded-wire tags within each survey section on the Hanford Reach, Return Years, 2010-2020.

Return Year	# 1		# 2		# 3		# 4		# 5		Total Sampled		Escapement
	N	%	N	%	N	%	N	%	N	%	N	%	
2010	1,832	2.1	519	0.6	3,129	3.6	3,362	3.9	937	1.1	9,779	11.2	87,016
2011	1,581	2.1	160	0.2	2,606	3.5	2,622	3.5	1,422	1.9	8,391	11.1	75,256
2012	1,091	1.9	149	0.3	1,685	2.9	2,213	3.8	1,676	2.9	6,814	11.8	57,715
2013	2,182	1.2	1,973	1.1	2,844	1.6	3,774	2.2	2,298	1.3	13,071	7.5	174,651
2014	2,682	1.5	1,142	0.6	5,544	3.0	4,573	2.5	2,815	1.5	16,756	9.1	183,680
2015	2,913	1.1	823	0.3	6,187	2.3	5,868	2.2	1,947	0.7	17,738	6.7	266,346
2016	1,141	1.0	513	0.4	2,796	2.4	2,977	2.6	1,459	1.3	8,886	7.6	116,388
2017	1,098	1.5	346	0.5	1,275	1.7	1,850	2.5	1,022	1.4	5,591	7.6	73,759

Return Year	# 1		# 2		# 3		# 4		# 5		Total Sampled		Escapement
	N	%	N	%	N	%	N	%	N	%	N	%	
2018	635	1.4	113	0.2	920	2.0	720	1.5	383	0.8	2,771	5.9	46,624
2019	904	1.4	202	0.3	1,028	1.6	1,223	1.9	655	1.0	4,012	6.1	65,991
2020	1,190	1.6	226	0.3	1,609	2.2	1,285	1.7	359	0.5	4,669	6.2	74,834
Mean	1,568	2.0	561	0.4	2,693	2.4	2,770	2.6	1,361	1.3	8,953	8.3	111,115

The survey effort was not equal for each section. Sections 3 and 4 were surveyed the most because these sections generally contain the largest number of carcasses (Table 26). As each season progresses, crews focused their effort in sections that provided greater chances to recover carcasses.

Table 26 Number of carcass surveys conducted by section in the Hanford Reach, Return Years 2010-2020.

Return Year	# 1	# 2	# 3	# 4	# 5	Total
2010	21	6	26	26	11	90
2011	33	5	38	29	13	118
2012	19	4	26	28	24	101
2013	18	15	16	17	13	79
2014	23	17	30	31	24	125
2015	23	8	35	37	13	116
2016	18	11	29	27	15	100
2017	19	14	30	31	17	111
2018	20	9	31	22	17	99
2019	24	9	31	25	21	110
2020	25	8	33	26	13	105
Mean	22	10	30	27	16	105

12.2 Proportion of Escapement Sampled: Section 1-5

The spawning escapement for sections 1 through 5 was estimated by the proportion of redds counted in aerial surveys to the estimated escapement of natural spawners in the Hanford Reach (see Section 14 - Redd Surveys). The calculations for estimating the escapement to the Hanford Reach are given in Appendix F.

We have identified through the carcass bias assessment that an unknown number of carcasses drift into downstream sections after spawning. The recovery of these carcasses may confound the estimate of the spawning escapement sampled by section as shown in Table 27.

As seen in prior years there were no redds identified in Section 5, but hundreds of carcasses were recovered in that section. It is likely that sections 1 and 3, having the greatest number of redds and largest spawning escapement, end up with a net loss of carcasses to downstream sections.

Table 27 Number of redds and carcasses, total spawning escapement, and proportion of escapement sampled for fall Chinook Salmon in Section 1 through 5 of the Hanford Reach, Return Year 2020.

Survey Section	Total Number of Redds	Total Number of Carcasses	Spawning Escapement ¹	Proportion of Escapement Sampled
1	3,527	1,190	26,004	0.046
2	170	226	1,253	0.180
3	5,946	1,609	43,839	0.037
4	507	1,285	3,738	0.344
5	0	359	0	--
Total	10,150	4,669	74,834	0.062

¹ Calculated based on proportion of redds by section

12.3 Carcass Distribution and Origin

Two methods were used to estimate the origin of carcasses recovered in the Sections 1 – 5. The first method includes the expansion of pooled CWT recoveries using juvenile tag rates and survey sample rate. The second method includes calculating the proportion of combined hatchery marks (i.e., otolith mark, adipose clips, and CWTs) in the demographic sample of the population. Estimates for both methods are given for the 2012 - 2020 adult returns: these years include otolith marks for all ages of PRH origin fish.

The assumption was made that all Chinook Salmon not accounted by hatchery origin CWT expansions were of natural origin. This assumption may underestimate the number of hatchery carcasses recovered in the annual surveys. We have compelling evidence to suggest this is the case with annual returns to PRH prior to return year 2014. The expansion of CWT recoveries suggests the proportion of hatchery origin fish was highest in section-1 and ranged from 0.178 to 0.252 among the five sections (Table 28). This estimate is higher than observed since return year 2013.

The second estimate of origin of carcasses recovered is based on the proportion of hatchery marked to non-marked fish recovered in the demographic sample which was every third fish. For this method, we assume that all hatchery origin carcasses recovered are marked in some manner (e.g., otolith marks, CWT, and adipose clips) and that we can accurately detect these marks and tags.

PRH has marked their entire juvenile releases with thermal marks on the otoliths beginning with progeny of brood year 2007: Hence, all PRH origin returns since 2013 were otolith marked. The age-6 PRH origin fish were not otolith marked during return year 2012. However, since there were no age-6 fish recovered in the carcass surveys or at PRH, it is assumed that few, if any PRH origin age-6 fish spawned in the Hanford Reach. Adipose clipped Chinook Salmon without a CWT and without a thermal otolith mark were classified as strays from other hatcheries. The natural origin fish were identified by either a Hanford Reach origin CWT or by the presence of an adipose fin and the absence of an otolith mark. The demographic sample data suggests the proportion of hatchery origin fish was highest in section-1 and ranged from 0.243 to 0.297 among the five sections (Table 29). These are the highest proportions of hatchery fish observed since return year 2013.

Table 28 Numbers of natural and hatchery origin fall Chinook Salmon carcasses sampled within Section 1 through 5 of Hanford Reach based on expansions of coded-wire tag recoveries, Return Years 2010-2020.

Return Year	Hanford Reach Sections						Total
	Origin	# 1	# 2	# 3	# 4	# 5	
2010	Natural	1,751	473	3,020	3,242	909	9,395
	Hatchery	81	46	116	125	28	396
	Proportion Hatchery	0.044	0.089	0.037	0.037	0.030	0.040
2011	Natural	1,350	155	2,520	2,475	1,347	7,847
	Hatchery	231	5	86	147	75	544
	Proportion Hatchery	0.146	0.031	0.033	0.056	0.053	0.065
2012	Natural	1,142	149	1,526	2,081	1,510	6,408
	Hatchery	49	0	159	132	166	506
	Proportion Hatchery	0.041	0.000	0.094	0.060	0.099	0.073
2013	Natural	1,572	1,587	2,433	2,895	1,748	10,235
	Hatchery	610	386	411	879	550	2,836
	Proportion Hatchery	0.280	0.196	0.145	0.233	0.239	0.217
2014	Natural	2,469	1,072	5,264	4,329	2,703	15,838
	Hatchery	213	70	280	244	112	918
	Proportion Hatchery	0.079	0.061	0.050	0.053	0.040	0.055
2015	Natural	2,654	709	5,745	5,490	1,858	16,456
	Hatchery	259	114	442	378	89	1,282
	Proportion Hatchery	0.089	0.139	0.071	0.064	0.046	0.072
2016	Natural	1,108	256	2,585	2,866	684	8,111
	Hatchery	162	33	257	211	111	775
	Proportion Hatchery	0.142	0.064	0.092	0.071	0.076	0.087
2017	Natural	1,015	260	1,173	1,648	863	4,958
	Hatchery	83	86	102	202	175	649
	Proportion Hatchery	0.076	0.249	0.080	0.109	0.169	0.116
2018	Natural	578	101	881	694	355	2,608
	Hatchery	57	12	39	26	28	163
	Proportion Hatchery	0.090	0.106	0.043	0.037	0.073	0.059
2019	Natural	848	171	998	1,193	609	3,820
	Hatchery	56	31	30	30	46	192
	Proportion Hatchery	0.062	0.152	0.029	0.024	0.070	0.048
2020	Natural	890	172	1,209	1,055	295	3,622
	Hatchery	300	54	400	230	64	1,047
	Proportion Hatchery	0.252	0.240	0.249	0.179	0.178	0.224
Mean	Proportion Hatchery	0.118	0.121	0.084	0.084	0.098	0.096

Table 29 Origin of Chinook Salmon carcasses recovered in the Hanford Reach by section based on recoveries of marked and unmarked carcasses within the biological sample, Return Years 2012-2020.

Year	Origin	# 1	# 2	# 3	# 4	# 5	Total	Proportion of Sample
2012 Biological sample Rate 1:4 N = 1,609	PRH ¹	23	2	26	18	38	107	0.067
	Other Hatchery ²	10	2	25	45	22	104	0.065
	Total Hatchery	33	4	51	63	60	211	0.131
	Natural ³	228	30	347	460	333	1,398	0.869
	Proportion Hatchery	0.126	0.118	0.128	0.120	0.153	0.131	
2013 ^a Biological sample rate = 1:5 and then randomly sub- sampled, N = 712	PRH ¹	32	19	34	30	32	147	0.206
	Other Hatchery ²	6	3	16	21	6	52	0.073
	Total Hatchery	38	22	50	51	38	199	0.279
	Natural ³	76	84	113	155	85	513	0.721
	Proportion Hatchery	0.333	0.208	0.307	0.248	0.309	0.279	
2014 ^a Biological sample rate = 1:5 and then randomly sub- sampled, N = 2,426	PRH ¹	37	7	45	35	11	135	0.056
	Other Hatchery ²	12	5	16	32	18	83	0.034
	Total Hatchery	49	12	61	67	29	218	0.090
	Natural ³	347	142	711	612	396	2,208	0.910
	Proportion Hatchery	0.124	0.078	0.079	0.099	0.068	0.090	
2015 Biological sample rate = 1:7 N = 2,485	PRH ¹	47	12	61	55	13	188	0.076
	Other Hatchery ²	6	2	17	20	7	52	0.021
	Total Hatchery	53	14	78	75	20	240	0.097
	Natural ³	346	101	792	752	254	2,245	0.903
	Proportion Hatchery	0.133	0.122	0.090	0.091	0.073	0.097	
2016 Biological sample rate = 1:5 N = 1,743	PRH ¹	27	12	42	22	10	113	0.066
	Other Hatchery ²	9	6	31	23	13	82	0.048
	Total Hatchery	36	18	73	45	23	195	0.114
	Natural ³	182	80	465	534	257	1,518	0.886
	Proportion Hatchery	0.165	0.184	0.136	0.078	0.082	0.114	
2017 Biological sample rate = 1:3 N = 1,813	PRH ¹	42	19	21	19	16	117	0.065
	Other Hatchery ²	7	2	4	14	6	33	0.018
	Total Hatchery	49	21	25	33	22	150	0.083
	Natural ³	311	86	391	564	311	1,663	0.917
	Proportion Hatchery	0.136	0.196	0.060	0.055	0.066	0.083	
2018 Biological sample rate = 1:2 N = 1,325	PRH ¹	28	6	11	11	6	63	0.047
	Other Hatchery ²	7	2	8	10	2	29	0.022
	Total Hatchery	35	8	19	21	8	92	0.069
	Natural ³	245	72	422	318	177	1,236	0.931
	Proportion Hatchery	0.127	0.100	0.043	0.062	0.043	0.069	
2019 Biological sample Rate = 1:2 M = 1,887	PRH ¹	57	10	44	35	31	177	0.094
	Other Hatchery ²	11	1	8	16	8	44	0.023
	Total Hatchery	68	11	52	51	39	221	0.117
	Natural ³	371	85	432	537	241	1,666	0.883

Year	Origin	# 1	# 2	# 3	# 4	# 5	Total	Proportion of Sample
	Proportion Hatchery	0.155	0.115	0.107	0.087	0.139	0.117	
2020 Biological sample Rate = 1:3 M = 1,446	PRH ¹	91	16	114	73	16	310	0.214
	Other Hatchery ²	19	1	19	43	10	92	0.064
	Total Hatchery	110	17	133	116	26	402	0.279
	Natural ³	260	53	349	309	73	1,044	0.722
	Proportion Hatchery	0.297	0.243	0.276	0.273	0.263	0.278	
Mean Proportion	PRH ¹	0.140	0.117	0.091	0.066	0.085	0.094	
	Other Hatchery ²	0.032	0.029	0.036	0.051	0.039	0.039	
	Total Hatchery	0.172	0.146	0.127	0.117	0.124	0.133	
	Natural ³	0.828	0.854	0.873	0.883	0.876	0.867	

^a Estimate of origin based on random sub-sample of biological sample.

¹ Priest Rapids Hatchery fish were identified by either the presence Priest Rapids Hatchery otolith mark or coded-wire tag

² Other hatchery strays were identified as adipose clipped Chinook Salmon without a Priest Rapids Hatchery coded-wire tag and without a thermal otolith mark or by the presence of other hatchery coded-wire tags.

³ Natural origin fish were identified by either a Hanford Reach origin coded-wire tag or by the presence of an adipose fin and the absence of an otolith mark.

12.4 Priest Rapids Dam Pool Carcass Survey: Section 6

Staff performed four surveys in Section 6 between November 5 and December 1 during return year 2020 (Table 30).

12.5 Number Sampled: Section 6

Survey crews recovered 71 Chinook Salmon in Section 6 during return year 2020 (Table 30). All fish recovered were scanned for the presence of a CWT. The locations of carcass recoveries in the lower portion of the pool suggest that carcasses drift downstream of the spawning areas below Wanapum Dam into deeper water where they are difficult to locate and recover.

12.5.1 Proportion of Escapement Sampled: Section 6

The spawning escapement for Section 6 was calculated by subtracting from the Priest Rapids Dam fall Chinook Salmon passage count, the fall Chinook Salmon passage at Wanapum Dam, tribal and sport harvest of fall Chinook Salmon in the Priest Rapids Dam pool, and the estimated fallback of fall Chinook Salmon at Priest Rapids Dam (Appendix F).

The 2020 fall Chinook Salmon spawning escapement estimate for Section 6 is 840 fish. Overall, 8.4% of the total estimated spawning escapement in Section 6 was sampled for a CWT and gender (Table 30).

Table 30 Carcasses sampled, total spawning escapement and proportion of escapement for fall Chinook Salmon in Section 6 (Priest Rapids Dam Pool), Return Years 2010-2020.

Return Year	# of Surveys	# of Carcasses	Spawning Escapement	Escapement Sampled
2010	8	123	11,121	0.011
2011	7	69	11,362	0.006
2012	4	72	21,919	0.003
2013	7	407	62,237	0.007
2014	7	237	25,179	0.009

Return Year	# of Surveys	# of Carcasses	Spawning Escapement	Escapement Sampled
2015	6	155	38,313	0.004
2016	8	139	13,162	0.011
2017	5	40	1,788	0.022
2018	2	57	2,876	0.020
2019	6	90	5,476	0.016
2020	4	71	840	0.084
Mean	6	133	17,661	0.018

12.5.2 Carcass Origin: Section 6

Similar to those methods described in detail in the previous section, the carcasses included in the 1:3 demographic sample were identified as hatchery origin based on a combination of hatchery marks and tags (i.e., otoliths marks, adipose clips, and CWTs). Natural origin carcasses were identified by the absence of any hatchery mark or the presence of a natural origin CWT. An estimated 68.2% of the fall Chinook Salmon carcasses recovered in Section 6 were hatchery origin of which most all were PRH origin (Table 31).

Table 31 Origin of fall Chinook Salmon spawning in Section 6 (Priest Rapids Dam Pool), Return Years 2012-2020.

Year	Origin	Total	Proportion of Sample
2012 N = 70	PRH ¹	18	0.257
	Other Hatchery ²	2	0.029
	Total Hatchery	20	0.286
	Natural ³	50	0.714
2013 N = 98	PRH ¹	62	0.633
	Other Hatchery ²	5	0.051
	Total Hatchery	67	0.684
	Natural ³	31	0.316
2014 N = 229	PRH ¹	81	0.354
	Other Hatchery ²	5	0.022
	Total Hatchery	86	0.376
	Natural ³	143	0.624
2015 N = 244	PRH ¹	83	0.535
	Other Hatchery ²	3	0.019
	Total Hatchery	155	0.554
	Natural ³	69	0.446
2016 N = 134	PRH ¹	66	0.475
	Other Hatchery ²	3	0.022
	Total Hatchery	69	0.497
	Natural ³	65	0.503
2017 N = 40	PRH ¹	15	0.375
	Other Hatchery ²	1	0.025
	Total Hatchery	16	0.4
	Natural ³	24	0.6
2018	PRH ¹	8	0.143
	Other Hatchery ²	1	0.018

Year	Origin	Total	Proportion of Sample
N = 56	Total Hatchery	9	0.161
	Natural ³	47	0.839
2019 N = 78	PRH ¹	23	0.295
	Other Hatchery ²	1	0.013
	Total Hatchery	24	0.308
	Natural ³	54	0.692
2020 N = 22	PRH ¹	15	0.652
	Other Hatchery ²	0	0.000
	Total Hatchery	15	0.652
	Natural ³	8	0.348
Proportions	PRH¹		0.413
	Other Hatchery²		0.022
	Total Hatchery		0.435
	Natural³		0.565

¹ Priest Rapids Hatchery fish were identified by either the presence of thermal otolith mark or by the presence of a PRH origin coded-wire tag

² Other hatchery strays were identified as adipose clipped Chinook Salmon without a Priest Rapids Hatchery coded-wire tag and without a thermal otolith mark.

³ Natural origin fish were identified by either a Hanford Reach origin coded-wire tag or by the presence of an adipose fin and the absence of an otolith mark.

12.6 Hatchery Discharge Channel: Section 7 and 8 Carcass Survey

During return year 2020, crews performed three carcass surveys in Section 8 by boat and one carcass survey in Section 7 by foot. It has been observed that many carcasses drift out of the discharge channel under full flow conditions. Performing carcass surveys in the discharge channel when it is at full flow is difficult and dangerous due to poor footing and high velocities. Staff performed the one survey in Section 7 on December 6 when discharge levels in the channel were still high. It is likely a portion of the carcasses may have drifted out of the discharge channel by the date that it was surveyed.

12.7 Number sampled: Section 7 and 8

Survey crews recovered 9 carcasses in Section 7 and 15 in Section 8 (Table 32). All fish recovered were scanned for the presence of a CWT.

Table 32 The number of fall Chinook Salmon carcass surveys within Section 7 (Priest Rapids Hatchery Discharge Channel) and Section 8 (Columbia River at the confluence of the hatchery discharge channel), Return Years 2010-2020.

Return Year	Section 7		Section 8		Total	
	# of Carcasses	# of Surveys	# of Carcasses	# of Surveys	# of Carcasses	# of Surveys
2010	87	1	123	9	210	10
2011	123	2	80	8	203	10
2012	99	3	108	10	207	13
2013	105	3	159	4	264	7
2014	9	1	52	7	61	8
2015	33	1	26	2	59	3
2016	3	1	7	1	10	2

Return Year	Section 7		Section 8		Total	
	# of Carcasses	# of Surveys	# of Carcasses	# of Surveys	# of Carcasses	# of Surveys
2017	9	1	16	1	25	2
2018	3	1	0	2	3	3
2019	10	1	47	4	57	5
2020	9	1	15	3	26	4
Mean	45	1	58	5	102	6

12.7.1 Proportion of Escapement Sampled: Section 7 and 8

The 2020 fall Chinook Salmon spawning escapement index for Sections 7 and 8 is 47 fish (Table 33). The spawning escapement for these Sections was calculated using the expansion factor of 1:1 female/redd ratio and a 0.68:1 male/female sex ratio including jacks, as estimated from the Hanford Reach 2020 escapement. Therefore, the assumption is made that each of the 28 redds represents one female and one male. In the past, we assumed that most of the carcasses recovered in Section 8 drifted downstream from Section 7. It's possible that some portion of post spawned fish from Section 7 may drift downstream into Sections 1 and 2 as well.

Table 33 Number of carcasses sampled, total spawning escapement and proportion of escapement sampled for fall Chinook Salmon within Section 7 (Priest Rapids Hatchery Discharge Channel) and Section 8 (Columbia River at confluence of the hatchery discharge channel), Return Year 2020.

Section	Total Number of Carcasses	Spawning Escapement	Escapement Sampled
# 7	9	47	0.468
# 8	13	0	
Total	22	47	0.468

12.7.2 Carcass Distribution and Origin: Section 7 and 8

The demographic sample rate was set at 1:1 in Section 7 and 1:3 in Section 8 to account for the low numbers of carcasses recovered. As described in detail previously, the carcasses included the demographic sample were identified as hatchery origin based on a combination of hatchery marks and tags (i.e., otoliths marks, adipose clips, and CWTs). Natural origin carcasses were identified by the absence of any hatchery mark or the presence of a natural origin CWT.

It is estimated that 77.3% of fall Chinook Salmon recovered in Sections 7 and 8 during 2020 were hatchery origin of which all were PRH origin (Table 34).

Table 34 The origin of Chinook Salmon carcasses recovered within Section 7 (Priest Rapids Hatchery Discharge Channel) and Section 8 (Columbia River at the confluence of the hatchery discharge channel), Return Years 2012-2020.

Return Year	Origin	Total	Proportion of Sample
2012 N = 70	PRH ¹	18	0.257
	Other Hatchery ²	2	0.029
	Total Hatchery	20	0.286
	Natural ³	50	0.714
2013 N = 33	PRH ¹	28	0.848
	Other Hatchery ²	2	0.061
	Total Hatchery	30	0.909

Return Year	Origin	Total	Proportion of Sample
	Natural ³	3	0.091
2014 N= 5	PRH ¹	3	0.600
	Other Hatchery ²	0	0.000
	Total Hatchery	3	0.600
	Natural ³	2	0.400
2015 N= 59	PRH ¹	19	0.322
	Other Hatchery ²	2	0.034
	Total Hatchery	21	0.356
	Natural ³	38	0.644
2016 N=6	PRH ¹	4	0.667
	Other Hatchery ²	1	0.167
	Total Hatchery	5	0.833
	Natural ³	1	0.167
2017 N=6	PRH ¹	6	0.750
	Other Hatchery ²	0	0.000
	Total Hatchery	6	0.750
	Natural ³	2	0.250
2018 N = 3	PRH ¹	1	0.333
	Other Hatchery ²	0	0.000
	Total Hatchery	1	0.333
	Natural ³	2	0.667
2019 N=31	PRH ¹	35	0.511
	Other Hatchery ²	0	0.000
	Total Hatchery	35	0.511
	Natural ³	33	0.489
2020 N=22	PRH ¹	16	0.727
	Other Hatchery ²	1	0.045
	Total Hatchery	17	0.773
	Natural ³	5	0.227
Means Proportions	PRH¹		0.557
	Other Hatchery²		0.037
	Total Hatchery		0.595
	Natural³		0.405

¹ Priest Rapids Hatchery fish were identified by either the presence of thermal otolith mark or by the presence of a PRH origin coded-wire tag

² Other hatchery strays were identified as adipose clipped Chinook Salmon without a Priest Rapids Hatchery coded-wire tag and without a thermal otolith mark.

³ Natural origin fish were identified by either a Hanford Reach origin coded-wire tag or by the presence of an adipose fin and the absence of an otolith mark.

13.0 Life History Monitoring

Migration timing of hatchery and natural origin Hanford Reach fall Chinook Salmon is estimated from arrival timing at Bonneville Dam based on PIT tag observations at the adult fish ladder for both PRH and Hanford Reach origin fall Chinook Salmon.

Life history characteristics of Hanford Reach fall Chinook Salmon were assessed by examining carcasses on spawning grounds, fish collected or examined at broodstock collection sites, and by reviewing tagging data and fisheries statistics.

For the 2012 - 2020 returns, the origin of fall Chinook Salmon for the comparison of age and length at maturity was based on a combination of hatchery marks and tags (i.e., otolith, adipose clips, and CWT). PRH origin fall Chinook Salmon were identified by either the presence of a thermal otolith mark specific to PRH or by the presence of a PRH origin CWT. Adipose clipped Chinook Salmon without a CWT and without an otolith mark were classified as fish from other hatcheries. The natural origin fish were identified by either a Hanford Reach origin CWT or by the presence of an adipose fin combined with the absence of any hatchery marks. The age composition for both the natural and hatchery origin fall Chinook Salmon recovered in return years 2012 - 2010 were assembled from the carcass recoveries in sections 1-8 of the Hanford Reach.

In order to make coarse comparisons between hatchery and natural origin fish prior to return year 2012, the designation of origin required the assumption that all fish collected in the Hanford Reach, except for those that were of known hatchery origin (e.g., adipose clipped or possessed a CWT), were natural origin. We know this was not the case, but we were not able to identify all the hatchery origin fish in the demographic samples and it was assumed that the majority of the fish sampled in the stream surveys were natural origin.

13.1 Adult Migration Timing

PIT tag observations for both PRH and Hanford Reach natural origin adult fall Chinook Salmon at the PIT tag arrays in the Bonneville Dam adult fish ladders were used to assess arrival timing. The PIT tag observation data was obtained from the PTAGIS website. Arrival date for each unique tagged adult was based on its first observation date and time at Bonneville Dam. The data presented encompasses return years 2010 – 2020. The annual number of adult PIT tag observations at Bonneville Dam varied for both hatchery and natural origin fish as a result of varying size tag groups, smolt to adult survival, and PIT tag detection efficiencies at the adult fishways. Roughly 3,000 juveniles were PIT tagged at PRH annually for release years 2005 – 2010. The annual tag group was roughly 43,000 between 2011 to 2019. The annual tag size for the Hanford Reach natural origin juvenile fall Chinook Salmon have ranged from a high of 22,433 in 2007 to a low of 4,183 in 2013. There was not a tag group for natural origin fish in 2006.

The adult PIT tag detections at Bonneville Dam are useful to compare migration timing between Hanford Reach natural origin and PRH origin fall Chinook Salmon because harvest and other losses upstream of Bonneville Dam reduce the number of potential detections at upstream sites.

The 10th, 50th, and 90th percentiles of the annual migration timing to Bonneville Dam are given in (Table 35). The observation sample size of both groups of PIT tagged fish at Bonneville Dam can be small and therefore, may not be representative of the populations. However, this may be the best migration information currently available.

Table 35 The week that 10%, 50% (median), and 90% of the natural and hatchery origin fall Chinook Salmon passed Bonneville Dam, 2010-2020. Migration timing is based on PIT tag passage of Hanford natural origin and Priest Rapids Hatchery in the adult fish ladder at Bonneville Dam.

Return Year	Origin	Hanford Reach Fall Chinook Migration Time (Date)									
		Priest Rapids Origin					Hanford Reach Natural Origin				
		Age 2	Age 3	Age 4	Age 5	Age 6	Age 2	Age 3	Age 4	Age 5	Age 6
2010	10 th Percentile	28-Aug	26-Aug	--	24-Aug	--	31-Aug	5-Sep	25-Aug	--	--
	50 th Percentile	9-Sep	17-Sep	--	4-Sep	--	21-Sep	17-Sep	9-Sep	--	--
	90 th Percentile	15-Sep	24-Sep	--	6-Sep	--	4-Oct	6-Oct	15-Sep	--	--
	N	5	20	0	3	0	8	22	18	0	0
2011	10 th Percentile	8-Aug	3-Sep	23-Aug	--	--	--	4-Sep	24-Aug	4-Aug	--
	50 th Percentile	8-Sep	20-Sep	8-Sep	--	--	--	4-Sep	10-Sep	30-Aug	--
	90 th Percentile	21-Sep	25-Sep	21-Sep	--	--	--	10-Sep	2-Oct	1-Sep	--
	N	6	7	10	0	0	0	2	65	3	--
2012	10 th Percentile	31-Aug	6-Sep	13-Sep	7-Sep	--	14-Sep	4-Sep	28-Aug	27-Aug	--
	50 th Percentile	16-Sep	11-Sep	13-Sep	7-Sep	--	23-Sep	16-Sep	5-Sep	8-Sep	--
	90 th Percentile	27-Sep	21-Sep	19-Sep	7-Sep	--	10-Oct	26-Sep	21-Sep	19-Sep	--
	N	7	13	2	1	0	10	11	19	26	0
2013	10 th Percentile	24-Aug	28-Aug	25-Aug	--	--	11-Sep	2-Sep	2-Sep	9-Aug	--
	50 th Percentile	8-Sep	9-Sep	3-Sep	--	--	11-Sep	22-Sep	9-Sep	27-Aug	--
	90 th Percentile	18-Sep	22-Sep	15-Sep	--	--	11-Sep	10-Oct	19-Sep	2-Oct	--
	N	40	55	16	0	0	1	29	22	10	--
2014	10 th Percentile	6-Sep	4-Sep	5-Sep	--	--	24-Sep	10-Sep	3-Sep	29-Aug	--
	50 th Percentile	16-Sep	13-Sep	12-Sep	--	--	25-Sep	11-Sep	12-Sep	1-Sep	--
	90 th Percentile	28-Sep	25-Sep	23-Sep	--	--	1-Oct	28-Sep	26-Sep	15-Sep	--
	N	175	228	50	0	0	3	4	62	5	0
2015	10 th Percentile	16-Oct	8-Sep	25-Aug	14-Sep	--	--	10-Sep	30-Aug	29-Aug	27-Sep
	50 th Percentile	16-Oct	21-Sep	6-Sep	26-Sep	--	--	20-Sep	10-Sep	9-Sep	27-Sep
	90 th Percentile	16-Oct	9-Oct	18-Sep	26-Sep	--	--	1-Oct	25-Sep	25-Sep	27-Sep
	N	1	345	323	2	0	0	5	13	32	1
2016	10 th Percentile	--	30-Aug	8-Aug	14-Aug	--	--	21-Sep	28-Aug	31-Aug	--
	50 th Percentile	--	13-Sep	7-Sep	1-Sep	--	--	21-Sep	10-Sep	7-Sep	--
	90 th Percentile	--	6-Oct	19-Sep	15-Sep	--	--	14-Oct	19-Sep	14-Sep	--
	N	0	41	182	41	0	0	2	10	5	0
2017	10 th Percentile	10-Sep	5-Sep	5-Sep	31-Aug	27-Sep	24-Sep	12-Sep	26-Aug	5-Sep	--
	50 th Percentile	20-Sep	18-Sep	14-Sep	12-Sep	27-Sep	24-Sep	12-Sep	12-Sep	15-Sep	--
	90 th Percentile	31-Oct	9-Oct	24-Sep	18-Sep	27-Sep	24-Sep	12-Sep	3-Nov	11-Oct	--
	N	8	19	63	48	1	1	1	19	13	0
2018	10 th Percentile	5-Sep	8-Sep	30-Aug	31-Aug	--	--	--	12-Sep	31-Aug	--
	50 th Percentile	20-Sep	20-Sep	3-Sep	31-Aug	--	--	--	12-Sep	15-Sep	--
	90 th Percentile	19-Oct	8-Oct	27-Sep	10-Sep	--	--	--	12-Sep	26-Oct	--
	N	10	37	13	2	0	0	0	1	5	0
2019	10 th Percentile	3-Sep	3-Sep	29-Aug	6-Nov	--	--	10-Sep	8-Sep	--	--
	50 th Percentile	13-Sep	20-Sep	8-Sep	6-Nov	--	--	10-Sep	12-Sep	--	--
	90 th Percentile	23-Sep	7-Oct	30-Sep	6-Nov	--	--	27-Sep	11-Oct	--	--
	N	21	82	62	1	0	0	2	12	0	0
2020	10 th Percentile	5-Sep	8-Sep	30-Aug	31-Aug	--	--	1-Sep	31-Aug	9-Sep	--
	50 th Percentile	20-Sep	20-Sep	3-Sep	31-Aug	--	--	6-Sep	31-Aug	20-Sep	--

Return Year	Hanford Reach Fall Chinook Migration Time (Date)										
	Origin	Priest Rapids Origin					Hanford Reach Natural Origin				
		Age 2	Age 3	Age 4	Age 5	Age 6	Age 2	Age 3	Age 4	Age 5	Age 6
	90 th Percentile	19-Oct	8-Oct	27-Sep	10-Sep	--	--	11-Oct	26-Sep	20-Sep	--
	N	20	83	95	3	0	0	4	6	3	0

13.2 Age at Maturity

Prior to return year 2012, the fish origin was assigned by location of survey due to the lack of identifiable hatchery marks and low CWT recoveries that may not have been representative of natural origin fish. Hence, the age composition for natural origin returns was generated from all the samples collected within the carcass survey regardless of true origin. Likewise, the age composition for hatchery origin fish was generated from all samples collected at PRH regardless of true origin. These 2007- 2015 brood year data suggests that between the two surveys, the proportions of age 3 fish returning to the hatchery were higher than observed in the escapement and the opposite for the other age classes of fish returning to the hatchery (Table 36)

The age compositions of the Hanford Reach escapement and the PRH returns are not directly comparable between locations without some adjustment or verification with another method. There is likely a recovery bias against smaller/younger fish in the stream surveys (Zhou 2002; Murdoch et al. 2010; Richards and Pearsons, 2013). Hence, the age composition for the Hanford Reach escapement is likely biased towards larger/older fish (Pearsons et al in preparation). All fish recovered from the PRH volunteer trap are available for systematic sampling, reducing the potential bias of the age composition data. Although this dataset is imperfect, the dataset is maintained for future reference should a method be established to correct the data for associated age and origin bias or if it is verified with another method.

The availability of otolith data combined with other hatchery mark data from the Hanford Reach carcass recoveries for the 2012 through 2020 return years provided the ability to estimate age compositions for both hatchery and natural origin fish within the demographic sample for the Hanford Reach escapement. However, the hatchery origin age composition may be influenced by

the low number of hatchery origin fish present in the demographic samples which is further reduced by sub-sampling the demographic origin. In addition, the age composition for both groups may be biased towards larger fish due to potential size recovery biases in the carcass surveys. Larger demographic samples per return year were required to better represent the age composition data before conclusions can be made. Beginning with return year 2014, the sub-sample size to determine origin was increased substantially to include up to 2,500 fish to obtain more hatchery origin fish in the sub-sample. Within the demographic sample of the escapement, the proportions of hatchery origin fish were higher than natural origin fish at age 3, and lower for ages 4, 5, and 6 during brood years 2007-2015 (Table 37, Table 38, and Table 39). Despite the differences in methods (trap vs. escapement, or hatchery vs. wild in the escapement), both methods identified a propensity for the hatchery to produce higher proportions of age 3 and lower proportions of age 4, 5, and 6 fish than the Hanford Reach.

Table 36 Age compositions for fall Chinook Salmon sampled in the Hanford Reach escapement compared to fall Chinook Salmon sampled at Priest Rapids Hatchery (genders combined), Brood Years 1998-2015.

Brood Year	Source ¹	Age Composition				
		Age-2	Age-3	Age-4	Age-5	Age-6
1998	Escapement	0.119	0.097	0.420	0.346	0.018
	PRH Returns	0.034	0.575	0.353	0.038	0.000
1999	Escapement	0.123	0.089	0.390	0.392	0.005
	PRH Returns	0.061	0.366	0.432	0.140	0.001
2000	Escapement	0.262	0.081	0.290	0.359	0.009
	PRH Returns	0.070	0.303	0.467	0.152	0.007
2001	Escapement	0.152	0.149	0.488	0.206	0.005
	PRH Returns	0.061	0.506	0.309	0.122	0.002
2002	Escapement	0.178	0.154	0.568	0.099	0.001
	PRH Returns	0.103	0.386	0.466	0.043	0.001
2003	Escapement	0.249	0.170	0.248	0.331	0.000
	PRH Returns	0.041	0.443	0.355	0.160	0.000
2004	Escapement	0.216	0.064	0.406	0.311	0.003
	PRH Returns	0.133	0.398	0.406	0.063	0.000
2005	Escapement	0.151	0.082	0.306	0.458	0.003
	PRH Returns	0.116	0.572	0.284	0.028	0.000
2006	Escapement	0.109	0.052	0.632	0.206	0.000
	PRH Returns	0.331	0.325	0.314	0.030	0.000
2007	Escapement	0.109	0.230	0.490	0.171	0.001
	PRH Returns	0.103	0.483	0.381	0.033	0.000
2008	Escapement	0.159	0.193	0.511	0.137	0.000
	PRH Returns	0.221	0.497	0.279	0.003	0.000
2009	Escapement	0.091	0.136	0.688	0.083	0.001
	PRH Returns	0.124	0.557	0.243	0.076	0.000
2010	Escapement	0.020	0.269	0.441	0.265	0.006
	PRH Returns	0.104	0.368	0.492	0.036	0.000
2011	Escapement	0.102	0.075	0.641	0.180	0.002
	PRH Returns	0.064	0.434	0.445	0.056	0.001
2012	Escapement	0.186	0.276	0.367	0.169	0.002
	PRH Returns	0.184	0.556	0.217	0.042	0.001
2013	Escapement	0.348	0.172	0.375	0.105	0.000
	PRH Returns	0.140	0.459	0.375	0.026	0.000
2014	Escapement	0.180	0.201	0.539	0.079	0.001
	PRH Returns	0.066	0.462	0.445	0.027	0.000
2015 ^a	Escapement	0.034	0.181	0.644	0.141	--
	PRH Returns	0.070	0.526	0.388	0.016	--
Mean 1998 – 2015	Escapement	0.155	0.148	0.469	0.224	0.003
	PRH Returns	0.113	0.456	0.370	0.061	0.001
Mean 2007 - 2015	Escapement	0.137	0.193	0.522	0.148	0.001
	PRH Returns	0.120	0.482	0.363	0.035	0.000

¹The origin is assigned by survey

^a Does not include age-6 returns

Table 37 Age compositions for male natural and hatchery origin fall Chinook Salmon sampled in the Hanford Reach escapement, Brood Years 2007-2015.

Brood Year	Origin ¹	N ²	Age Composition				
			Age-2	Age-3	Age-4	Age-5	Age-6
2007	Natural	1,093	No otolith data	0.377	0.483	0.139	0.002
	Hatchery	121		0.801	0.116	0.083	0.000
2008	Natural	1,234	0.044	0.336	0.502	0.118	0.000
	Hatchery	49	0.255	0.299	0.353	0.092	0.000
2009	Natural	816	0.034	0.231	0.66	0.076	0.000
	Hatchery	139	0.033	0.27	0.678	0.019	0.000
2010	Natural	2,097	0.008	0.361	0.454	0.176	0.000
	Hatchery	333	0.043	0.814	0.108	0.034	0.000
2011	Natural	838	0.182	0.157	0.547	0.112	0.002
	Hatchery	72	0.113	0.232	0.577	0.078	0.000
2012	Natural	858	0.058	0.527	0.319	0.095	0.001
	Hatchery	86	0.077	0.683	0.223	0.017	0.000
2013	Natural	517	0.029	0.463	0.442	0.066	0.000
	Hatchery	44	0.067	0.629	0.271	0.033	0.000
2014	Natural	297	0.057	0.420	0.461	0.062	0.000
	Hatchery	26	0.000	0.557	0.355	0.088	0.000
2015 ^a	Natural	822	0.023	0.325	0.572	0.080	--
	Hatchery	42	0.000	0.514	0.346	0.140	--
Mean	Natural	952	0.054	0.355	0.493	0.103	0.001
	Hatchery	101	0.074	0.533	0.336	0.065	0.000

¹Origin based on the presence of otoliths marks, hatchery coded-wire tags, and adipose clips present in the sub-sample.

²N equals the number fish included in the demographic sample for a specific brood year. Sample rates varied between return years; therefore, the age composition is based on pooled sample data expanded for total returns by year.

^a Does not include age-6 returns

Table 38 Age compositions for female natural and hatchery origin fall Chinook Salmon sampled in the Hanford Reach escapement, Brood Years 2007-2015.

Brood Year	Origin ¹	N ²	Age Composition				
			Age-2	Age-3	Age-4	Age-5	Age-6
2007	Natural	1,299	No otolith data	0.047	0.706	0.247	0.000
	Hatchery	167		0.532	0.317	0.151	0.000
2008	Natural	426	0.000	0.117	0.679	0.204	0.000
	Hatchery	74	0.000	0.176	0.651	0.172	0.000
2009	Natural	486	0.000	0.033	0.789	0.175	0.003
	Hatchery	188	0.000	0.06	0.918	0.021	0.000
2010	Natural	1,934	0.000	0.026	0.542	0.432	0.000
	Hatchery	353	0.000	0.418	0.448	0.133	0.000
2011	Natural	926	0.000	0.005	0.775	0.217	0.002
	Hatchery	118	0.000	0.022	0.782	0.195	0.000
2012	Natural	1,072	0.000	0.133	0.536	0.33	0.001
	Hatchery	165	0.000	0.382	0.479	0.138	0.000
2013	Natural	693	0.000	0.056	0.867	0.077	0.000
	Hatchery	91	0.000	0.219	0.586	0.195	0.000
2014	Natural	448	0.000	0.149	0.617	0.234	0.000
	Hatchery	33	0.000	0.106	0.849	0.045	0.000

Brood Year	Origin ¹	N ²	Age Composition				
			Age-2	Age-3	Age-4	Age-5	Age-6
2015 ^a	Natural	935	0.000	0.040	0.749	0.211	--
	Hatchery	111	0.000	0.248	0.664	0.088	--
Mean	Natural	913	0.000	0.067	0.696	0.236	0.001
	Hatchery	144	0.000	0.240	0.633	0.126	0.000

¹Origin based on the presence of otoliths marks, hatchery coded-wire tags, and adipose clips present in the sub-sample.

²N equals the number fish included in the demographic sample for a specific brood year. Sample rates varied between return years; therefore the age composition is based on pooled sample data expanded for total returns by year.

^a Does not include age-6 returns

Table 39 Age compositions for natural and hatchery origin fall Chinook Salmon sampled in the Hanford Reach escapement, Brood Years 2007-2015.

Brood Year	Sex Combined ¹	N ²	Age Composition				
			Age-2	Age-3	Age-4	Age-5	Age-6
2007	Natural	2,392	No Otolith	0.201	0.602	0.196	0.001
	Hatchery	288		0.656	0.225	0.119	0.000
2008	Natural	1,660	0.022	0.23	0.587	0.16	0.002
	Hatchery	123	0.1	0.224	0.535	0.141	0.000
2009	Natural	1,302	0.019	0.147	0.715	0.118	0.001
	Hatchery	327	0.012	0.136	0.831	0.021	0.000
2010	Natural	4,052	0.004	0.185	0.501	0.304	0.006
	Hatchery	686	0.022	0.617	0.278	0.084	0.000
2011	Natural	1,764	0.088	0.079	0.665	0.166	0.002
	Hatchery	190	0.038	0.093	0.713	0.156	0.000
2012	Natural	1,930	0.030	0.335	0.424	0.209	0.002
	Hatchery	251	0.030	0.5	0.378	0.091	0.000
2013	Natural	1,210	0.015	0.275	0.638	0.071	0.000
	Hatchery	135	0.024	0.372	0.462	0.143	0.000
2014 ^a	Natural	745	0.033	0.304	0.528	0.135	0.000
	Hatchery	59	0.000	0.258	0.682	0.060	0.000
2015 ^a	Natural	1,757	0.011	0.169	0.669	0.151	--
	Hatchery	153	0.000	0.344	0.550	0.106	--
Mean	Natural	1,868	0.028	0.214	0.592	0.168	0.002
	Hatchery	246	0.028	0.356	0.517	0.102	0.000

¹Origin based on the presence of otoliths marks, hatchery coded-wire tags, and adipose clips present in the sub-sample.

²N equals the number fish included in the demographic sample for a specific brood year. Sample rates varied between return years; therefore, the age composition is based on pooled sample data expanded for total returns by year.

^a Does not include age-6 returns

13.3 Size at Maturity

Prior to return year 2012, the size (fork length) at maturity comparisons between fall Chinook Salmon recovered at PRH and the Hanford Reach stream survey were calculated in a similar manner as the age composition data for the same period. Likewise, the assignment of origin was based on the survey (i.e., stream or hatchery). The estimates based on this method may not be representative of natural and hatchery origin fish due to possible size bias during recovery of carcasses.

Comparisons of the size at maturity data between the two surveys for brood years 2007 through 2015 suggests that ages 2 and 3 fish are similar in size and that ages 4, 5, and 6 fish are smaller in the hatchery survey (Table 40). The demographic sample for the 2012 through 2020 return

years provide the ability to estimate size at maturity for both hatchery and natural origin fish within the Hanford Reach escapement. These data suggest that either by gender or combined genders, hatchery origin fish are larger at age 2, similar at age 3, and smaller at ages 4, 5, and 6 (Table 41, Table 42 and Table 43). Again, the results between the two different methods provided similar findings.

Table 40 Mean fork length (cm) at age (total age) of fall Chinook Salmon sampled in the Hanford Reach escapement compared to fall Chinook Salmon sampled at Priest Rapids Hatchery, Brood Years 1999-2015. N = sample size and SD = 1 standard deviation.

Brood Year	Origin	Fall Chinook fork length (cm)														
		Age-2			Age-3			Age-4			Age-5			Age-6		
		N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
1999	Escapement	83	44	4	227	70	6	1,423	86	7	1,085	93	7	22	103	10
	PRH Returns	85	46	5	488	70	5	762	84	6	170	92	6	2	94	11
2000	Escapement	17	44	4	118	65	7	428	82	6	669	94	8	6	96	9
	PRH Returns	25	44	5	136	69	6	196	82	6	58	93	7	2	103	10
2001	Escapement	32	44	5	251	69	6	1,157	84	6	288	93	7	18	97	5
	PRH Returns	121	48	4	1,040	69	5	628	81	6	183	91	6	9	94	9
2002	Escapement	31	46	4	229	70	6	194	86	8	239	95	8	2	99	6
	PRH Returns	80	52	4	281	70	5	246	84	6	61	91	6	1	73	--
2003	Escapement	19	48	5	42	69	7	395	85	6	450	96	8	0	--	--
	PRH Returns	12	49	6	93	70	6	215	83	6	20	91	4	0	--	--
2004	Escapement	34	47	4	71	68	6	386	84	6	208	94	8	2	91	1
	PRH Returns	19	55	4	115	69	5	51	84	5	9	95	7	0	--	--
2005	Escapement	25	50	5	202	70	6	532	84	7	744	96	8	5	96	6
	PRH Returns	31	49	4	429	73	4	428	84	6	180	91	6	0	--	--
2006	Escapement	20	48	4	85	69	6	962	86	6	340	92	7	0	--	--
	PRH Returns	3	45	3	42	71	4	170	84	6	13	92	7	0	--	--
2007	Escapement	24	46	5	642	72	6	1,468	84	7	482	92	7	1	105	--
	PRH Returns	5	50	4	1,149	71	4	1,419	80	5	179	87	6	0	--	--
2008	Escapement	34	50	4	243	70	5	620	84	7	72	92	8	1	84	--
	PRH Returns	22	52	5	652	69	4	573	81	6	1	84	0	0	--	--
2009	Escapement	50	48	4	421	69	6	931	81	6	183	92	10	1	73	--
	PRH Returns	308	48	4	1,690	68	5	218	77	5	66	86	7	0	--	--
2010	Escapement	63	47	7	1,040	68	5	2,754	82	7	826	88	7	25	90	6
	PRH Returns	883	48	4	1,375	69	4	1,413	78	5	55	84	4	1	65	--
2011	Escapement	58	46	4	266	67	5	1,151	80	6	465	88	7	8	91	12
	PRH Returns	111	47	3	694	67	4	355	77	5	109	84	6	1	87	--
2012	Escapement	79	47	4	489	67	5	936	80	6	670	85	7	9	89	5
	PRH Returns	335	48	5	607	67	5	568	78	5	484	81	6	4	81	3
2013	Escapement	9	47	6	241	67	6	823	77	6	284	85	7	1	82	--
	PRH Returns	40	44	6	464	67	5	1,645	75	5	112	82	6	0	0	0
2014	Escapement	9	44	2	170	67	6	561	80	6	83	87	7	9	89	5
	PRH Returns	78	50	4	609	66	5	258	77	8	19	83	9	--	--	--
2015 ^a	Escapement	17	46	4	380	67	6	1,366	81	6	200	88	7	--	--	--
	PRH Returns	244	45	4	283	66	5	661	78	6	15	81	9	--	--	--
Mean	Escapement	37	47	4	296	69	6	920	83	6	443	91	8	7	92	6

Brood Year	Origin	Fall Chinook fork length (cm)														
		Age-2			Age-3			Age-4			Age-5			Age-6		
		N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
99 -15	PRH Returns	135	48	4	617	69	5	572	81	6	107	88	6	2	75	7
Mean	Escapement	45	47	4	459	68	5	1,097	81	6	337	88	7	6	88	5
07- 15	PRH Returns	211	48	4	801	68	5	805	78	6	171	84	6	3	64	3

^a Does not include age-6 returns.

Table 41 Mean fork length (cm) at age (total age) of male natural and hatchery origin fall Chinook Salmon that spawned naturally in the Hanford Reach, Brood Years 2007-2015. N = sample size and SD = 1 standard deviation.

Brood Year	Males by Origin	Fork Length (cm)														
		Age-2			Age-3			Age-4			Age-5			Age-6		
		N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
2007	Natural	No otolith Data			364	70	5	205	84	8	143	98	9	0	--	--
	Hatchery	No otolith Data			44	72	4	16	82	5	6	94	7	0	--	--
2008	Natural	22	49	4	134	69	5	260	85	8	25	99	7	0	--	--
	Hatchery	8	52	3	20	69	5	7	86	4	2	91	15	0	--	--
2009	Natural	3	48	3	325	68	6	123	82	6	40	99	7	0	--	--
	Hatchery	2	55	5	34	71	6	21	79	10	2	96	6	0	--	--
2010	Natural	31	45	4	291	68	7	855	83	8	135	94	8	4	97	8
	Hatchery	28	49	5	58	69	6	35	79	8	7	92	7	0	--	--
2011	Natural	31	45	4	176	66	5	403	81	8	137	94	7	3	104	3
	Hatchery	27	49	5	19	68	4	31	80	6	7	88	7	0	--	--
2012	Natural	45	47	4	312	67	6	316	80	8	140	92	8	1	88	--
	Hatchery	7	49	5	49	69	5	25	83	6	3	88	10	0	--	--
2013	Natural	8	47	6	179	67	6	269	79	8	48	91	9	0	--	--
	Hatchery	1	50	--	23	67	6	17	77	6	0	--	--	0	--	--
2014	Natural	10	49	4	116	67	6	151	82	9	20	95	7	0	--	--
	Hatchery	0	--	--	16	69	6	24	81	7	3	96	9	0	--	--
2015 ^a	Natural	16	46	5	289	67	3	468	84	8	46	96	8	0	--	--
	Hatchery	0	--	--	14	65	5	24	81	7	2	91	3	0	--	--
Mean	Natural	20	46	4	240	68	6	362	82	8	86	95	8	2	96	6
	Hatchery	12	50	4	39	69	5	22	81	7	4	92	9	0	--	--

^a Brood year does not include age-6 returns

Table 42 Mean fork length (cm) at age (total age) of female natural and hatchery origin fall Chinook Salmon that spawned naturally in the Hanford Reach, Brood Years 2007-2015. N = sample size and SD = 1 standard deviation.

Brood Year	Females by Origin	Fork Length (cm)														
		Age-2			Age-3			Age-4			Age-5			Age-6		
		N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
2007	Natural	No otolith Data			83	72	5	375	83	5	314	89	4	0	--	--
	Hatchery	No otolith Data			48	72	4	48	80	4	8	85	5	0	--	--
2008	Natural	0	--	--	36	70	3	344	83	5	49	88	5	1	84	--
	Hatchery	0	--	--	23	70	5	21	82	4	7	85	6	0	--	--
2009	Natural	0	--	--	44	71	5	105	80	4	82	87	11	1	73	--
	Hatchery	0	--	--	12	68	4	49	78	6	4	85	4	0	--	--
2010	Natural	0	--	--	33	71	5	999	87	5	528	85	4	20	89	5

Brood Year	Females by Origin	Fork Length (cm)														
		Age-2			Age-3			Age-4			Age-5			Age-6		
		N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
	Hatchery	0	--	--	22	69	4	144	79	5	29	82	4	0	--	--
2011	Natural	0	--	--	7	67	5	597	80	5	283	85	5	5	84	7
	Hatchery	0	--	--	4	65	2	72	77	4	34	84	4	0	--	--
2012	Natural	0	--	--	77	68	3	449	80	4	480	83	6	0	--	--
	Hatchery	0	--	--	42	68	3	83	78	6	38	81	5	0	--	--
2013	Natural	0	--	--	20	67	6	457	77	5	218	84	5	--	--	--
	Hatchery	0	--	--	12	67	4	58	75	5	12	80	7	1	82	--
2014	Natural	0	--	--	33	68	5	361	79	4	57	84	4	0	--	--
	Hatchery	0	--	--	6	69	4	22	78	5	2	86	1	0	--	--
2015 ^a	Natural	0	--	--	57	69	4	748	79	4	135	85	5	0	--	--
	Hatchery	0	--	--	9	69	3	84	78	5	8	83	6	0	--	--
Mean	Natural	0	--	--	43	69	5	493	81	5	238	86	5	5	83	6
	Hatchery	0	--	--	20	69	4	65	78	5	16	83	5	1	82	--

^a Brood year does not include age-6 returns

Table 43 Mean fork length (cm) at age (total age) of natural and hatchery origin fall Chinook Salmon that spawned naturally in the Hanford Reach, Brood Years 2007-2015. N = sample size and SD = 1 standard deviation.

Brood Year	Origin	Sex Combined Fork Length (cm)														
		Age-2			Age-3			Age-4			Age-5			Age-6		
		N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
2007	Natural	No otolith Data			447	70	5	580	83	6	457	92	6	0	--	--
	Hatchery	No otolith Data			92	72	4	64	81	4	28	87	6	0	--	--
2008	Natural	22	49	4	170	69	5	604	84	6	74	92	6	1	84	--
	Hatchery	8	52	3	43	70	5	28	83	4	9	86	8	0	--	--
2009	Natural	3	48	3	369	68	6	228	81	5	122	91	10	1	73	--
	Hatchery	2	55	5	46	70	5	70	78	7	6	89	5	0	--	--
2010	Natural	31	45	4	324	69	6	1,854	82	8	663	88	5	24	90	7
	Hatchery	27	50	6	80	69	6	179	79	7	36	84	5	0	--	--
2011	Natural	31	45	4	183	66	5	1,000	80	6	420	88	7	8	91	12
	Hatchery	28	50	6	23	67	4	103	78	5	41	84	5	0	--	--
2012	Natural	45	47	4	389	67	5	760	80	6	624	85	7	0	--	--
	Hatchery	7	49	5	91	68	4	108	79	6	41	81	6	0	--	--
2013	Natural	8	47	6	199	67	6	726	77	6	266	85	6	0	--	--
	Hatchery	1	50	--	35	67	5	75	76	5	12	80	7	0	--	--
2014	Natural	10	49	4	149	67	6	512	80	6	77	87	7	1	82	--
	Hatchery	0	--	--	22	69	5	46	80	7	5	92	8	1	89	--
2015 ^a	Natural	16	46	5	346	67	6	1,216	81	6	181	88	7	--	--	--
	Hatchery	0	--	--	23	67	5	108	79	5	10	85	7	--	--	--
Mean	Natural	21	47	4	286	68	6	831	81	6	320	88	7	6	84	10
	Hatchery	12	51	5	51	69	5	87	79	6	21	85	6	0	89	--

^a Brood year does not include age-6 returns

13.4 Gender Composition for Adult Escapement

Prior to return year 2012, the gender ratio comparisons between fall Chinook Salmon recovered at PRH and the Hanford Reach stream survey were based on the survey type (i.e., stream or hatchery). Although the estimates based on this method are imperfect, we continue to present this information to maintain the longest data set available (Table 44).

Gender ratios (male/females) by brood year and origin of adult fall Chinook Salmon sampled in the Hanford Reach carcass survey are presented in (Table 45). Annually, higher male to female ratios have been observed in the natural origin fish than that of the hatchery origin fish.

Table 44 Comparisons of male to female ratio of fall Chinook Salmon sampled at Priest Rapids Hatchery and in the Hanford Reach stream surveys, Brood Years 1996-2015.

Brood Year	Origin	Male ¹ : Female Ratio
1996	Stream	0.94:1
	Hatchery	1.98:1
1997	Stream	0.48:1
	Hatchery	1.88:1
1998	Stream	0.66:1
	Hatchery	1.38:1
1999	Stream	0.71:1
	Hatchery	2.15:1
2000	Stream	1.51:1
	Hatchery	2.40:1
2001	Stream	0.74:1
	Hatchery	2.31:1
2002	Stream	1.43:1
	Hatchery	1.94:1
2003	Stream	1.33:1
	Hatchery	1.64:1
2004	Stream	1.14:1
	Hatchery	1.63:1
2005	Stream	0.98:1
	Hatchery	2.15:1
2006	Stream	0.85:1
	Hatchery	2.57:1
2007	Stream	1.08:1
	Hatchery	1.60:1
2008	Stream	1.29:1
	Hatchery	1.89:1
2009	Stream	1.31:1
	Hatchery	2.57:1
2010	Stream	1.09:1
	Hatchery	1.47:1
2011	Stream	0.92:1
	Hatchery	2.00:1
2012	Stream	1.32:1
	Hatchery	1.82:1
2013	Stream	1.84:1
	Hatchery	1.91:1

Brood Year	Origin	Male ¹ : Female Ratio
2014	Stream	1.07:1
	Hatchery	2.27:1
2015 ^a	Stream	0.85:1
	Hatchery	2.35:1
Mean	Stream	1.08:1
	Hatchery	2.00:1

¹ Includes both adult males and jacks.

^a Includes age-2 through 5.

Table 45 Comparisons of male to female ratio of fall Chinook Salmon sampled in the Hanford Reach stream surveys, Brood Years 2007-2015.

Brood Year	Origin	Male ¹ : Female Ratio
2007 ^a	Natural	0.86:1.00
	Hatchery	0.74:1.00
2008	Natural	1.06:1.00
	Hatchery	0.64:1.00
2009	Natural	1.36:1.00
	Hatchery	0.56:1.00
2010	Natural	1.04:1.00
	Hatchery	1.01:1.00
2011	Natural	0.81:1.00
	Hatchery	0.50:1.00
2012	Natural	1.03:1.00
	Hatchery	0.64:1.00
2013	Natural	1.17:1.00
	Hatchery	0.57:1.00
2014	Natural	1.34:1.00
	Hatchery	0.51:1.00
2015 ^b	Natural	0.83:1.00
	Hatchery	0.56:1.00
Mean	Natural	1.06:1.00
	Hatchery	0.64:1.00

¹ Includes both adult males and jacks. ^a Does not include age-2. ^b Includes age-2 through 5.

13.5 Egg Retention

All female Chinook included in the demographic sample for the Hanford Reach stream surveys were examined for egg retention to assess spawn success. The females sampled were partitioned into the egg retention categories of 0%, 25%, 50%, 75% and 100%. The assignment of origin for each female for years 2010 and 2011 were based on the presence or absence of an adipose fin. The adipose intact group may include non-adipose clipped fish from PRH. A combination of hatchery marks (i.e., adipose clips, CWTs, and otolith marks) were used to identify hatchery origin fish in years 2012 - 2020. For all years, we assume that fish not possessing any hatchery marks are natural origin fish.

The assessment of egg retention was influenced by the loss of eggs during the collection and transport of carcasses prior to sampling. Therefore, our estimates of egg retention were likely to be underestimates and our estimates of egg loss were likely to be overestimates. In addition, the

methods for quantifying egg retention and assignment of origin for each female have varied among years. The amount of egg retention for years 2004 through 2013 were determined by visual estimates; whereas, during 2014 through 2020, the amount of retention was based on egg counts when the gametes were not completely intact. For these recent data sets, the percent of egg retention was calculated by dividing the amount of egg retained by an estimated fecundity based on length versus fecundity regressions by origin (Hatchery or Natural). An explanation of these regressions is provided in the fecundity section of this report.

The data from the egg counts were categorized into the standard egg retention categories based on the following ranges: 1 = 100-88%, 2 = 87-63%, 3 = 62-38%, 4 = 37-11%, and 5 = 10-0%. A comparison between visual and egg count methods was performed for years 2015 to 2020 to assess the egg retention estimates based on methods used prior to 2015. The difference between two methods was less than 1 percentage point by category for each year which provides some confidence that the visual methods of the past may provide reasonable indices of spawning success (Richards and Pearsons 2018).

We also calculated a mean spawn success for the female escapement for years 2004 through 2020 and a mean spawn success for the natural and hatchery origin female escapement for years 2012 through 2020. The calculations for the mean spawn success are weighted by the percentage of females sampled within each of the five egg retention categories.

The mean percentage of females recovered for years 2004 to 2020 that voided all of their eggs was 96.5% (Table 46). The mean percentage of females voiding all of their eggs for years 2012 to 2020 of natural and hatchery origin was 96.9% and 90.8%, respectively. The mean spawn success for year 2004 through 2020 was 98.2%. The mean spawn success for natural and hatchery origin females was 98.8% and 95.0%, respectively. The egg voidance means and resulting spawn success means were reduced by the 2013 outlier year.

Table 46 Comparison of egg retention of natural and hatchery origin fall Chinook sampled in the Hanford Reach stream survey, Return Years 2010-2020.

Return Year	Origin	Females Sampled	Egg Retention Categories					No Egg Retention (%)	Adj Spawn Success for Escapement (%)
			0 %	25%	50%	75%	100%		
2004	Combined	1,176	1,151	NA	21	NA	4	97.9	98.8
2005	Combined	1,323	1,310	NA	6	NA	7	99.0	99.2
2006	Combined	352	343	NA	8	NA	1	97.4	98.6
2007	Combined	454	443	NA	8	NA	3	97.6	98.5
2008	Combined	No spawn success data collected							
2009	Combined	499	484	NA	5	NA	10	97.0	97.5
2010	Combined	1,173	1,147	6	13	1	6	97.8	98.7
2011	Combined	1,264	1,203	1	52	5	3	95.2	97.4
2012 ^b	Natural	681	658	14	5	1	3	96.6	98.6
	Hatchery	90	89	0	0	0	1	98.9	98.9
	Total	771	747	14	5	1	4	96.9	98.6
2013 ^b	Natural	461	392	51	9	3	6	85.0	94.5
	Hatchery	224	144	39	11	13	17	64.3	81.3
	Total	685	536	90	20	16	23	78.2	90.1
2014 ^b	Natural	1,082	1,074	1	0	0	7	99.3	99.3

	Hatchery	153	141	3	0	0	9	92.2	93.6
	Total	1,235	1,215	4	0	0	16	98.4	98.6
2015 ^b	Natural	1256	1237	14	3	2	0	98.5	99.5
	Hatchery	149	135	7	5	2	0	90.6	96.1
	Total	1,405	1,372	21	8	4	0	97.7	99.1
2016 ^b	Natural	857	842	7	3	1	0	98.2	99.5
	Hatchery	138	127	11	3	1	0	92.0	96.4
	Total	995	969	18	6	2	0	97.4	99.1
2017 ^b	Natural	1,071	1,062	8	1	0	0	99.2	99.8
	Hatchery	109	100	5	2	2	0	91.7	96.6
	Total	1,180	1,162	13	3	2	0	98.5	99.5
2018 ^b	Natural	712	705	4	2	1	0	99.0	99.6
	Hatchery	46	43	1	0	1	1	93.5	95.7
	Total	758	748	4	2	1	0	98.7	99.6
2019 ^b	Natural	978	968	4	3	3	0	99.0	99.5
	Hatchery	128	124	3	1	0	0	96.9	99.0
	Total	1,106	1,092	7	4	3	0	98.7	99.5
2020 ^b	Natural	752	736	11	2	2	1	97.7	99.2
	Hatchery	267	260	2	0	0	5	97.4	97.9
	Total	1,109	996	13	2	2	6	97.7	98.9
Mean Natural Spawn Success (RY 2012 – 2020)								96.9	98.8
Mean Hatchery Spawn Success (RY 2012 – 2020)								90.8	95.0
Mean Combined Spawn Success (RY 2004 – 2020)								96.5	98.2

The measure for reporting egg retention changed from that used for previous years beginning in 2010

^b Origins were determined the presence or absence of otolith marks, adipose clips and CWTs

14.0 Contribution to Fisheries

The contribution of fish produced at PRH to fisheries was estimated by querying the Regional Mark Processing Center (RMPC) database. This is central repository for all CWT and otherwise associated release, catch, sample, and recovery data of anadromous Salmonids in the greater Pacific Coast Region of the United States of America. The Regional Mark Information System database (RMIS) within the RMPC provides specific recovery data for individual tag codes, along with the sample rate used to derive the estimated total number of recoveries by fishery type.

The CWT data reported to RMPC are expanded by sample rates generated by the agency reporting the data. In some cases, the estimated number of tags reported is less than the number reported as observed. This typically occurs when the sample rate is unknown, not reported, or biased (Gilbert Lensegrav, WDFW, personal communication). In these instances, the observed number was used instead of the estimated number to calculate the numbers of PRH origin fish recovered by location.

The RMIS database was queried on April 6, 2021 to provide CWT recoveries for active broods of PRH origin fish. The database for the 2014 brood should be complete for age-2 through age-5. The age-6 recovered during RY2020 may not be included until January 1, 2022 due to the lag in reporting field data to RMPC.

Beginning with the 2010 release year, portions of the non-adipose clipped smolts released from PRH received a CWT as part of a double index tag (DIT) study to evaluate the effect of various mark-selective fisheries occurring in Oregon, Washington, and British Columbia waters (PSC 2013). We are currently reviewing the data reported to the RMPC database to evaluate the results of the double index tagging for the PRH origin fish. Data for brood years 2014 through 2017 (some are incomplete due to time lag for reporting) show that adipose clipped fish from the DIT groups are being recovered in mark selective commercial and sport fisheries occurring in ocean, marine and freshwater zones. Comparisons of the demographics between the DIT groups recovered at PRH are very similar (Appendix G). Therefore, mark selective fisheries did not appear to markedly influence the demographic data collected at PRH.

Fall Chinook Salmon released from PRH supplement Pacific Ocean harvest for both commercial and sport fisheries from Washington to Southeast Alaska as well as Columbia River commercial, sport, and treaty tribal harvest. The Hanford Reach sport fishery for fall Chinook Salmon is an extremely popular fishery. This fishery typically runs annually from August 16 to late October. In 2020, an estimated 16,046 fall Chinook Salmon were harvested during this fishery (14,651 adults and 1,395 jacks). Estimates generated from CWT recoveries from the Hanford Reach sport fishery suggest that 30.4% (4,882 fish) of the total sport harvest in the Hanford Reach was comprised of fall Chinook Salmon released from PRH (Table 47). In comparison, fall Chinook Salmon released from Ringold Springs Hatchery comprised 8.4% (1,346 fish) of the sport fishery. Strays from other hatcheries combined represent 2.2% (346 fish) of the harvest. Sport harvest monitoring in the Hanford Reach and lower Yakima includes surveying both adipose intact and adipose clipped fish for CWT sampling.

The CWT data for PRH origin fall Chinook Salmon that were marked with an adipose clip were reviewed to assess contributions to marine and freshwater, commercial, tribal, and sport fisheries. The largest proportion of the harvest of PRH origin fall Chinook Salmon occurred in ocean fisheries followed by Zone-6 tribal harvest. For brood years 1997 through 2014, 48.8% of the reported harvest was in ocean fisheries and the other 55.9% in the Columbia River fisheries (Table 48). The adipose clip CWT rate for the broods after 2008 notably increased from previous brood years. Not all CWT survey locations check harvested adipose intact fish for the presence of a CWT. Therefore, the data presented in Table 48 includes harvest estimates based on recoveries of adipose clipped CWT tagged fish.

Table 47 Hatchery fall Chinook Salmon contributions to harvest in the Hanford Reach fall Chinook Salmon fishery. Coded-wire tag recoveries provided from RMIS database were expanded by sample rate and juvenile tag rate, Return Years 2003-2020.

Return Year	Harvest & CWT Sampling			CWT Expansions			% of Harvest		
	Harvest	Sampled	%	PRH	RSH	Other Hatcheries	PRH	RSH	Other Hatcheries
2003	7,190	1,848	25.7	510	424	43	7.1	5.9	0.6
2004	8,787	2,255	25.7	276	62	23	3.1	0.7	0.3
2005	7,974	1,834	23.0	1,200	265	35	15.0	3.3	0.4
2006	4,508	1,296	28.7	683	66	10	15.2	1.5	0.2
2007	6,466	1,812	28.0	929	50	89	14.4	0.8	1.4
2008	7,013	1,593	22.7	304	66	22	4.3	0.9	0.3
2009	8,806	1,741	19.8	520	0	10	5.9	0.0	0.1
2010	12,499	2,475	19.8	1,157	399	10	9.3	3.2	0.1

Return	Harvest & CWT Sampling			CWT Expansions			% of Harvest		
2011	14,262	2,715	19.0	1,558	663	121	10.9	4.6	0.8
2012	18,854	3,615	19.2	3,974	1,974	237	21.1	10.5	1.3
2013	27,630	5,555	20.1	6,570	3,947	537	23.8	14.3	1.9
2014	32,417	8,319	25.7	3,987	1,419	332	12.3	4.4	1.0
2015	35,419	10,327	29.2	4,144	992	319	11.7	2.8	0.9
2016	17,927	5,544	30.9	2,177	822	339	12.1	4.6	1.9
2017	12,368	4,435	35.9	1,585	843	105	12.8	6.8	0.8
2018	9,756	3,639	37.3	1,367	102	217	14	1.00	2.2
2019	13,149	4,569	34.7	1,205	120	88	9.2	0.9	0.7
2020	16,046	4,847	30.2	4,882	1,346	346	30.4	8.4	2.2
Mean	14,504	3,801	26.4	2,057	753	160	12.9	4.1	0.9

Table 48 Priest Rapids Hatchery coded-wire tag recoveries provided from RMIS by brood year and harvest type expanded by sample rate and juvenile tag rate, Brood Years 1997-2014. Data only includes coded-wire tag recoveries from adipose clipped fish expanded by the juvenile tag rate.

Brood Year	Ocean Fisheries		Columbia River Fisheries						Total Harvest	Ad-CWT Rate
	#	%	Tribal		Commercial		Recreational			
	#	%	#	%	#	%	#	%		
1997	1,100	36.7	1,506	50.2	304	10.1	91	3	3,001	3.0
1998	6,580	48.4	3,956	29.1	1,066	7.8	1,981	14.6	13,583	3.0
1999	14,190	54.6	5,908	22.8	2,410	9.3	3,458	13.3	25,966	2.9
2000	4,938	61.5	1,583	19.7	1,099	13.7	412	5.1	8,032	3.2
2001	17,758	56.5	6,612	21.1	1,554	4.9	5,484	17.5	31,408	5.2
2002	3,779	50.6	1,240	16.6	576	7.7	1,869	25	7,464	5.2
2003	1,871	54.6	570	16.6	226	6.6	757	22.1	3,424	5.9
2004	562	49.3	364	31.9	214	18.8	0	0	1,140	5.9
2005	10,699	52.1	5,975	29.1	998	4.9	2,871	14	20,543	3.0
2006	1,023	44.1	713	30.7	288	12.4	298	12.8	2,322	2.9
2007	13,838	44.4	10,620	34.1	2,160	6.9	4,523	14.5	31,141	3.0
2008	5,763	43.7	4,447	33.7	887	6.7	2,080	15.8	13,177	3.2
2009	24,872	43.4	21,121	36.8	2,581	4.5	8,761	15.3	57,335	9.1
2010	46,584	43.5	34,275	32	7,886	7.4	18,299	17.1	107,044	8.9
2011	18,235	44.2	11,813	28.6	3,874	9.4	7,310	17.7	41,232	8.4
2012	29,017	55.7	13,390	25.7	610	1.2	9,040	17.4	52,057	8.9
2013	9,027	37.2	6,824	28.1	1,023	4.2	7,378	30.4	24,252	8.3
2014	5,562	51.6	3,242	30.1	140	1.3	1,842	17.1	10,786	8.6
Mean	11,967	48.4	7,453	28.7	1,550	7.7	4,247	15.1	25,217	5.5

15.0 Straying

The distribution of PRH origin fish spawning in areas outside of the target stream is presented to assess the level of straying and potential impacts on other populations. Natural origin stray rates were presented in Pearsons and O'Connor 2020. The presumptive target spawning location for PRH origin fish includes the section of Columbia River from McNary Dam to Wanapum Dam as well as the lower Yakima River below Prosser Dam.

The spawning escapement of PRH origin fish by brood year is determined from CWT recoveries collected during spawning surveys. The CWT recoveries are expanded by the juvenile mark rates and survey sampling rates to estimate the number of PRH origin fish recovered on spawning grounds.

The stray rates (i.e., fish that spawned outside of the presumptive target area ÷ total escapement) for each brood year were calculated from the estimated recoveries of PRH origin fish from spawning grounds within and outside of the presumptive target area. CWT recoveries at non-target hatcheries and adult fish traps are not included. These fish were not considered strays because these fish were not able to leave the facilities on their own volition.

There are three stray rate metrics for recipient populations given in the Monitoring and Evaluation Plan for PUD Hatchery Programs based on return year stray rates and brood year stray rates (Hillman et al. 2017). The two stray rates based on return year for PRH origin fish are as follows:

- 1). Stray rate for PRH origin fall Chinook Salmon should be less than 5% of the spawning escapement for other non-target independent populations based on run year.
- 2). Stray rate for PRH origin fall Chinook Salmon should be less than 10% of the spawning escapement of any non-target streams within the independent population based on run year.
- 3). The donor stray rate for each hatchery brood year is monitored to determine if hatchery operations affect the homing and straying of specific brood years but does not include a specific target

The CWT recoveries by return year for presumptive non-target streams or areas suggest that PRH fall Chinook Salmon seldom exceeded more than 5% of the spawning escapement for other independent populations of fall Chinook Salmon. However, for multiple return years, greater than 5% of the spawning escapement for the Chelan River may have consisted of PRH origin fall Chinook Salmon (Table 49). The Chelan River spawning population is a mix of both summer and fall Chinook Salmon strays as well as fish returning from the Chelan hatchery program and is not considered an independent native population. This location was included to show contributions of PRH strays to this group of fish. With one exception (brood year 2006), less than 5% of the PRH origin returns for each brood year are estimated to have spawned outside of the presumptive target spawning area (Table 50).

Table 49 Percent of fall/summer Chinook spawning populations by return year (2000-2019) comprised of Priest Rapids Hatchery fall Chinook from 1998-2016 brood releases based on coded-wire tag recoveries.

Return Year	Presumptive Non-Target Stream													
	Snake Fall Chinook		Yakima Fall Chinook		Wenatchee Summer Chinook		Entiat River ¹		Chelan River ¹		Methow Summer Chinook		Okanogan Summer Chinook	
	#	%	#	%	#	%	#	%	#	%	#	%	#	%
2000	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
2001	0	0.0	0	0.0	0	0.0	0	0.0	334	33.9	0	0.0	0	0.0
2002	0	0.0	0	0.0	0	0.0	0	0.0	274	47.1	0	0.0	0	0.0

Return Year	Presumptive Non-Target Stream													
	Snake Fall Chinook		Yakima Fall Chinook		Wenatchee Summer Chinook		Entiat River ¹		Chelan River ¹		Methow Summer Chinook		Okanogan Summer Chinook	
	#	%	#	%	#	%	#	%	#	%	#	%	#	%
2003	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
2004	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
2005	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
2006	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
2007	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
2008	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	57	1.6
2009	0	0.0	0	0.0	0	0.0	0	0.0	228	36.5	0	0.0	0	0.0
2010	0	0.0	0	0.0	0	0.0	0	0.0	359	32.1	0	0.0	0	0.0
2011	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
2012	0	0.0	0	0.0	0	0.0	0	0.0	50	3.8	0	0.0	0	0.0
2013	0	0.0	0	0.0	0	0.0	0	0.0	102	6.1	0	0.0	0	0.0
2014	0	0.0	0	0.0	0	0.0	0	0.0	83	7.5	0	0.0	0	0.0
2015	0	0.0	0	0.0	0	0.0	22	1.9	323	22.5	0	0.0	0	0.0
2016	11	0.0	0	0.0	0	0.0	0	0.0	49	5.4	0	0.0	0	0.0
2017	0	0.0	0	0.0	0	0.0	0	0.0	89	10.4	0	0.0	0	0.0
2018	0	0.0	0	0.0	0	0.0	0	0.0	51	5.3	0	0.0	0	0.0
2019	0	0.0	0	0.0	0	0.0	0	0.0	48	4.1	0	0.0	0	0.0
Mean	1	0.0	0	0.0	0	0.0	1	0.1	100	10.7	0	0.0	3	0.1

¹The Chelan and Entiat River spawning populations are a mix of both summer and fall Chinook Salmon and are not considered independent native populations. These locations were included to show contributions of PRH strays to these groups of fish.

Table 50 Estimated number and percentage of Priest Rapids Hatchery fall Chinook Salmon spawning escapement to Priest Rapids Hatchery and stream within and outside of the presumptive target stream by brood year (1992-2014). Coded-wire tag recoveries are expanded by juvenile mark rate and survey sample rate for each brood year.

Brood Year	Number of PRH Origin Recoveries	Homing				Straying Outside of Target Stream	
		Target Hatchery		Target Stream ¹		Number	%
		Number	%	Number	%		
1992	9,037	7,630	84.4	1,037	11.5	370	4.1
1993	25,965	21,144	81.4	4,821	18.6	0	0.0
1994	1,693	1,385	81.8	308	18.2	0	0.0
1995	30,655	23,414	76.4	7,207	23.5	34	0.1
1996	13,551	10,034	74.0	3,517	26	0	0.0
1997	3,173	2,690	84.8	483	15.2	0	0.0
1998	18,167	11,833	65.1	5,867	32.3	467	2.6
1999	27,334	15,467	56.6	11,867	43.4	0	0.0
2000	4,759	3,690	77.5	1,069	22.5	0	0.0
2001	25,375	15,875	62.6	9,469	37.3	31	0.1

Brood Year	Number of PRH Origin Recoveries	Homing				Straying Outside of Target Stream	
		Target Hatchery		Target Stream ¹		Number	%
		Number	%	Number	%		
2002	5,288	3,769	71.3	1,519	28.7	0	0.0
2003	3,034	2,034	67.0	949	31.3	51	1.7
2004	1,133	1,133	100.0	0	0.0	0	0.0
2005	21,378	17,103	80.0	4,241	19.8	34	0.2
2006	1,001	641	64.0	0	0.0	367	36.7
2007	22,206	19,220	86.6	2,964	13.3	22	0.1
2008	11,866	9,002	75.9	2,864	24.1	0	0.0
2009	28,153	18,373	65.3	14,689	52.2	22	0.1
2010	107,961	67,940	62.9	40,574	37.6	327	0.3
2011	49,396	37,477	75.9	13,258	26.8	95	0.2
2012	71,635	59,189	82.6	12,341	17.2	105	0.1
2013	33,910	23,364	68.9	10,947	32.3	48	0.1
2014	8,652	5,714	66.0	2,938	34.0	0	0.0
Median	18,167	11,833	75.9	3,517	24.1	22	0.1

¹ Target stream includes the Columbia River between McNary and Wanapum dams as well as the Yakima River below Prosser Dam.

As previously described in Section 4, approximately 3,000 smolts at PRH were annually PIT tagged at PRH from brood years 1995 through 2010. The annual release of PIT tagged smolts was increased to ~43,000 beginning with brood year 2011. There were no PIT tagged smolts released from Priest Rapids Hatchery for brood year 2019 due to the COVID 19 pandemic. The last known observations of individual PIT tag adult fall Chinook Salmon originating from PRH at detection locations above McNary Dam are given in Table 51 for brood years 1999 through 2016. The number of observed PRH PIT tagged adults is increasing as anticipated due to the increased number of tags.

The majority of the PIT tagged PRH adults observed at McNary Dam have been observed at Priest Rapids Dam (PRD) adult fishways and/or PRH. Very few fish have been detected in the Snake River, which is an area of high concern for straying. In addition, notable proportions of the returns for several brood years have been observed at sites upstream of PRD. It is unclear whether fish spawned outside of the target areas because fish could return to a target location after being detected at a PIT tag array outside of the target stream without being detected again. Observations for PIT tagged presumptive Hanford Reach natural origin adults show very few detections above PRD.

Table 51 Last observations of unique PIT tagged adult fall Chinook from Priest Rapids Hatchery at detection sites upstream of McNary Dam, Brood Years 1999-2016.

Brood Year	# PIT tagged	Number of last known detections of unique Priest Rapids Origin PIT tags by site												
		MCN	ICH	PRO	PRH	PRD	RIA	LWE	RRF	EBO	ENL	WEA	LMR	Total
1999	3000	9				7	1					1		18
2000	3000	3				4								7
2001	3000	5				6								11
2002	3000	7				1								8
2003	3000													0

Brood Year	# PIT tagged	Number of last known detections of unique Priest Rapids Origin PIT tags by site												Total
		MCN	ICH	PRO	PRH	PRD	RIA	LWE	RRF	EBO	ENL	WEA	LMR	
2004	3000													0
2005	3000	9				4	1							14
2006	3000													0
2007	3,000	20			1	12	2		2			1	1	39
2008	2,994	5				6			1					12
2009	1,995	4			16		2							22
2010	3,000	8			34	23	5	1	3			3		77
2011	42,844	81			276	160	8	3	28	1		22	5	584
2012	42,908	101			435	122	6	1	20	1	1	14	2	703
2013	42,988	41			22	19	1		5			1	1	90
2014	42,621	9			33	4	1		3			1		51
2015 (age 2-5)	42,999	12			59	17			3			2		93
2016 (age 2-4)	42,858	59			100	32	2	2	8			3		206
MCN	McNary Dam Adult Fishways RKM 470						LWE	Lower Wenatchee River RKM 754						
ICH	Ice Harbor Dam Adult Fishways RKM 522						RRF	Rocky Reach Dam Adult Fishway RKM 763						
PRO	Prosser Diversion Dam RKM 539						EBO	East Bank Hatchery Outfall RKM 764						
PRH	Priest Rapids Hatchery Outfall RKM 635						ENL	Lower Entiat River RKM 778						
PRD	Priest Rapids Dam Adult Fishways RKM 639						WEA	Well Dam Adult Fishways RKM 830						
RIA	Rock Island Dam Adult Fishways RKM 730						LMR	Lower Methow River at Pateros RKM 843						

16.0 Genetics

Genetic tissue was collected from each Chinook Salmon spawned at PRH during 2020 by staff from the Columbia River Inter-Tribal Fish Commission (CRITFC). In total 5,626 specimens were collected at PRH to support their work associated with genetic stock identification and parentage-based tagging. The tissue samples collected from return years 2011 through 2020 are currently being archived by CRITFC. During 2010, WDFW staff collected 100 genetic tissue samples from both the Priest Rapids Hatchery broodstock and naturally spawning broodstock from the Hanford Reach.

17.0 Proportionate of Natural Influence

The intent of integrated hatchery programs is to achieve management objectives while having hatchery and natural origin fish share a common gene pool. Gene flow and the associated risks within and between the hatchery and natural environments can be estimated using a simple ratio estimator using the proportion of natural origin fish in the hatchery broodstock (pNOB) and the proportion of hatchery origin fish in the natural spawning escapement (pHOS). This ratio of pNOB/(pHOS+pNOB) is termed the Proportionate Natural Influence (PNI). The larger the PNI ratio, the greater selection that the natural environment has on the population relative to that of the hatchery environment. Alternatively, PNI estimates addressing gene flow from multiple sources/hatchery programs can be calculated from a multiple population gene flow model based on the Ford model which has been extended to three or more populations (Busack 2015, 2016; Pearsons et al. 2020).

In order for the natural environment to dominate selection, PNI for either calculation should be greater than 0.5 and for integrated hatchery programs the Hatchery Scientific Review Group

(HSRG) recommends a $PNI \geq 0.67$ (HSRG/WDFW/NWIFC 2004). The HSRG recommends a minimum target of 0.15 for the proportion of natural origin Chinook Salmon to be incorporated into the hatchery broodstock (pNOB) as well as a maximum target of 0.30 for the proportion of hatchery origin Chinook allowed to spawn in the natural environment (pHOS) for the Hanford Reach if it is to be managed as an integrated hatchery program.

Several estimates of PNI have been calculated to show the contributions of multiple programs on the overall PNI for the Hanford Reach. These programs include the hatchery production associated with the Grant PUD and USACE mitigation and the influence of strays. The different PNI estimates are based on pNOB and pHOS estimates specific to each source of spawning adults. The methods used to allocate pNOB and pHOS are described in the following sections.

17.1 Estimate of pNOB

Otolith marking began with the 2007 brood. Therefore, otolith marks are only available for specific age classes of PRH origin fish during return years 2010 and 2011 and do not provide representative samples for estimating pNOB for the PRH broodstock. Thus, estimates of pNOB based on otolith samples are limited to return years 2012 through 2020

The 2020 broodstock included 5,606 fish, some of which were used for off-site production for other programs. The Grant PUD and USACE programs utilized 5,231 fish from the broodstock that comprised of 4,314 fish from the volunteer trap and 1,007 from the ABC. In general, broodstock from ABC and OLAFT are held in a specific holding pond (Pond 4) and mated with fish from this pond or with fish collected from the PRH volunteer trap and held in holding ponds 3 and 2. The fish culturists segregate the progeny resulting from these matings for release from PRH. Broodstock utilized for non-Grant PUD programs are collected from the PRH volunteer trap and generally held in pond 2. Large portions of the progeny from the pond 2 broodstock are shipped to other facilities for use in other programs.

Grant PUD funds the collection of non-marked or tagged broodstock from the ABC and OLAFT with the intent of improving the pNOB associated with the production of their 5.6 million smolt mitigation requirement. The inclusion of these fish contributed greatly to the Grant PUD program's egg-take goal and the resulting pNOB (Table 52).

Table 52 Origin of broodstock and pNOB apportioned to program for fall Chinook Salmon spawned at Priest Rapids Hatchery, Brood Year 2020.

Program	Egg-Take	Facility Mean Fecundity	Natural Females	Hatchery Females	Natural Males	Hatchery Males	Total Natural	Total Hatchery	pNOB
GCPUD	6,681,816	3,775	408	1,362	449	448	857	1,810	0.321
GCPUD Alt Mating ¹	--	--	--	--	--	--	--	--	--
GCPUD Combined	6,681,816	3,775	408	1,362	449	448	857	1,810	0.321
USACE – PRH	2,017,054	4,150	52	434	18	219	70	653	0.097
USACE – RSH	4,585,067	3,574	9	1,274	9	639	18	1,913	0.009
USACE Combined	6,602,121	3,732	61	1,708	27	858	88	2,566	0.033
Combined PRH and RSH Programs	13,283,937	3,754	469	3,070	476	1,306	945	4,376	0.178
Other Programs ²	676,486	3,579	1	188	1	94	2	282	0.007

¹ Alternative mating strategy incorporates 1 natural origin male x 4 females did not occur during 2019.

² Includes eggs from presumed hatchery x hatchery crosses shipped to educational organizations.

The annual pNOB for fish spawned at PRH and used for Grant PUD and USACE smolt releases into the Hanford Reach during return years 2012 through 2020 is provided in Table 53.

Table 53 Number of broodstock spawned and pNOB apportioned to program for fall Chinook Salmon spawned at Priest Rapids Hatchery, Return Years 2012-2020.

Return Year	N	GCPUD pNOB	USACE pNOB	GCPUD and USACE Combined pNOB	Other Programs pNOB ¹
2012	4,974	0.182	0.057	0.119	N/A
2013	5,442	0.225	0.026	0.127	N/A
2014	5,443	0.343	0.076	0.206	0.000
2015	5,524	0.313	0.045	0.179	0.000
2016	4,938	0.259	0.073	0.163	0.000
2017	5,668	0.433	0.091	0.254	0.000
2018	5,836	0.650	0.156	0.387	0.141
2019	5,482	0.666	0.170	0.368	0.115
2020	5,606	0.321	0.033	0.178	0.007

¹ Represents pNOB associated with egg-takes utilized outside of the Hanford Reach.

17.2 Estimates of pHOS

Otolith marking began with the 2007 brood. Hence, otolith marks are only available for specific age classes of PRH origin fish during return years 2010 and 2011 and do not provide representative samples for estimating population level pHOS. In addition, estimates of pHOS for RSH based on otolith samples may be limited to return years 2012 through 2018 because ages 2-3 RSH origin returns were not otolith marked during return year 2020. However, all fish released from RSH were adipose fin clipped. The population level pHOS estimates for recent annual Hanford Reach spawning escapements are presented in Table 54.

Table 54 Proportion of hatchery Chinook Salmon on the spawning grounds (pHOS) in the Hanford Reach, Brood Years 2012-2020.

Return Year	N	Total Escapement	Hatchery Origin Escapement (pHOS)			
			PRH	RSH ¹	Other ¹	Total
2012	1,609	57,715	0.062	0.066	0.005	0.135
2013	927	174,841	0.203	0.054	0.018	0.275
2014	2,426	183,807	0.052	0.015	0.028	0.096
2015	2,485	266,328	0.076	0.017	0.004	0.097
2016	1,648	116,287	0.066	0.022	0.027	0.115
2017	1,813	73,759	0.063	0.017	0.001	0.081
2018	1,208	42,277	0.051	0.005	0.015	0.071
2019	1,686	65,991	0.077	0.004	0.024	0.105
2020	1,248	74,717	0.220	0.037	0.011	0.268
Mean	1,672	117,302	0.097	0.026	0.015	0.138

¹ Includes fish with coded-wire tags recovered within the demographic sample.

Estimates of pHOS were calculated for contributing sources of hatchery origin fall Chinook escapement in the Hanford Reach (Table 55). The pHOS associated with the PRH origin escapement was apportioned between the Grant PUD and USACE programs at PRH based on the annual program release goals from PRH of 5.6 million for Grant PUD and 1.7 million for

USACE. The pHOS estimate for return year 2020 includes 16,404 PRH origin fish in the escapement. Of these, 76.7% and 23.3% were allocated respectively to Grant PUD (12,608 fish) and USACE (3,830 fish) programs at PRH. The USACE’s 23.3% portion of the PRH origin escapement was combined with the escapement associated with the USACE’s RSH program (2,750 fish) to estimate the pHOS associated with the USACE programs in the Hanford Reach. There were 800 hatchery fish in the escapement associated with other hatchery programs located outside of the Hanford Reach.

Table 55 Origin of pHOS apportioned by program source for fall Chinook Salmon spawning naturally in the Hanford Reach, Return Years 2012-2020.

Return Year	Natural Origin	Hatchery Origin Spawners				pHOS by Source			
		GCPUD ¹	USACE ^{1,2}	Other ³	Total	GCPUD ¹	USACE ^{1,2}	Other ³	Combined
2012	50,072	3,943	3,598	261	7,803	0.068	0.062	0.005	0.135
2013	126,782	26,507	18,427	3,123	48,057	0.152	0.105	0.018	0.275
2014	166,183	7,185	5,262	5,120	17,567	0.039	0.029	0.028	0.096
2015	240,511	15,101	9,669	1,065	25,835	0.057	0.036	0.004	0.097
2016	103,033	5,732	4,513	3,143	13,388	0.049	0.039	0.027	0.115
2017	67,807	3,463	2,409	79	5,951	0.047	0.033	0.001	0.081
2018	42,277	1,760	841	696	3,297	0.038	0.018	0.015	0.071
2019	59,052	3,919	1,469	1,551	6,939	0.059	0.022	0.024	0.105
2020	54,759	12,608	6,580	800	19,964	0.168	0.088	0.011	0.268
Mean	101,164	8,913	5,863	1,760	16,536	0.075	0.048	0.015	0.138

¹Estimated number of PRH origin fish that spawned naturally in the Hanford Reach. Of these, 76.7% and 23.3% were apportioned to Grant PUD-PRH and USACE-PRH, respectively.

²Includes hatchery origin fish released from Ringold Springs Hatchery.

³Includes hatchery origin fish released from other hatcheries based on the presence of a hatchery mark without an otolith mark.

17.3 Estimates of PNI

We present PNI estimates based on pNOB and pHOS values calculated to reflect differing methodologies driven by the type of data available to assign origin of adult Chinook Salmon returns. The population level PNI for the Hanford Reach provided for each method includes all hatchery origin fish regardless of hatchery program or funding source.

Prior to return year 2012, pHOS, pNOB and PNI rates were based on CWT recoveries from the adult returns. Historically, we used juvenile mark rate expansions of CWT recoveries in the hatchery and stream surveys for these calculations. The pNOB estimated from CWT requires the assumption that fish unaccounted for by the juvenile mark rate expansions were natural origin fish. As discussed in Appendix A of this report, this assumption may overestimate pNOB and PNI. This method of estimated pNOB of the broodstock for the years following 2014 was not calculated due to culling fish possessing a CWT and or an adipose clip. Hence, the broodstock origin was poorly represented by CWT.

The pHOS estimates based on juvenile mark rate expansions of CWT recoveries also may underestimate the presences of PRH and RSH origin fish as explained in Appendix A. For comparison, we present CWT based estimates of PNI derived from CWT adult-to-adult expansions for PRH and RSH origin adult recoveries at their respective hatcheries. An explanation of methods is given in Appendix H. Estimates of pNOB, pHOS, and PNI based on both methods of CWT expansions are presented in Table 56.

The pHOS and pNOB estimates from limited otolith datasets for recent complete brood years are more like the estimates produced by adult-to-adult CWT expansions versus juvenile mark rate expansions of CWT recoveries of returning adults.

Table 56 PNI of the Hanford Reach fall Chinook Salmon supplementation program based on expanded coded-wire tag recoveries of all fish surveyed, Return Years 2001-2020.

Return Year	pNOB ¹	pHOS ¹	pNOB ²	pHOS ²	PNI based on pNOB ¹ and pHOS ¹	PNI based on pNOB ² and pHOS ²
2001	0.155	0.094	0.046	0.066	0.622	0.411
2002	0.145	0.101	0.046	0.125	0.589	0.269
2003	0.132	0.099	0.046	0.117	0.571	0.282
2004	0.229	0.081	0.046	0.099	0.739	0.317
2005	0.370	0.106	0.046	0.156	0.777	0.229
2006	0.507	0.057	0.046	0.124	0.899	0.271
2007	0.326	0.041	0.046	0.065	0.888	0.414
2008	0.501	0.046	0.046	0.087	0.916	0.346
2009	0.568	0.077	0.046	0.174	0.881	0.209
2010	0.392	0.040	0.046	0.076	0.907	0.377
2011	0.381	0.076	0.046	0.154	0.836	0.230
2012	0.304	0.074	0.119 ^a	0.118	0.871	0.529
2013	0.252	0.217	0.127 ^a	0.287	0.537	0.300
2014	0.443	0.054	0.206 ^a	0.069	0.888	0.760
2015	N/A ³	0.072	0.179 ^a	0.075	N/A ³	0.691
2016	N/A ³	0.092	0.163 ^a	0.097	N/A ³	0.627
2017	N/A ³	0.116	0.254 ^a	0.102	N/A ³	0.713
2018	N/A ³	0.071	0.372 ^a	0.091	N/A ³	0.803
2019	N/A ³	0.050	0.368 ^a	0.050	N/A ³	0.880
2020	N/A ³	0.216	0.178 ^a	0.237	N/A ³	0.429

pNOB¹ assumes that all fish not accounted for by juvenile coded-wire tag expansions are natural origin.

pHOS¹ based on hatchery origin coded-wire recoveries expanded by juvenile mark rate and survey sample rate.

pNOB² is assigned to years 2001 – 2011 based on an average proportion of natural origin returns to PRH for return years 2012 -2014 as determined by otolith and other hatchery marks.

pHOS² is based on an adult coded-wire tag expansion rate for PRH and RSH origin adults recovered in the Hanford Reach escapement combined with juveniles coded-wire tag mark rate expansions for other hatchery strays. Both groups were expanded by the survey sample rate.

³ Brood stock was generally high-graded to remove coded-wire tagged fish during ponding.

^apNOB of broodstock used for production of PRH and RSH programs as determined from otoliths and other hatchery marks.

For return years 2012-2020 we present PNI estimates calculated from the multiple population gene flow model (Busack 2015) and otoliths (Table 57). This is likely our most accurate estimate of PNI. The output from this model indicates that the PNI values for return years 2014 and 2019 have exceeded the goal of 0.670. The PNI for return year 2020 was 0.546. High pHOS during return year 2020 contributed to a lower PNI than observed in recent years.

Table 57 PNI estimates for the Hanford Reach fall Chinook Salmon supplementation programs based on otoliths, Return Years 2012-2020. Calculated from multiple population gene flow model based on the Ford model which has been extended to three or more populations.

Return Year	pNOB			pHOS			pHOS Reach ⁷	PNI Population ⁸
	GCPUD ¹	USACE ²	Facility ³	GCPUD ⁴	USACE ⁵	Other ⁶		
2012	0.182	0.057	0.119	0.068	0.062	0.005	0.135	0.599
2013	0.225	0.027	0.127	0.152	0.105	0.018	0.275	0.463
2014	0.343	0.076	0.206	0.039	0.029	0.028	0.096	0.775
2015	0.313	0.045	0.179	0.057	0.036	0.004	0.097	0.762
2016	0.259	0.072	0.163	0.049	0.039	0.027	0.115	0.700
2017	0.433	0.091	0.254	0.047	0.033	0.001	0.081	0.835
2018	0.650	0.156	0.387	0.038	0.018	0.015	0.071	0.891
2019	0.666	0.170	0.368	0.059	0.022	0.024	0.105	0.852
2020	0.321	0.033	0.178	0.169	0.088	0.011	0.268	0.546
Mean	0.377	0.081	0.220	0.075	0.048	0.015	0.138	0.714

¹Includes broodstock associated with Grant PUD production at PRH.

²Includes broodstock associated with USACE production at PRH and RSH.

³Includes broodstock spawned at PRH for all production

⁴Includes pHOS associated with Grant PUD mitigation smolt releases at PRH

⁵Includes pHOS associated with USACE mitigation smolt releases at PRH and RSH

⁶Includes pHOS associated with strays from hatcheries outside of the Hanford Reach

⁷Population level pHOS in the Hanford Reach

⁸Population level PNI for the Hanford Reach. Assumes strays from hatcheries outside of the Hanford Reach have an associated pNOB of zero.

18.0 Natural and Hatchery Replacement Rates

The numbers of hatchery origin recruits (HOR) are estimated from CWT recoveries for brood year returns to the PRH and the Hanford Reach of the Columbia River. The recovered CWTs are expanded by sample rate of the survey and then by the juvenile tag rate. CWTs recovered from natural origin recruits (NOR) originating from the Hanford Reach are difficult to expand accurately because the juvenile tag rates are unknown. While imperfect, we make the assumption that returns not accounted for by HOR CWT recoveries are NOR.

Hatchery replacement rates (HRR) were calculated as the ratio of HOR to the parent broodstock at PRH. This broodstock is an estimate of the number of fish spawned at PRH to produce the target release of subyearling fall Chinook Salmon. Similarly, natural replacement rates (NRR) for the Hanford Reach URB fall Chinook Salmon were calculated as the ratio of NOR to the parent population spawning naturally in the Hanford Reach natural environment. This spawning population was based on the escapement estimate to the Hanford Reach without adjustments for spawn success.

Harvest estimates for HOR were calculated from the proportion of the expanded CWT recoveries in the fisheries to the total number of the expanded CWTs recoveries included in fisheries, stream surveys, and hatchery racks. The CWT recoveries are expanded by sample rate of the survey and juvenile mark rate for the CWT group. Since there is not a CWT mark rate for NOR, the harvest rates for PRH origin returns (HOR) were used as an indicator for similar brood years of NORs.

The HRR and NRR for brood year 2014 that includes harvest was 6.79 and 0.38, respectively (Table 58). In comparison, the HRR and NRR for this brood year without harvest included was 3.41 and 0.19, respectively. The HRR should be greater than or equal to 5.30 (the target value in Murdoch and Peven 2005).

Table 58 Broodstock spawned at Priest Rapids Hatchery, estimated escapement to the Hanford Reach, natural and hatchery origin recruits (NOR and HOR), and natural and hatchery replacement rates (NRR and HRR, with and without harvest) for natural origin fall Chinook Salmon in the Hanford Reach, Brood Years 1996-2013.

Brood Year	Broodstock Spawned	Hanford Reach Escapement ¹	Harvest not included				Harvest included ²			
			HOR	NOR	HRR	NRR	HOR	NOR	HRR	NRR
1996	2,859	43,249	13,584	28,849	4.75	0.67	26,205	59,899	9.17	1.38
1997	2,726	43,493	3,002	44,416	1.10	1.02	6,037	88,349	2.21	2.03
1998	3,027	35,393	18,464	93,999	6.10	2.66	31,932	222,865	10.55	6.30
1999	2,619	29,812	27,093	115,237	10.34	3.87	52,099	240,090	19.89	8.05
2000	2,619	48,020	4,665	56,422	1.78	1.17	12,508	89,983	4.78	1.87
2001	3,621	59,848	25,059	71,359	6.92	1.19	55,789	129,548	15.41	2.16
2002	3,630	84,509	5,277	47,813	1.45	0.57	12,744	81,600	3.51	0.97
2003	3,003	100,508	3,021	31,788	1.01	0.32	5,974	64,307	1.99	0.64
2004	3,014	87,696	1,109	22,747	0.37	0.26	3,262	34,465	1.08	0.39
2005	2,898	71,967	21,107	64,011	7.28	0.89	61,122	97,777	21.09	1.36
2006	2,911	51,701	998	54,288	0.34	1.05	3,347	77,344	1.15	1.50
2007	2,096	22,274	22,184	101,753	10.58	4.57	52,832	175,404	25.21	7.87
2008	2,959	29,058	11,867	41,809	4.01	1.44	25,166	79,116	8.51	2.72
2009	3,177	36,720	28,154	97,834	8.86	2.66	85,489	145,874	26.91	3.97
2010	3,320	87,016	97,567	281,364	29.38	3.23	209,338	526,972	63.05	6.06
2011	3,121	75,256	49,396	168,864	15.83	2.24	90,628	371,161	29.04	4.93
2012	3,008	57,715	70,175	160,185	23.33	2.78	124,058	368,804	41.24	6.39
2013	4,370	174,651	33,904	82,464	7.76	0.47	58,181	197,627	13.31	1.13
2014	3,189	183,749	10,867	34,826	3.41	0.19	21,653	69,916	6.79	0.38
Mean	3,061	69,612	23,552	84,212	7.73	1.64	49,388	164,268	16.41	3.16
Median	3,008	57,715	18,464	64,011	6.10	1.17	31,932	97,777	10.55	2.03

¹ Includes estimated adult and jack escapement to the Hanford Reach natural environment.

² Harvest rates for NORs was estimated using the HRRs harvest rates for similar brood years as an indicator stock.

19.0 Smolt-to-Adult Survivals

Smolt-to-adult survival ratios (SAR) were calculated by dividing the expanded number of adults possessing a CWT recovered by the number of CWT smolts released. This estimate could be biased low for both hatchery and natural origin fish because of some of CWT bias identified previously in this report. The following data was obtained from the RMPC’s RMIS online database: <http://www.rmpec.org/>. The SAR by brood year data was re-queried on April 13, 2021. This query should account for age 2 through 6 fall Chinook Salmon sampled through December 2019. The lag in reporting field data for the 2020 return year likely excludes recoveries of a limited number of age-6 fish from the 2014 brood.

Annual brood year SAR for hatchery fall Chinook Salmon released from PRH from 1992 through 2014 had a mean of 0.0069 with a median of 0.0049 (Table 59). The SAR for the PRH origin 2010 brood was 0.0304, which was the highest SAR on record for PRH releases.

Table 59 Smolt-to-adult Survival Ratios (SAR) for Priest Rapids Hatchery fall Chinook Salmon, Brood Years 1992-2014. Data includes all coded-wire tag recoveries from adipose clipped fish.

Brood Year	Number of Tagged Smolts Released	Estimated Adult Captures	SAR
1992	194,622	456	0.0023
1993	185,683	1,474	0.0079
1994	174,033	108	0.0006
1995	196,089	1,789	0.0091
1996	193,195	762	0.0039
1997	196,203	184	0.0009
1998	193,545	943	0.0049
1999	204,506	1,578	0.0077
2000	200,784	368	0.0018
2001	219,918	1,829	0.0083
2002	355,373	672	0.0019
2003	399,119	351	0.0009
2004	200,072	95	0.0005
2005	199,445	1,751	0.0088
2006	202,000	96	0.0005
2007	202,568	2,338	0.0115
2008	218,011	727	0.0033
2009	619,568	7,903	0.0128
2010	603,790	18,724	0.0310
2011	595,608	4,184	0.0129
2012	603,930	6,485	0.0175
2013	603,797	2,539	0.0080
2014	604,850	1,196	0.0020
Mean	320,292	2,437	0.0069
Median	202,568	943	0.0049

Annual SAR for Hanford Reach natural origin fall Chinook Salmon for brood years 1992 through 2014 had a mean of 0.0043 with a median of 0.0020 (Table 60). The SAR for the Hanford Reach natural origin 2010 brood was 0.0178 which was the highest SAR on record for the Hanford Reach natural origin stock.

Table 60 Smolt-to-adult Survival Ratios (SAR) for Hanford Reach natural origin fall Chinook Salmon, Brood Years 1992-2014. Data includes all coded-wire tag recoveries from adipose clipped fish. Source Regional Mark Processing Center.

Brood Year	Number of Tagged Smolts Released	Estimated Adult Captures	SAR
1992	203,591	820	0.0040
1993	95,897	486	0.0051
1994	148,585	74	0.0005
1995	146,887	340	0.0022
1996	92,262	110	0.0012
1997	199,896	358	0.0018
1998	129,850	783	0.0060
1999	213,259	2,367	0.0111

Brood Year	Number of Tagged Smolts Released	Estimated Adult Captures	SAR
2000	204,925	362	0.0018
2001	127,758	519	0.0041
2002	203,557	340	0.0017
2003	207,168	201	0.0010
2004	163,884	143	0.0009
2005	203,929	381	0.0019
2006	263,478	357	0.0017
2007	53,618	446	0.0083
2008	203,947	556	0.0027
2009	201,606	1,616	0.0080
2010	179,727	2,919	0.0178
2011	166,610	1,063	0.0064
2012	148,107	1,770	0.0120
2013	179,952	487	0.0027
2014	159,209	91	0.0006
Mean	216,731	718	0.0043
Median	202,939	414	0.0020

20.0 ESA/HCP Compliance

20.1 Broodstock Collection

Section 10(a)(1)(B) Permit 23194 (NMFS 2019) authorizes collection of hatchery origin Chinook Salmon at the Priest Rapids Volunteer Trap for broodstock and surplus operations as well as up to 1,323 natural origin Chinook broodstock from the OLAFT and by angler hook and line for the Priest Rapids hatchery program. A total of 1,166 natural origin broodstock were collected by hook and line during 2020. Broodstock was not collected at the OLAFT during 2020. The permit also allows for the handling of up to 15 steelhead and a take of 3 steelhead for operation of the PRH volunteer trap. A total of 3 steelhead, one of which was not adipose clipped, were encountered at the PRH volunteer trap with no incidental mortality reported. No steelhead mortalities were associated with broodstock collection in the ABC fishery (Table 61).

Table 61 Recoveries and disposition of steelhead at the Priest Rapids Hatchery volunteer trap, Return Year 2020.

		No Mark	Ad Only	Ad-RV	Total
Released	Males	1	1	0	2
	Females	0	1	0	1
	Sub Total	1	2	0	3
Killed	Males	0	0	0	0
	Females	0	0	0	0
	Sub Total	0	0	0	0
Total		1	2	0	3

20.2 Hatchery Rearing and Release

The juvenile fall Chinook Salmon from the 2020 brood year reared throughout their life-stages at PRH without incident. The reported smolt release for the 2020 brood totaled 7,542,518 URB fall Chinook Salmon, representing 103% of the production objective and was compliant with the 10% overage allowable in the ESA Permit. There were no precocious smolts observed during

pre-release sampling (Section 9.4, Table 15). The median travel time between the initial detections at the PRH PIT-tag array and the McNary Dam PIT-tag arrays by release group ranged from 22 days for the first release group to 8 days for the last release group. The median travel time for the entire release was 12 days and did not exceed the 13 day of residence target stated in the permit.

20.3 Hatchery Effluent Monitoring

Per ESA Permit WAG-13-7013 states that the permit holder shall monitor and report hatchery effluents in compliance with applicable National Pollution Discharge Elimination Systems (NPDES) (EPA 1999) permit limitations. There were no NPDES violations reported at Grant PUD Hatchery facilities during the September 2020 through June 2021 collection and rearing periods.

20.4 Ecological Risk Assessment

One of the regional objectives in the Grant PUD M&E plan is to conduct an ecological risk assessment on non-target taxa of concern to determine if additional M&E is necessary (Pearsons and Langshaw 2009). The methodology that was used to assess risks was presented in Pearsons et al. (2012) and Pearsons and Busack (2012). This objective was completed through an approved report that summarized the methods and results of the risk assessment (Mackey et al. 2014).

21.0 Distribution of Adult fall Chinook Salmon carcass from Priest Rapids Hatchery

All adult Chinook Salmon recovered at PRH were eventually distributed to multiple organizations depending on the condition and treatment of the individual fish while at the hatchery. Many of these fish were suitable for consumption and transported to foodbanks (Table 62). The numbers in Table 62 do not include fish that were surplus and used for other hatchery programs.

Table 62 Disposition of Chinook Salmon removed from Priest Rapids Hatchery volunteer trap, Return Year 2001-2020. Surplus numbers do not include shipped to other hatchery programs.

Return Year	Disposal of Mortalities		WDFW Nutrient Enhancement Projects	Donations to Educational Programs & Research	Donations to Foodbanks	Donations to Tribes	Sold to Fish Buyers	Fish Removed from Priest Rapids Hatchery
	Pet Food	Landfill						
2001	0	6,597	2,054	0		525	6,139	15,315
2002	0	6,572	2,192	0	3,130	502	0	12,396
2003	0	5,144	3,211	9	881	98	0	9,343
2004	350	2,661	2,756	88	9,371		595	15,821
2005	153	5,635	318	2	0		4,503	10,611
2006	0	5,467	0	250	0	340	2,146	8,203
2007	2,595	0	0	0	0	159	3,345	6,099
2008	5,384	90	0	340	0	375	13,428	19,617
2009	5,846	0	0	310	0	201	6,502	12,859
2010	5,412	1,937	1,937	452	3,548	8	8,259	21,553
2011	6,951	0	1,500	412	11,217	588	0	20,668
2012	7,554	0	0	460	20,628		0	28,642
2013	10,108	0	0	489	31,647	626	0	42,870
2014	10,805	0	0	237	67,684	783	0	79,509

2015	7,402	0	0	398	52,987	4,228	0	65,015
2016	7,833	0	0	411	19,424	1,948	0	29,616
2017	10,905	0	0	436	6,413	1,505	0	19,259
2018	8,670	0	0	480	8,647	350	0	18,147
2019	9,743	0	0	409	10,878	113	0	21,601
2020	9,422	0	0	160	28,072	2,345	0	39,999
Mean	5,457	1,705	698	267	14,449	864	2,246	24,857

22.0 Acknowledgments

We thank the many individuals and organizations that helped collect the data or provided data for inclusion in this report: Shawnaly Meehan and Dennis Werlau for the leadership of their WDFW M&E crews that sorted and sampled 51,793 fall Chinook Salmon, along with entering and reviewing all the sample data collected in 2020; Paul Hoffarth for supervising the creel and carcass staff and summarizing the sport harvest data; and Alf Haukenes for providing helpful suggestions for the improvement of this report. Deb Firestone (Grant PUD) assisted with final formatting of the report. Furthermore, we would like to thank the hatchery staff at Priest Rapids and Ringold Springs Hatcheries: Brian Lyon, Glen Pearson, Mike Erickson, and their crews for accommodating the activities associated with M&E. This work was primarily funded by Grant PUD, the USACE, WDFW, and the CRCWTRP.

23.0 Literature Cited

- AFS-FHS (American Fisheries Society-Fish Health). 2014. FHS blue book: suggested procedures for the detection and identification of certain finfish and shellfish pathogens, 2014 edition.
- Busack, C. 2015. Extending the Ford model to three or more populations. August 31, 2015. Sustainable Fisheries Division, West Coast Region, National Marine Fisheries Service. 5p.
- Busack , C. 2016. Methow Gene Flow 2.xlsx. NMFS West Coast Region Portland, Oregon.
- Elliot, D. G. 2012. Bacterial kidney disease. *In* AFS-FHS (American Fisheries Society- Fish Health Section. FHS blue book: suggested procedures for the detection and identification of certain finfish and shellfish pathogens, 2014 edition.
- Fryer, J. K. 2018. Expansion of the 2017 Hanford Reach fall Chinook Salmon juvenile coded-wire tagging and PIT tagging project. Report submitted to the Pacific Salmon Commission Technical Committee from the Columbia River Inter-Tribal Fish Commission, Portland, Oregon.
- Ham, K. D., and T. N. Pearsons. 2001. A practical approach for containing ecological risks associated with fish stocking programs. *Fisheries* 25(4):15-23.
- Harnish, R. A., R. Sharma, G. A. McMichael, R. B. Langshaw, and T. N. Pearsons. 2014. Effect of hydroelectric dam operations on the freshwater productivity of a Columbia River fall Chinook Salmon population. *Canadian Journal of Fisheries and Aquatic Sciences* 71:602-615.
- Harnish, R.A., R. Sharma, G. A. McMichael, R.B. Langshaw, T.N. Pearsons, and D.A. Bernard. 2012. Effects of Priest Rapids Dam Operations on Hanford Reach Fall Chinook Salmon Productivity and Estimation of Maximum Sustainable Yield, 1975-2004. Prepared for: Public Utility District No. 2 of Grant County, Ephrata, WA. Contract Number 430-2464.
- Harnish, R. A. 2017. Hanford Reach Upriver Bright Productivity Analysis Update. Pacific Northwest National Laboratory, Richland, Washington.
- Hillman, T., T. Kahler, G. Mackey, A. Murdoch, K. Murdoch, T. Pearsons, M. Tonseth and C. Willard. 2017. Monitoring and evaluation plan for PUD hatchery programs: 2017 update. Report to the HCP and PRCC Hatchery Committees, Wenatchee and Ephrata WA.
- Hoffarth, P. A. and T. N. Pearsons. 2012a. Priest Rapids Hatchery Monitoring and Evaluation: Annual Report for 2010. Grant County Public Utility District, Ephrata, Washington.
- Hoffarth, P. A. and T. N. Pearsons. 2012b. Priest Rapids Hatchery Monitoring and Evaluation: Annual Report for 2011. Grant County Public Utility District, Ephrata, Washington.
- Langshaw, R. B., P. J. Graf and T. N. Pearsons. 2015. Effects of the Hanford Reach Fall Chinook Protection Program on Fall Chinook Salmon in the Hanford Reach –Summary, Conclusions, and Future Monitoring. Grant County Public Utility District, Ephrata, Washington.

- Langshaw, R. B., P. J. Graf and T. N. Pearsons. 2017. Hydropower and high productivity in the Hanford Reach: A synthesis of how flow management may benefit fall Chinook Salmon in the Columbia River, USA. *WIREs Water*. 2017;e1275. <https://doi.org/10.1002/wat2.1275>
- Mackey, G., T. N. Pearsons, M. R. Cooper, K. G. Murdoch, A. R. Murdoch, and T. W. Hillman. 2014. Ecological risk assessment of upper Columbia hatchery programs on non-target taxa of concern. Report produced by the Hatchery Evaluation Technical Team (HETT) for the HCP Wells Hatchery Committee, HCP Rocky Reach Hatchery Committee, HCP Rock Island Hatchery Committee, and the Priest Rapids Hatchery Sub-Committee.
- Murdoch, A.R., and C. Peven. 2005. Conceptual approach to monitoring and evaluating the Chelan County Public Utility District Hatchery Programs. Final report to the Chelan PUD Habitat Conservation Plan's Hatchery Committee.
- Murdoch, A. R., T. N. Pearsons, and T. W. Maitland. 2010. Estimating the spawning escapement of hatchery- and natural-origin spring Chinook Salmon using redd and carcass data. *North American Journal of Fisheries Management* 30:361-375.
- Norris, J.G., S.Y. Hyun, and J.J. Anderson, 2000. Ocean distribution of Columbia River upriver bright fall chinook Salmon stocks. *North Pacific Anadromous Fish Commission Bulletin* 2:221-232.
- NMFS. 2019. National Marine Fisheries Service Section 10(a)(1)(B) permit 23194 to Public Utility District #2 of Grant County for takes of endangered/threatened species. Portland, Oregon.
- Ohlberger, J., E.J. Ward, D.E. Schnindler, and B. Lewis, 2018. Demographic changes in Chinook Salmon across the Northeast Pacific Ocean. *Fish and Fisheries*, 2018; DOL: 10.1111/faf.12272.
- Oldenburg, E.W., B.J. Goodman, G.A. McMichael, and R.B. Langshaw. 2012. Forms of Production Loss During the Early Life History of Fall Chinook Salmon in the Hanford Reach of the Columbia River. Prepared for the Public Utility District No. 2 of Grant County, Ephrata, WA. Contract Number 430-2464.
- Pearsons, T. N. and C. A. Busack. 2012. PCD Risk 1: A tool for assessing and reducing ecological risks of hatchery operations in freshwater. *Environmental Biology of Fishes* 94:45-65. DOI:10.1007/s10641-011-9926-8.
- Pearsons, T. N., A. R. Murdoch, G. Mackey, K. G. Murdoch, T. W. Hillman, M. R. Cooper, and J. L. Miller. 2012. Ecological risk assessment of multiple hatchery programs in the upper Columbia watershed using Delphi and modeling approaches. *Environmental Biology of Fishes* 94:87-100. DOI 10.1007/s10641-011-9884-1.
- Pearsons, T. N., and C. W. Hopley. 1999. A practical approach for assessing ecological risks associated with fish stocking programs. *Fisheries* 24(9):16-23.
- Pearsons, T. N., and R. B. Langshaw. 2009. Monitoring and evaluation plan for Grant PUDs Salmon and steelhead supplementation Programs. Grant PUD, Ephrata, Washington.
- Pearsons, T. N. and R. R. O'Connor. 2020. Stray rates of natural-origin Chinook Salmon and Steelhead in the Upper Columbia Watershed. *Transactions of the American Fisheries Society* 149:147–158. DOI: 10.1002/tafs.10220

- Pearsons, T. N., A. H. Haukenes, P. A. Hoffarth, and S. P. Richards. 2020. Expanding partnerships and innovations to implement reform of a large Columbia River hatchery program. *Fisheries* 45(9):484-491. DOI: 10.1002/fsh.10437
- PSC (Pacific Salmon Commission). 2013. 2013 Exploitation Rate Analysis and Model Calibration - Volume One. A report of the Pacific Salmon Commission Joint Chinook Technical Committee. Technical Report (14)-1 V.1.
- Hoffarth, P. A., and T. N. Pearsons. 2012a. Priest Rapids Hatchery Monitoring and Evaluation Annual Report for 2012-13. Public Utility District Number 2 of Grant County, Ephrata, Washington.
- Hoffarth, P. A., and T. N. Pearsons. 2012b. Priest Rapids Hatchery Monitoring and Evaluation Annual Report for 2012-13. Public Utility District Number 2 of Grant County, Ephrata, Washington.
- Richards, S. P., P. A. Hoffarth, and T. N. Pearsons. 2013. Priest Rapids Hatchery Monitoring and Evaluation Annual Report for 2012-13. Public Utility District Number 2 of Grant County, Ephrata, Washington.
- Richards, S. P., and T. N. Pearsons. 2014. Priest Rapids Hatchery Monitoring and Evaluation Annual Report for 2013-14. Public Utility District Number 2 of Grant County, Ephrata, Washington.
- Richards, S. P., and T. N. Pearsons. 2015. Priest Rapids Hatchery Monitoring and Evaluation Annual Report for 2014-15. Public Utility District Number 2 of Grant County, Ephrata, Washington.
- Richards, S. P., and T. N. Pearsons. 2016. Priest Rapids Hatchery Monitoring and Evaluation Annual Report for 2015-16. Public Utility District Number 2 of Grant County, Ephrata, Washington.
- Richards, S. P., and T. N. Pearsons. 2017. Priest Rapids Hatchery Monitoring and Evaluation Annual Report for 2016-17. Public Utility District Number 2 of Grant County, Ephrata, Washington.
- Richards, S. P., and T. N. Pearsons. 2018. Priest Rapids Hatchery Monitoring and Evaluation Annual Report for 2017-18. Public Utility District Number 2 of Grant County, Ephrata, Washington.
- Richards, S. P., and T. N. Pearsons. 2019. Priest Rapids Hatchery Monitoring and Evaluation Annual Report for 2018-19. Public Utility District Number 2 of Grant County, Ephrata, Washington.
- Richards, S. P., and T. N. Pearsons. 2020. Priest Rapids Hatchery Monitoring and Evaluation Annual Report for 2019-20. Public Utility District Number 2 of Grant County, Ephrata, Washington.
- Volk E.C., S.L. Schroder, and J.G. Grimm. 1999. Otolith Thermal Marking. *Fisheries Research* 43 (1999), pp. 205-219
- USDOE – U.S. Department of Energy. In Press. Hanford Site Ecological Monitoring Report for Calendar Year 2016. HNF-61231, Rev. 0. U.S. Department of Energy, Richland Operations Office, Richland, Washington.

Weitkamp, L.A. 2010: Marine Distributions of Chinook Salmon from the West Coast of North America Determined by Coded Wire Tag Recoveries, Transactions of the American Fisheries Society, 139 (1), pp. 147-170

Zhou, S., 2002. Size-dependent recovery of Chinook Salmon in carcass surveys. Transactions of the American Fisheries Society, 131(6), pp. 1194-1202.

Appendix A Evaluation of Coded-Wire Tag Bias

We annually evaluate bias associated with estimates of the number of hatchery origin returns to PRH generated using coded-wire tags (CWT). Results from demographic sampling of the fall Chinook returns for 2010 through 2014 indicate that estimates of hatchery contributions to broodstock, the terminal sport fishery, and to escapement of the Hanford Reach and to the PRH trap calculated from otolith marks were substantially different from estimates generated using CWTs expanded by sampling rates and juvenile mark rates. This was of significant concern because many estimates such as stray rate, survival, origin, and harvest are dependent upon estimates generated from CWTs.

To assess the level of CWT recovery bias for any brood year, we used the following equation:

$$\text{CWT Recovery Bias} = \frac{(\# \text{ of PRH Origin CWT Fish Recovered} / \# \text{ of PRH Origin Fish Collected})}{\text{CWT Mark Rate for Brood Year}}$$

Where:

of PRH origin fish collected = Estimate of the number of PRH origin fish for a specific age/brood year as determined by otoliths, scale aging, and expansion and pooling of age samples to represent total returns by age

of PRH Origin CWT Fish Recovered = Number of PRH origin CWT fish for a specific age/brood recovered at the hatchery (100% sample rate)

CWT Mark Rate = CWT marking rate for the specific brood year which is the number of CWT placed in fish divided by the estimated total number of fish at the time of marking.

If no CWT bias exists, the proportion of PRH CWT returns to the PRH CWT mark rate should equal 1.000. The values for CWT Recovery bias ranged from 0.258 to 4.851 for the different age/broods examined (Table A.1). Even though the datasets are not complete for recent brood years, it appears that the CWT Recovery bias is less pronounced since brood year 2011. The source of any bias is likely due to inappropriate expansion rate estimates resulting from non-representative placement of CWT groups within the general population of rearing in the channel ponds. However, several other factors may contribute to the variation in CWT Recovery bias such as tag loss, CWT detection efficiency, or differential survival of tagged fish. In addition, the estimate of bias may be influenced by the level of precision of the estimated # of PRH origin fish collected which varies for each age class of a given brood year due to size of the otolith sub-sample pulled from the demographic sample. In some cases, there are relatively few samples for age-2 and 5 fish for a given brood year for this estimate.

Verification of the juvenile CWT rate at time of release is necessary to determine level of potential bias associated with reported juvenile CWT rates. Sampling for CWT rates at time of release has occurred at PRH since brood year 2014. Shortly prior to release, roughly 1,000 subyearlings from each of the five rearing ponds were captured and scanned with a V-detector to determine the proportions of adipose clipped CWT fish and adipose intact CWT fish within the sample. These proportions at release were compared to the proportions reported as ponded. In general, these two groups of proportions were generally similar for each brood year with the exception of brood years 2018 and 2019 (Table A.2). The results of the quality control sampling for the 2018 brood year found that the observed CWT rates were higher than the expected rates for rearing ponds D and E; whereas during 2019 the observed CWT rates were notably lower for

Pond D. The observed CWT rates were notably higher than the expected rates for Pond E and similar for the other ponds.

Table A.1 Estimate of coded-wire tags bias for Priest Rapids origin returns to the hatchery, Brood Years 2007- 2017.

Brood	Age	Proportion CWT Marked	# of PRH Origin CWT Fish Recovered	Estimated # of PRH origin Fish Collected	Proportion of PRH Origin Brood Return CWT	Proportion of PRH CWT Returns to the PRH CWT Mark Rate (CWT Recovery Bias)	Primary Detector Type	
2007	5	0.045	48	928	0.052	1.161	Blue Wand	
2007	4	0.045	280	10,977	0.026	0.573	Blue Wand	
2007	3	0.045	410	14,073	0.029	0.654	Blue Wand	
2007	2	No otolith data collected during return year 2009						
2008	5	0.032	2	31	0.065	2.026	Blue Wand	
2008	4	0.032	81	3,029	0.027	0.840	Blue Wand	
2008	3	0.032	124	5,606	0.022	0.695	Blue Wand	
2008	2	0.032	57	2,578	0.022	0.694	Blue Wand	
2009	5	0.243	407	1,980	0.206	0.846	R9500	
2009	4	0.243	1,081	6,025	0.179	0.739	Blue Wand	
2009	3	0.243	2,309	13,713	0.168	0.693	Blue Wand	
2009	2	0.243	628	3,083	0.204	0.839	Blue Wand	
2010	6	0.237	23	20	1.150	4.851	R9500	
2010	5	0.237	999	2,375	0.421	1.778	R9500	
2010	4	0.237	8,719	39,621	0.220	0.928	R9500	
2010	3	0.237	5,828	32,014	0.182	0.768	Blue Wand	
2010	2	0.237	1,498	8,932	0.168	0.707	Blue Wand	
2011	6	0.169	10	47	0.213	0.258	R9500	
2011	5	0.169	395	2,520	0.157	0.927	R9500	
2011	4	0.169	2,988	19,536	0.153	0.904	R9500	
2011	3	0.169	2,596	19,692	0.132	0.779	R9500	
2011	2	0.169	349	3,008	0.116	0.686	R9500	
2012	6	0.177	7	19	0.368	2.086	R9500	
2012	5	0.177	1,913	11,259	0.170	0.961	R9500	
2012	4	0.177	2,206	13,821	0.160	0.904	R9500	
2012	3	0.177	5,933	34,082	0.174	0.986	R9500	
2012	2	0.177	1,910	11,259	0.170	0.961	R9500	
2013	6	0.166	0	0	0	N/A	R9500	
2013	5	0.166	109	527	0.207	1.245	R9500	
2013	4	0.166	1,530	8,695	0.164	0.998	R9500	
2013	3	0.166	1,805	10,967	0.165	0.991	R9500	
2013	2	0.166	545	3,327	0.164	0.986	R9500	
2014	6	0.172	0	0	0	N/A	R9500	
2014	5	0.172	15	189	0.080	0.462	R9500	
2014	4	0.172	407	2,685	0.152	0.883	R9500	
2014	3	0.172	483	3,289	0.147	0.856	R9500	
2014	2	0.172	78	462	0.169	0.984	R9500	
2015	5	0.167	71	290	0.245	1.473	R9500	
2015	4	0.167	1,158	7,081	0.164	0.979	R9500	

Brood	Age	Proportion CWT Marked	# of PRH Origin CWT Fish Recovered	Estimated # of PRH origin Fish Collected	Proportion of PRH Origin Brood Return CWT	Proportion of PRH CWT Returns to the PRH CWT Mark Rate (CWT Recovery Bias)	Primary Detector Type
2015	3	0.167	1,343	8,596	0.156	0.936	R9500
2015	2	0.167	183	1,219	0.150	0.899	R9500
2016	4	0.171	1,386	8,105	0.171	1.000	R9500
2016	3	0.171	1,152	7,879	0.146	0.855	R9500
2016	2	0.171	138	1,061	0.130	0.760	R9500
2017	3	0.151	3,161	22,778	0.139	0.919	R9500
2017	2	0.151	269	2,515	0.107	0.708	R9500
CWT Recovery Bias	Ages						
	Mean	2	3	4	5	6	
		0.822	0.830	0.875	1.209	2.398	

Table A.2. Proportions of coded-wire tagged juvenile fish reported ponded and the proportions of coded-wire tagged fish sampled at time of release, Brood Years 2014-20.

Coded-wire sampling at release, Brood Year 2014						
# of Fish	Pond E	Pond D	Pond C	Pond B	Pond A	Total
Fish Released	1,425,371	1,457,198	1,400,956	1,444,918	1,311,100	7,039,543
N =	1,040	1,024	1,018	1,023	1,565	5,670
CWT Only Sampled	98	85	79	67	220	549
Ad-CWT Sampled	102	69	73	86	165	495
Proportion of Release Tagged						
CWT Only	8.5%	8.3%	8.6%	8.2%	9.0%	8.5%
Ad-CWT	8.5%	8.2%	8.6%	8.7%	8.7%	8.5%
Proportion of Sample Tagged						
CWT Only	9.4%	8.3%	7.8%	6.5%	14.1%	9.7%
Ad-CWT	9.8%	6.7%	7.2%	8.4%	10.5%	8.7%
Coded-wire sampling at release, Brood Year 2015						
# of Fish	Pond E	Pond D	Pond C	Pond B	Pond A	Total
Fish Released	1,445,733	1,448,510	1,507,753	1,512,437	1,327,621	7,242,054
N =	1,015	995	991	1,048	1,021	5,070
CWT Only Sampled	91	86	77	62	76	392
Ad-CWT Sampled	71	87	79	71	80	388
Proportion of Release Tagged						
CWT Only	8.1%	8.6%	8.3%	7.5%	9.1%	8.3%
Ad-CWT	8.3%	8.6%	7.7%	8.0%	9.1%	8.3%
Proportion of Sample Tagged						
CWT Only	9.0%	8.6%	7.8%	5.9%	7.4%	7.7%
Ad-CWT	7.0%	8.7%	8.0%	6.8%	7.8%	7.7%

Table A.2 Continued

Coded-wire sampling at release, Brood Year 2016						
# of Fish	Pond E	Pond D	Pond C	Pond B	Pond A	Total
Fish Released	1,401,157	1,455,960	1,450,776	1,487,339	1,211,019	7,006,251
N =	1,031	1,317	2,228	1,117	1,181	6,874
CWT Only Sampled	119	103	205	116	120	663
Ad-CWT Sampled	101	96	224	112	117	650
Proportion of Release Tagged						
CWT Only	8.6%	8.3%	8.3%	8.1%	10.0%	8.6%
Ad-CWT	8.6%	8.3%	8.3%	8.1%	10.0%	8.6%
Proportion of Sample Tagged						
CWT Only	11.5%	7.8%	9.2%	10.4%	10.2%	9.6%
Ad-CWT	9.8%	7.3%	10.1%	10.0%	9.9%	9.5%
Coded-wire sampling at release, Brood Year 2017						
# of Fish	Pond E	Pond D	Pond C	Pond B	Pond A	Total
Fish Released	1,632,887	1,573,080	1,615,297	1,588,038	1,594,137	8,003,439
N =	1,046	1,260	1,022	1,173	1,044	5,545
CWT Only Sampled	88	143	74	87	85	477
Ad-CWT Sampled	81	164	71	77	67	460
Proportion of Release Tagged						
CWT Only	7.5%	7.6%	7.5%	7.6%	7.6%	7.6%
Ad-CWT	7.2%	7.7%	7.5%	7.6%	7.6%	7.5%
Proportion of Sample Tagged						
CWT Only	8.4%	11.3%	7.2%	7.4%	8.1%	8.6%
Ad-CWT	7.7%	13.0%	6.9%	6.6%	6.4%	8.3%
Coded-wire sampling at release, Brood Year 2018						
# of Fish	Pond E	Pond D	Pond C	Pond B	Pond A	Total
Fish Released	1,471,868	1,452,947	1,430,194	1,464,134	1,394,773	7,213,916
N =	1,201	1,197	1,099	1,100	1,100	5,697
CWT Only Sampled	192	136	99	83	92	602
Ad-CWT Sampled	200	158	99	109	91	657
Proportion of Release Tagged						
CWT Only	8.2%	8.2%	8.4%	8.3%	8.7%	8.3%
Ad-CWT	8.2%	8.2%	8.5%	8.3%	8.7%	8.4%
Proportion of Sample Tagged						
CWT Only	16.0%	11.4%	9.0%	7.5%	8.4%	10.6%
Ad-CWT	16.7%	13.2%	9.0%	9.9%	8.4%	11.5%
Coded-wire sampling at release, Brood Year 2019						
# of Fish	Pond E	Pond D	Pond C	Pond B	Pond A	Total
Fish Released	1,661,233	1,631,517	1,521,956	1,499,259	1,345,921	7,659,886
N =	1,070	1,093	1,145	1,212	1,704	6,224
CWT Only Sampled	75	50	94	98	75	602

Ad-CWT Sampled	86	54	74	83	96	657
Proportion of Release Tagged						
CWT Only	7.3%	7.4%	7.9%	8.0%	9.0%	8.3%
Ad-CWT	7.3%	7.4%	7.9%	8.0%	8.9%	8.4%
Proportion of Sample Tagged						
CWT Only	7.0%	4.6%	8.2%	8.1%	7.0%	10.6%
Ad-CWT	8.0%	4.6%	6.5%	6.8%	8.9%	11.5%
Coded-wire sampling at release, Brood Year 2020						
# of Fish	Pond E	Pond D	Pond C	Pond B	Pond A	Total
Fish Released	1,534,299	1,495,653	1,567,246	1,507,210	1,438,110	7,542,518
N =	1,218	1,183	1,040	1,038	1,038	5,517
CWT Only Sampled	115	92	72	82	82	443
Ad-CWT Sampled	112	99	78	90	90	469
Proportion of Release Tagged						
CWT Only	7.8%	8.3%	7.6%	8.0%	8.3%	8.0%
Ad-CWT	7.8%	8.1%	7.7%	7.9%	8.3%	8.0%
Proportion of Sample Tagged						
CWT Only	9.4%	7.8%	6.9%	7.9%	7.9%	8.0%
Ad-CWT	9.2%	8.4%	7.5%	8.7%	8.7%	8.5%

Assessment of CWT detection efficiency has been conducted annually at PRH since 2010 during adult fish sampling with enhancement to these procedures developed over time. In 2013, M&E staff randomly selected a total of 1,063 quality control fish being surplus with no CWT detected using the T-wand (Table A.3). These fish were then re-scanned with the older blue-wand. If CWT was detected using a blue wand the fish was again scanned using the T-wand. In such a manner the missed CWT could be inferred as a result of operator error or the inability of the T-wand to detect the CWT. On a few occasions the T-wand did not detect a CWT identified by the blue-wand. In these instances, the snouts were removed from the fish to increase the likelihood of detection and then passed through a V-detector. Similar to quality control results for previous years, there were only a few (2 tags) of the sample CWT detections observed in the quality control fish sampled that were not detected initially by the T-wands during 2020.

Table A.3 Quality control results for coded-wire tag detection at Priest Rapids Hatchery, Brood Years 2013- 2019.

Brood Year	Initial Device	QC Device	# Sampled	# Missed CWT
2013	T-Wand	Blue Wand	1,063	4
2014	R9500	T-Wand	2,000	3
2015	R9500	T-Wand	4,596	2
2016	R9500	T-Wand	5,943	3
2017	R9500	T-Wand	1,744	3
2018	R9500	T-wand	1,679	6
2019	R9500	T-wand	2,011	2
2020	R9500	T-wand	1886	2

During 2013 and 2014, we found the T-wands to be overly sensitive leading to false positive detections and additional work related to processing snouts to extract CWTs. On October 2, 2014 we set up two series R9500 detectors to expedite scanning for CWTs (Figure 1). The detectors were checked for proper operation each day prior to scanning any fish. Informal quality control checks occurred daily during the first two weeks of operation in order to identify the detection efficiency of each detector. These checks involved running 100 fish through each machine and then re-scanning the fish with the T-wands. A total of 2,000 fish were passed through the R9500 units of which 422 were identified to possess a CWT. Of these fish, 419 signaled positive for a CWT during the initial scanning. The three fish possessing a CWT that were not identified by the R9500 during the initial scanning were correctly detected when re-ran through the detectors. The missed fish were likely the result of passing fish through the detectors too rapidly which can interfere with the operation of the flip gates.

R9500 detectors were used to scan the vast majority of fish surplused at PRH during 2015 through 2019. During each of these years, the first group of fish handled each day was used to test the CWT detection of each R9500 detector. The test fish that a CWT was not detected were re-scanned with a T-wand to assess the performance of the R9500 detectors. The results for all three years suggest that very few possessing a CWT are missed by the R9500 detectors.

The methods describe here do not provide a definitive estimate of undetected CWTs for fish sampled at PRH. We make an assumption, that if the CWT detection wands and R9500 units do not detect a CWT in a given fish, then it did not possess a tag. Based on this assumption, the CWT detection efficiency at PRH is likely greater than 99%. Therefore, the magnitude of the CWT recovery bias expressed in Table 1 is not likely due to poor CWT detection efficiency.

Appendix B
Recovery of coded-wire tags collected from adult returns to the Priest Rapids
Hatchery Volunteer Trap during Return Year 2020

Code	Tag #	BY	Race	Age	Stock	Release Year
56163	1	2019	Fall	1	Spring Cr	2020
637730	1	2019	Fall	1	Priest Rapids	2020
91177	8	2018	Fall	2	Umatilla River	2019
91286	4	2018	fall	2	Snake River	2019
91377	5	2018	Fall	2	Priest Rapids	2019
200146	1	2018	Summer	2	Okanogan River	2020
220267	1	2018	Fall	2	Snake River	2019
220509	1	2018	Fall	2	Lyons Ferry	2019
220512	2	2018	Fall	2	Lyons Ferry	2019
220514	1	2018	Fall	2	Lyons Ferry	2019
637354	1	2018	Fall	2	Skokomish River	2019
637381	1	2018	Fall	2	Cowlitz Hatchery	2019
637420	2	2018	Fall	2	Snake River	2019
637467	1	2018	Spring	2	Chiwawa River	2020
637512	22	2018	Fall	2	Priest Rapids	2019
637513	53	2018	Fall	2	Priest Rapids	2019
637514	57	2018	Fall	2	Priest Rapids	2019
637515	52	2018	Fall	2	Priest Rapids	2019
637516	71	2018	Fall	2	Priest Rapids	2019
637517	29	2018	Fall	2	Priest Rapids	2019
637520	55	2018	Fall	2	Priest Rapids	2019
637521	63	2018	Fall	2	Priest Rapids	2019
637522	58	2018	Fall	2	Priest Rapids	2019
637523	69	2018	Fall	2	Priest Rapids	2019
637524	1	2018	Fall	2	Skokomish River	2019
55928	1	2017	Fall	3	Spring Cr	2018
91183	32	2017	Fall	3	Col R Upriver brights	2018
91184	41	2017	Fall	3	Col R Upriver brights	2018
91185	7	2017	Fall	3	Snake River	2018
91254	129	2017	Fall	3	Priest Rapids	2018
91255	59	2017	Fall	3	Priest Rapids	2018
91277	1	2017	Fall	3	WA Brights	2019
91280	1	2017	Fall	3	WA Brights	2019
220258	1	2017	Fall	3	Lyons Ferry	2018
220268	2	2017	Fall	3	Lyons Ferry	2018
220271	3	2017	Fall	3	Lyons Ferry	2018
220395	1	2017	Fall	3	Snake River	2019
220502	4	2017	Fall	3	Lyons Ferry	2018
220503	4	2017	Fall	3	Lyons Ferry	2018
220506	2	2017	Fall	3	Lyons Ferry	2018
220507	1	2017	Fall	3	Lyons Ferry	2018
610485	1	2017	Fall	3	Hanford URB Wild	2018
610487	5	2017	Fall	3	Hanford URB Wild	2018
610488	3	2017	Fall	3	Hanford URB Wild	2018
637240	2	2017	Fall	3	Little White Salmon	2018
637355	463	2017	Fall	3	Priest Rapids	2018
637356	370	2017	Fall	3	Priest Rapids	2018
637357	312	2017	Fall	3	Priest Rapids	2018
637358	330	2017	Fall	3	Priest Rapids	2018
637359	447	2017	Fall	3	Priest Rapids	2018
637360	276	2017	Fall	3	Priest Rapids	2018

Code	Tag #	BY	Race	Age	Stock	Release Year
637361	330	2017	Fall	3	Priest Rapids	2018
637362	288	2017	Fall	3	Priest Rapids	2018
637363	167	2017	Fall	3	Priest Rapids	2018
637364	178	2017	Fall	3	Priest Rapids	2018
637385	1	2017	Spring	3	Lewis River Hatchery	2019
637387	2	2017	Spring	3	Lewis River Hatchery	2019
637394	1	2017	Fall	3	Snake River	2018
637395	6	2017	Fall	3	Snake River	2018
637397	4	2017	Fall	3	Snake River	2019
637398	2	2017	Fall	3	Snake River	2019
91011	12	2016	Fall	4	WA Brights	2017
91084	14	2016	Fall	4	WA Brights	2017
91112	18	2016	Fall	4	Priest Rapids	2017
91113	40	2016	Fall	4	Priest Rapids	2017
91134	1	2016	Fall	4	WA Brights	2017
91179	3	2016	Fall	4	Yakima River	2018
200144	1	2016	Summer	4	Okanogan River	2018
200145	1	2016	Summer	4	Okanogan River	2018
220252	1	2016	Fall	4	Snake River	2017
220253	2	2016	Fall	4	Snake River	2017
220259	1	2016	Fall	4	Snake River	2017
220260	1	2016	Fall	4	Snake River	2017
220261	1	2016	Fall	4	Snake River	2017
220262	1	2016	Fall	4	Snake River	2017
220384	1	2016	Fall	4	Lyons Ferry	2017
220388	1	2016	Fall	4	Lyons Ferry	2018
610476	2	2016	Fall	4	Hanford URB Wild	2017
610478	2	2016	Fall	4	Hanford URB Wild	2017
610479	2	2016	Fall	4	Hanford URB Wild	2017
610480	2	2016	Fall	4	Hanford URB Wild	2017
610483	1	2016	Fall	4	Hanford URB Wild	2017
610484	1	2016	Fall	4	Hanford URB Wild	2017
636986	20	2016	Fall	4	Priest Rapids	2017
637123	1	2016	Spring	4	Lewis River Hatchery	2017
637148	139	2016	Fall	4	Priest Rapids	2017
637179	114	2016	Fall	4	Priest Rapids	2017
637180	140	2016	Fall	4	Priest Rapids	2017
637181	149	2016	Fall	4	Priest Rapids	2017
637182	120	2016	Fall	4	Priest Rapids	2017
637183	114	2016	Fall	4	Priest Rapids	2017
637184	140	2016	Fall	4	Priest Rapids	2017
637185	123	2016	Fall	4	Priest Rapids	2017
637186	213	2016	Fall	4	Priest Rapids	2017
637187	114	2016	Fall	4	Priest Rapids	2017
637194	1	2016	Fall	4	Little White Salmon	2017
637199	1	2016	Fall	4	Snake River	2017
91083	2	2015	Fall	5	Bonneville Hatchery	2017
636967	50	2015	Fall	5	Priest Rapids	2016
636968	21	2015	Fall	5	Priest Rapids	2016
90909	71	Blank wire			Oregon Blank wire	
Total	5,679					

Appendix C

Juvenile fish health inspections for Priest Rapids Hatchery fall Chinook Salmon, Brood Years 1998-2020. The description in the Condition column indicates the presence of a certain condition within at least one of the fish examined.

Hatchery/Stock	Date	Brood	Condition
Priest Rapids	23-Feb-99	1998	Healthy
	22-Mar-99	1998	Healthy
	23-Apr-99	1998	Healthy
	25-May-99	1998	Dropout Syndrome & Bacterial Gill Disease
	08-Jun-99	1998	Bacterial Kidney Disease
Priest Rapids	06-Mar-00	1999	Healthy
	14-Apr-00	1999	Healthy
	16-May-00	1999	Healthy
	12-Jun-00	1999	Healthy
Priest Rapids	23-Feb-01	2000	Healthy
	05-Apr-01	2000	Healthy
	07-May-01	2000	Healthy
	06-Jun-01	2000	Healthy
Priest Rapids	13-Feb-02	2001	Healthy
	01-Mar-02	2001	Coagulated Yolk Syndrome
	22-Apr-02	2001	Healthy
	10-Jun-02	2001	Healthy
Priest Rapids	07-Mar-03	2002	Healthy
	15-Apr-03	2002	Healthy
	02-Jun-03	2002	Healthy
Priest Rapids	01-Apr-04	2003	Healthy
	06-May-04	2003	Healthy
	07-Jun-04	2003	Healthy
Priest Rapids	11-Mar-05	2004	Healthy
	14-Apr-05	2004	Healthy
	1-Jun-05	2004	Healthy
Priest Rapids	6-Mar-06	2005	Healthy
	25-Apr-06	2005	Healthy
	13-Jun-06	2005	Healthy
Priest Rapids	9-Mar-07	2006	Healthy
	19-Apr-07	2006	Healthy
	1-Jun-07	2006	Healthy
Priest Rapids	12-Feb-08	2007	Coagulated Yolk Syndrome observed in some fish sampled
	23-Apr-08	2007	Healthy
	4-Jun-08	2007	Healthy
Priest Rapids	12-Feb-09	2008	Coagulated Yolk Syndrome observed in some fish sampled
	22-Apr-09	2008	Healthy
	8-Jun-09	2008	Healthy
Priest Rapids	18-Feb-10	2009	Coagulated Yolk Syndrome observed in some fish sampled
	1-Apr-10	2009	Healthy
	19-May-10	2009	Healthy
Priest Rapids	25-Mar-11	2010	Healthy
	18-Apr-11	2010	Healthy
	06-Jun-11	2010	Healthy
Priest Rapids	01-Mar-12	2011	Healthy
	26-Apr-12	2011	Healthy
	24-May-12	2011	Healthy
Priest Rapids	11-Feb-13	2012	Healthy
	3-Mar-13	2012	Healthy

Hatchery/Stock	Date	Brood	Condition
	29-Apr-13	2012	Healthy
	28-May-13	2012	Healthy
Priest Rapids	27-Mar-14	2013	Dropout Syndrome present
	23-Apr-14	2013	Dropout Syndrome present
	29-May-14	2013	Healthy
Priest Rapids	26-Feb-15	2014	Coagulated Yolk Syndrome observed in some fish sampled
	26-Mar-15	2014	Healthy
	21-Apr-15	2014	Healthy
	28-May-15	2014	Healthy
	22-June-15	2014	Columnaris present in some fish sampled from CH Pond B.
Priest Rapids	24-Feb-16	2015	Healthy
	15-Mar-16	2015	Coagulated Yolk Syndrome observed in some fish sampled
	15-June-16	2015	Mild Ich infection but healthy and ready for release
Priest Rapids	24-Feb-17	2016	Presence of bacterial gill disease in Raceway Bank D and E
	21-Mar-17	2016	Presence of bacterial gill disease in Raceway Pond B2
	6-June-17	2016	Mild Ich infection in Channel Ponds A, B, C
Priest Rapids	21-Mar-18	2017	Healthy
	19-Apr-18	2017	Bacterial gill diseases present in Raceway Pond C4
	7-May-18	2017	Bacterial gill diseases present in Raceway Ponds C2 and C3
	17-May-18	2017	Re-examine Raceway Ponds C2 and C3 found fish healthy
	17-May-18	2017	Pre-release examine Raceway Banks D and E found fish healthy C2 and C3 found fish healthy
	6-June-18	2017	Pre-release examine of Raceway Banks A and B found fish healthy
Priest Rapids	2-Feb-19	2018	Examines of Raceway Banks C, D, E resulted from reports of elevated mortalities. Some fish were found to appear thin and pin-headed. Results of internal necropsies were within normal limits.
	5-May-19	2018	Pre-release examine of Raceway Pond E found fish healthy
	5-May-19	2018	Pre-release examine of Raceway Pond D resulted no significant findings of diseases however elevated mortalities were observed. Mortalities examined showed lower levels of coelomic fat and ingesta in GI tracts compared to live fish examined.
	6-June-19	2018	Pre-release examines of Raceway Ponds A, B, and C found very low levels of bacterial gill disease
	2-Feb-19	2018	Examines of Raceway Banks C, D, E resulted from reports of elevated mortalities. Some fish were found to appear thin and pin-headed. Results of internal necropsies were within normal limits.
	5-May-19	2018	Pre-release examine of Raceway Pond E found fish healthy
Priest Rapids	19-Mar-20	2019	Examinations of Raceway Bank A resulted from reports of flashing with minimal increase in observed mortality. Some fish examined revealed moderate infestation of Trichodina sp.
	28-Apr-20	2019	Examinations of Raceway Banks A, B, C, D, E resulted from reports of elevated mortalities. Some fish were found to appear thin and pin-headed. Results of internal necropsies were within normal limits.
	20-May-20	2019	Pre-release examine Raceway Ponds D and E found generally fish healthy.
	4-Jun-20	2019	Pre-release examine Raceway Ponds A, B, and C found generally fish healthy.
Priest Rapids	21-May-21	2020	Pre-release examination of Raceway Ponds D and E found fish generally healthy.
	7-June-21	2020	Pre-release examination of Raceway Ponds A, B, and C found fish generally healthy.

Appendix D

Number and percent of fall Chinook Salmon redds counted in different reaches of the Columbia River, 2001-2020. Data for years 2001-2010 was collected by staff with Pacific Northwest National Laboratory. Data for years 2001-2020 was collected by staff with Environmental Assessment Services, LLC.

Location	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Islands 11-21	297	509	554	337	708	36	302	371	176	562
Islands 8-10	480	865	1,133	867	1,067	435	338	416	722	870
Near Island 7	350	280	455	415	500	873	311	360	380	457
Island 6 (lower)	750	940	1,241	1,084	1,229	289	615	753	878	1,135
Island 4, 5,6	1,130	1,165	1,242	1,655	1,130	934	655	960	796	1,562
Near Island 3	460	249	475	325	345	1,305	152	230	285	244
Near Island 2	780	955	850	960	895	523	455	555	459	657
Near Island 1	35	235	270	330	255	253	47	148	160	324
Coyote Rapids	16	63	354	180	304	150	10	29	34	49
China Bar	20	25	85	75	28	52	3	35	1,090	299
Vernita Bar	1,930	2,755	2,806	2,240	1,430	1,658	1,135	1,731	16	2,658
Total	6,248	8,041	9,465	8,468	7,891	6,508	4,023	5,588	4,996	8,817
Location	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Islands 11-21	5%	6%	6%	4%	9%	1%	8%	7%	4%	6%
Islands 8-10	8%	11%	12%	10%	14%	7%	8%	7%	14%	10%
Near Island 7	6%	3%	5%	5%	6%	13%	8%	6%	8%	5%
Island 6 (lower)	12%	12%	13%	13%	16%	4%	15%	13%	18%	13%
Island 4, 5, 6	18%	14%	13%	20%	14%	14%	16%	17%	16%	18%
Near Island 3	7%	3%	5%	4%	4%	20%	4%	4%	6%	3%
Near Island 2	12%	12%	9%	11%	11%	8%	11%	10%	9%	7%
Near Island 1	1%	3%	3%	4%	3%	4%	1%	3%	3%	4%
Coyote Rapids	<1%	1%	4%	2%	4%	2%	<1%	1%	1%	1%
China Bar	<1%	<1%	1%	1%	<1%	1%	<1%	1%	<1%	3%
Vernita Bar	31%	34%	30%	26%	18%	25%	28%	31%	22%	30%
Location	2011	2012	2013	2014	2015	2016	2017	2018	2019	(10-19) Mean
Islands 11-21	676	533	798	906	1,193	861	280	88	0	590
Islands 8-10	814	807	2,200	1,565	3,145	1,735	900	485	166	1,269
Near Island 7	670	700	655	1,100	800	670	670	350	723	680
Island 6 (lower)	1,181	1,375	3,340	2,530	2,315	1,807	900	950	408	1,594
Island 4, 5,6	1,524	1,195	2,650	2,080	2,540	2,270	911	605	810	1,615
Near Island 3	525	475	1,000	1,000	1,100	600	500	310	939	669
Near Island 2	653	528	1,700	2,050	1,900	1,140	790	550	300	1,027
Near Island 1	295	340	900	500	1,000	340	330	170	720	492
Coyote Rapids	44	29	520	500	765	255	80	51	150	244
China Bar	67	68	100	60	1,730	80	75	25	112	262
Vernita Bar	2,466	2,318	3,535	3,650	4,190	3,510	3210	1,845	3,541	3,092
Total	8,915	8,368	17,398	15,951	20,678	13,268	8,646	5,429	7,869	11,533
Location	2011	2012	2013	2014	2015	2016	2017	2018	2019	(10-19) Mean
Islands 11-21	8%	6%	5%	6%	6%	6%	3%	2%	0%	5%
Islands 8-10	9%	10%	13%	10%	15%	13%	10%	9%	2%	11%
Near Island 7	8%	8%	4%	7%	4%	5%	8%	6%	9%	6%
Island 6 (lower)	13%	16%	19%	16%	11%	14%	10%	17%	5%	14%
Island 4, 5, 6	17%	14%	15%	13%	12%	17%	11%	11%	10%	14%
Near Island 3	6%	6%	6%	6%	5%	5%	6%	6%	12%	6%
Near Island 2	7%	6%	10%	13%	9%	9%	9%	10%	4%	9%
Near Island 1	3%	4%	5%	3%	5%	3%	4%	3%	9%	4%
Coyote Rapids	<1%	<1%	3%	3%	4%	2%	1%	1%	2%	2%
China Bar	1%	1%	1%	0%	8%	1%	1%	<1%	1%	2%
Vernita Bar	28%	28%	20%	23%	20%	26%	37%	34%	45%	27%

Location	2020	2021	2022	2023	2024	2025	2026	2027	2028	(11-20) Mean
Islands 11-21	507									639
Islands 8-10	524									1,317
Near Island 7	650									738
Island 6 (lower)	1,310									1,685
Island 4, 5, 6	1,562									1,727
Near Island 3	800									771
Near Island 2	1,100									1,129
Near Island 1	100									503
Coyote Rapids	70									268
China Bar	20									257
Vernita Bar	3,507									3,325
Total	10,150									1,124
Location	2020	2021	2022	2023	2024	2025	2026	2027	2028	(11-20) Mean
Islands 11-21	5%									5%
Islands 8-10	5%									10%
Near Island 7	6%									7%
Island 6 (lower)	13%									13%
Island 4, 5, 6	15%									14%
Near Island 3	8%									7%
Near Island 2	11%									9%
Near Island 1	1%									4%
Coyote Rapids	1%									2%
China Bar	0%									1%
Vernita Bar	35%									30%

Appendix E

Historical numbers of Chinook Salmon carcasses recovered during the annual Hanford Reach fall Chinook Salmon carcass survey, Return Years 1991-2020.

Return Year	Total Recoveries	Total Escapement	Proportion of Escapement Recovered
1991	2,519	52,196	0.048
1992	2,221	41,952	0.053
1993	3,340	37,347	0.089
1994	5,739	63,103	0.091
1995	3,914	55,208	0.071
1996	4,529	43,249	0.105
1997	5,053	43,493	0.116
1998	4,456	35,393	0.126
1999	4,412	29,812	0.148
2000	10,556	48,020	0.220
2001	6,072	59,848	0.101
2002	8,402	84,509	0.099
2003	13,573	100,840	0.135
2004	11,030	87,696	0.126
2005	8,491	71,967	0.118
2006	5,972	51,701	0.116
2007	3,115	22,272	0.140
2008	5,455	29,058	0.188
2009	5,318	36,720	0.145
2010	9,779	87,016	0.112
2011	8,391	75,256	0.111
2012	6,814	57,710	0.118
2013	13,071	174,651	0.075
2014	16,756	183,749	0.091
2015	17,738	266,346	0.086
2016	8,886	116,421	0.076
2017	5,591	73,759	0.076
2018	2,771	46,624	0.059
2019	4,069	64,664	0.063
2020	4,669	74,834	0.062
Mean	7,090	73,847	0.105
Median	5,665	58,779	0.103

Appendix F
Estimated escapements for fall Chinook spawning in Hanford Reach and Priest Rapids Dam pool, Return Year 2020.

Count Source		2020 Hanford Reach Escapement		
		Adult	Jack	Total
Adult Fish Counts	McNary ¹	186,097	27,171	213,268
	Rock Island ⁸	13,877	1,236	15,113
	Wanapum ²	29,712	2,619	32,331
	Priest Rapids ³	46,966	4,426	51,392
	Adjustment ⁴	16,042	1,512	17,554
	Ice Harbor ⁵	29,172	9,251	38,423
	Prosser ⁶	2,061	118	2,179
Hatcheries	Priest Rapids Hatchery	34,839	4,109	38,948
	Priest Rapids Hatchery Channel	54	3	57
	Angler Broodstock Collection	1,166	2	1,168
	OLAFT Broodstock Collection	0	0	0
	Ringold Springs Hatchery	6,737	354	7,091
Harvest	Hanford Sport Harvest	14,651	1,395	16,046
	Yakima River Sport Harvest	230	55	285
	Wanapum Tribal (above PRD)	91	0	91
	Wanapum Tribal (below PRD)	12	0	12
Escapement	Yakima River (Lower) ⁷	473	31	504
	Hanford Reach + Priest Pool	66,899	9,234	76,133
	Priest Pool Return	1,121	295	1,416
	Priest + Wanapum Pool Return	16,956	1,678	18,634
	Hanford Reach Escapement	65,778	8,939	74,717

¹ McNary fish counts: August 9 - October 31

² Wanapum AFC: August 16 - November 7

³ Priest Rapids AFC: August 14 - November 5

⁴ Fallback (34.16) adjustment based on 217 run of the river (BOAFF) PIT tagged fish observed at PRD

⁵ Ice Harbor fish counts: August 12 -October 31

⁶ Prosser counts, August 14 through November 11 (SU+FA), missing counts Sept 8-13 estimated

⁷ Escapement estimated by proportion of spawning below Prosser v Prosser passage (2010-2018)

⁸ Rock Island AFC: August 18 - November 9

Count Source	2020 Priest Rapids Pool Escapement		
	Adult	Jack	Total
Priest Rapids Adult Passage ³	46,966	4,426	51,392
Priest Rapids Fallback Adjustment ²	16,042	1,512	17,554
Wanapum Adult Passage ¹	29,712	2,619	32,331
Wanapum Dam Fallback Adjustment	Unknown	Unknown	Unknown
Wanapum Tribal Fishery Above PRD	91	0	91
Priest Rapids Pool Sport Fishery	555	77	576
Priest Rapids Dam Pool Escapement	566	218	840

¹ Wanapum Dam fish counts, August 14 through November 5.

² Fallback/Reascension Adjustment estimate (34.15%) based on 68 run of the river PIT tagged fish from the BOAFF and the lower Columbia River test fishery observed at Priest Rapids Dam and Priest Rapids Hatchery PIT tag arrays.

³ Priest Rapids passage for fall Chinook based on counts from August 18 through November 15.

Appendix G

Demographic comparisons for double index tag groups released from Priest Rapids Hatchery, Brood Years 2009-2017.

Double Index Tag (DIT) groups of fall Chinook Salmon have been released annually from Priest Rapids Hatchery (PRH) starting with the progeny of the 2009 brood. Adipose clipped fish from these DIT groups have been recovered in various mark selective fisheries (MSF) occurring in ocean, marine, and freshwater zones. The Regional Mark Processing Center database was queried to identify mark selective fisheries occurring since 2012 that included recoveries of PRH DIT groups (Table G.1). Detailed descriptions of these fisheries are available at websites maintained by the RMPC, Oregon Department of Fish and Game, and WDFW. The level of contribution to these fisheries, some of which are summer Chinook Salmon fisheries, is beyond the scope of this document.

Survival estimates for DIT groups from release and recovery at PRH was calculated by dividing the total DIT recoveries at PRH for each brood year (ages 1 – 6) by the corresponding number of juveniles marked for each DIT group. Similarities in gender composition, survival, age at maturity, and size at age between DIT groups within a brood year strongly suggest there is no difference for fish recovered at PRH (Tables G.2, G.3, G.4, and G.5).

Table G.1 Regional Mark Processing Center mark selective fisheries showing recoveries of Priest Rapids Hatchery origin coded-wire tagged adipose clipped fall Chinook Salmon from brood years 2009-2017.

Sampling Agency	Fishery
Alaska Dept. of Fish and Game	Ocean Selective Troll
Oregon Dept. of Fish and Game	Ocean Sport Columbia River Sport Columbia River Test Net Columbia River Purse Seine Columbia River Gillnet
Washington Dept. of Fish and Wildlife	Marine Sport Columbia River Sport

Table G.2 Gender Composition of DIT groups recovered at Priest Rapids Hatchery by brood year. Brood years 2015-2017 not complete. Data current through Return Year 2020.

Brood Year	Males		Females	
	Ad-CWT	CWT Only	Ad-CWT	CWT Only
2009	0.730	0.711	0.270	0.289
2010	0.546	0.542	0.454	0.458
2011	0.648	0.634	0.352	0.366
2012	0.643	0.640	0.357	0.360
2013	0.641	0.638	0.359	0.362
2014	0.636	0.603	0.364	0.397
2015	0.623	0.644	0.377	0.356
2016	0.627	0.603	0.373	0.397
2017	0.830	0.839	0.170	0.161
Mean	0.658	0.651	0.342	0.349

Table G.3 Smolt to adult return proportion comparisons between DIT Groups recovered at Priest Rapids Hatchery by brood year. Brood years 2015-2017 not complete. Data current through Return Year 2020.

Brood Year	Mark plus CWT	P^ Survival by Age					
		Age 2	Age 3	Age 4	Age 5	Age 6	Total
2009	Ad-Clipped	0.0004	0.0014	0.0006	0.0003	0.0000	0.0026
	No Mark	0.0004	0.0014	0.0007	0.0002	0.0000	0.0027
2010	Ad-Clipped	0.0009	0.0033	0.0051	0.0005	0.0000	0.0098
	No Mark	0.0009	0.0035	0.0051	0.0006	0.0000	0.0101
2011	Ad-Clipped	0.0003	0.0021	0.0024	0.0003	0.0000	0.0051
	No Mark	0.0003	0.0023	0.0026	0.0003	0.0000	0.0055
2012	Ad-Clipped	0.0015	0.0047	0.0018	0.0003	0.0000	0.0083
	No Mark	0.0017	0.0052	0.0019	0.0004	0.0000	0.0091
2013	Ad-Clipped	0.0005	0.0014	0.0012	0.0001	0.0000	0.0032
	No Mark	0.0004	0.0016	0.0013	0.0001	0.0000	0.0035
2014	Ad-Clipped	0.0001	0.0004	0.0004	0.0000	0.0000	0.0008
	No Mark	0.0001	0.0004	0.0003	0.0000	0.0000	0.0008
2015	Ad-Clipped	0.0001	0.0012	0.0010	0.0001	--	0.0024
	No Mark	0.0002	0.0011	0.0009	0.0000	--	0.0022
2016	Ad-Clipped	0.0001	0.0009	0.0011	--	--	0.0021
	No Mark	0.0001	0.0010	0.0013	--	--	0.0024
2017	Ad-Clipped	0.0001	0.0028	--	--	--	0.0029
	No Mark	0.0003	0.0025	--	--	--	0.0028
Mean	Ad-Clipped	0.0004	0.0020	0.0015	0.0002	0.0000	0.0041
	No Mark	0.0005	0.0021	0.0016	0.0002	0.0000	0.0043

Table G.4 Age composition of DIT Groups recovered at Priest Rapids Hatchery by brood year. Brood years 2015-2017 not complete. Data current through Return Year 2020.

Brood Year	Mark plus CWT	N	Age Composition (Genders Combined)				
			Age-2	Age-3	Age-4	Age-5	Age-6
2009	Ad-Clipped	1,635	0.138	0.522	0.244	0.096	0.001
	No Mark	2,795	0.144	0.521	0.245	0.089	0.000
2010	Ad-Clipped	5,914	0.088	0.337	0.518	0.056	0.002
	No Mark	11,198	0.087	0.347	0.505	0.060	0.001
2011	Ad-Clipped	3,034	0.053	0.412	0.465	0.069	0.002
	No Mark	3,306	0.057	0.408	0.477	0.057	0.002
2012	Ad-Clipped	4,990	0.184	0.565	0.212	0.038	0.001
	No Mark	5,472	0.182	0.569	0.210	0.039	0.000
2013	Ad-Clipped	1,917	0.147	0.436	0.382	0.035	0.000
	No Mark	2,084	0.127	0.465	0.388	0.020	0.000
2014	Ad-Clipped	508	0.079	0.484	0.419	0.018	0.000
	No Mark	480	0.079	0.498	0.410	0.013	0.000
2015	Ad-Clipped	1,442	0.062	0.481	0.422	0.035	--
	No Mark	1,324	0.071	0.498	0.415	0.016	--
2016	Ad-Clipped	1,248	0.056	0.423	0.521	--	--
	No Mark	1,453	0.047	0.431	0.522	--	--
2017	Ad-Clipped	1,722	0.042	0.958	--	--	--
	No Mark	1,725	0.114	0.886	--	--	--
Mean	Ad-Clipped	N/A	0.101	0.566	0.298	0.035	0.001
	No Mark	N/A	0.107	0.567	0.295	0.031	0.000

Table G.5 Size at age for DIT Groups recovered at Priest Rapids Hatchery by brood year. Brood years 2015-2017 not complete. Data current through Return Year 2020.

Brood Year	Mark plus CWT	Fall Chinook fork length (cm)														
		Age-2			Age-3			Age-4			Age-5			Age-6		
		N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD	N	Mean	SD
2009	Ad-Clipped	225	49	4	853	67	5	399	78	5	157	86	5	1	72	--
	No Mark	403	48	4	1,456	66	5	685	77	6	250	84	6	1	86	--
2010	Ad-Clipped	520	48	4	1,991	68	4	3,065	77	5	329	81	5	9	89	5
	No Mark	978	48	4	3,881	68	5	5,655	77	5	670	82	5	14	81	6
2011	Ad-Clipped	161	47	4	1,249	65	5	1,411	76	5	208	82	6	5	84	2
	No Mark	189	47	4	1,348	66	5	1,577	77	5	187	82	6	5	79	10
2012	Ad-Clipped	916	49	5	2,820	67	5	1,060	78	5	188	82	5	5	83	2
	No Mark	994	49	5	3,113	67	5	1,148	78	5	215	81	6	2	95	11
2013	Ad-Clipped	281	45	5	836	66	5	732	75	5	68	81	7	--	--	--
	No Mark	264	45	5	970	66	5	808	75	5	41	80	7	--	--	--
2014	Ad-Clipped	40	49	3	246	66	5	212	76	5	9	83	5	--	--	--
	No Mark	38	50	4	239	66	5	197	76	5	6	82	1	--	--	--
2015	Ad-Clipped	89	45	4	694	66	5	608	77	7	51	82	7	--	--	--
	No Mark	94	44	4	659	66	5	550	77	6	21	82	9	--	--	--
2016	Ad-Clipped	70	46	3	528	67	5	650	78	7						
	No Mark	67	46	4	626	68	5	759	79	6						
2017	Ad-Clipped	73	44	3	1649	66	4									
	No Mark	196	46	4	1528	66	5									
Mean	Ad-Clipped	264	47	4	1,207	66	5	1,017	77	5	144	82	6	5	82	3
	No Mark	358	47	4	1,536	67	5	1,422	77	5	199	82	6	6	85	9

Appendix H

Explanation of methods for calculating adult-to-adult expansions based on coded-wire tag recoveries at Priest Rapids Hatchery

Expanding adult coded wire tag (CWT) recoveries of either PRH or RSH origin fish by the corresponding brood's juvenile CWT rates has historically resulted in an underestimate of adult returns to locations within the Hanford Reach for each brood. A variety of factors may contribute to this problem; however, inappropriate juveniles tag expansion rates resulting from non-representative placement of tag groups within the general population is likely the greatest contributing factor. For many years, WDFW fish management staff have addressed the issues related to problematic juvenile tag rates by employing adult-to-adult CWT expansions for the PRH origin returns to PRH for run-reconstruction associated with their annual fall Chinook Salmon forecast. We used similar methods to expand PRH and RSH origin adult CWT recoveries in the vicinity of Hanford Reach to calculate PNI. An example of the calculations for the adult-to-adult expansion for the 2010 brood during return year 2014 is provided below. We make the assumption that the total number of PRH origin returns to PRH can be determined by removing other hatchery fish from the return: this is done by expanding the few other hatchery CWT recoveries by their corresponding juvenile CWT rates. Other hatchery CWT groups often have tag rates exceeding 50%; therefore, we assume juvenile tag rate expansions are representative for these groups. In addition, we make the assumption that very few natural origin fish return to PRH.

$$\text{Adult-to-Adult Expansion}_{\text{BY2010}} = \frac{\text{Total}_{\text{BY2010}} \text{ CWT Recoveries at PRH}}{\text{Total}_{\text{BY2010}} \text{ PRH Origin Returns to PRH}}$$

$$\text{Adult-to-Adult Expansion}_{\text{BY2010}} = \frac{8,719}{41,348} = 0.211$$

We then use the Adult-to-Adult Expansion_{BY2010} to expand all recoveries of PRH_{BY2010} in the Hanford Reach stream survey for return year 2014. This method is duplicated for each brood present in the given return year for both PRH and RSH to determine the total number of PRH and RSH origin fish in the escapement. The estimated number of PRH origin fish in the RY2014 Hanford Reach escapement based on the adult-to-adult expansion is higher than the number calculated using the conventional juvenile tag rate (Table 1).

Table I.1 The number of Priest Rapids Hatchery origin fish in the RY 2014 Hanford Reach escapement calculated from Adult-to-Adult Expansions versus Juvenile Tag Rates.

BY	CWT Recovered	Adult-to-Adult Exp	Expanded CWT	Survey Sample Rate	Total PRH origin in Escapement
2009	5	0.216	23	0.1063	218
2010	139	0.211	659	0.1063	6,197
2011	18	0.127	142	0.1063	1,333
2012	5	0.160	31	0.019	1,645
Adult-to-Adult Exp estimate for PRH origin fish in the Hanford Reach Escapement					9,393
Juvenile Tag Rate estimate for PRH origin fish in the Hanford Reach Escapement					7,934